

BS

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property
Organization
International Bureau



(43) International Publication Date
8 January 2004 (08.01.2004)

PCT

(10) International Publication Number
WO 2004/002481 A1

(51) International Patent Classification⁷: **A61K 31/433**,
A61P 3/10, 3/06, 3/04, C07D 277/48, 277/52, 213/75,
417/12, 277/56, 417/06, 417/14, 285/16, 487/04, 277/54

(21) International Application Number:
PCT/DK2003/000449

(22) International Filing Date: 27 June 2003 (27.06.2003)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
PA 2002 00999 27 June 2002 (27.06.2002) DK
60/394,144 3 July 2002 (03.07.2002) US
PA 2003 00286 25 February 2003 (25.02.2003) DK
60/452,228 5 March 2003 (05.03.2003) US

(71) Applicant (for all designated States except US): **NOVO
NORDISK A/S** [DK/DK]; Novo Allé, DK-2880 Bagsværd
(DK).

(72) Inventors; and

(75) Inventors/Applicants (for US only): **POLISETTI**,
Dharma, Rao [IN/US]; 4460 Platinum Drive, Apt #
1B, Greensboro, NC 27409 (US). **KODRA**, János,
Tibor [DK/DK]; Ryesgade 111B, 4, DK-2100 Copen-
hagen Ø (DK). **LAU**, Jesper [DK/DK]; Dybedalsvej
27, DK-3520 Farum (DK). **BLOCH**, Paw [DK/DK];
Solbærhøget 2, DK-2630 Taastrup (DK). **VAL-
CARCE-I-ÓPEZ**, Maria, Carmen, [ES/SE]; Strandgatan
100, S-21611 Limhamn (SE). **BLUME**, Niels [DK/DK];
Steenstrups Allé 11, 4th, DK-1924 Frederiksberg C (DK).
GUZEL, Mustafa [TR/US]; 4058 Gunsmith Court,
Jamestown, NC 27282 (US). **SANTHOSH**, Kalpathy,
Chidambareswaran [IN/US]; 3712A Laurel Bluff Circle,

High Point, NC 27265 (US). **MJALLI**, Adnan, M., M.
[US/US]; 2902 Ellington Court, Jamestown, NC 27282
(US). **ANDREWS**, Robert, Carl [US/US]; 3312 Morris
Farm Drive, Jamestown, NC 27282 (US). **SUBRAMA-
NIAN**, Govindan [IN/US]; 1835 Morgans Mill Way, High
Point, NC 27265 (US). **ANKERSEN**, Michael [DK/DK];
Frøkær 8, Ganløse, DK-3660 Stenløse (DK). **VEDSØ**,
Per [DK/DK]; Dalgas Have 4, 2.th., DK-2000 Frederiks-
berg (DK). **MURRAY**, Anthony [GB/DK]; Esthersvej
32, 1. th., DK-2900 Hellerup (DK). **JEPPESEN**, Lone
[DK/DK]; Malmrosevej 121, DK-2830 Virum (DK).

(74) Common Representative: **NOVO NORDISK A/S**; Cor-
porate Patents, Novo Allé, DK-2880 Bagsværd (DK).

(81) Designated States (national): AE, AG, AL, AM, AT, AU,
AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU,
CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH,
GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC,
LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW,
MX, MZ, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC,
SD, SE, SG, SK, SL, SY, TJ, TM, TN, TR, TT, TZ, UA,
UG, US, UZ, VC, VN, YU, ZA, ZM, ZW.

(84) Designated States (regional): ARIPO patent (GH, GM,
KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW),
Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM),
European patent (AT, BE, BG, CH, CY, CZ, DE, DK, EE,
ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PT, RO,
SE, SI, SK, TR), OAPI patent (BF, BJ, CF, CG, CI, CM,
GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).


Published:

— with international search report

For two-letter codes and other abbreviations, refer to the "Guid-
ance Notes on Codes and Abbreviations" appearing at the begin-
ning of each regular issue of the PCT Gazette.

WO 2004/002481 A1

(54) Title: **ARYL CARBOXYL DERIVATIVES AS THERAPEUTIC AGENTS**



(57) Abstract: This invention relates to aryl carbonyl derivatives of the general formula (I), which are activators of glucokinase which may be useful for the management, treatment, control, or adjunct treatment of diseases, where increasing glucokinase activity is beneficial. (Formul 1).

GLUCOKINASE ACTIVATORS

FIELD OF THE INVENTION

This invention relates to compounds which are activators of glucokinase (GK), which may be useful for the management, treatment, control, or adjunct treatment of diseases,
5 where increasing glucokinase activity is beneficial.

BACKGROUND OF THE INVENTION

Diabetes is characterised by an impaired glucose metabolism manifesting itself among other things by an elevated blood glucose level in the diabetic patients. Underlying defects lead to a classification of diabetes into two major groups: Type 1 diabetes, or insulin
10 demanding diabetes mellitus (IDDM), which arises when patients lack β -cells producing insulin in their pancreatic glands, and type 2 diabetes, or non-insulin dependent diabetes mellitus (NIDDM), which occurs in patients with an impaired β -cell function besides a range of other abnormalities.

Type 1 diabetic patients are currently treated with insulin, while the majority of type 2
15 diabetic patients are treated either with sulphonylureas that stimulate β -cell function or with agents that enhance the tissue sensitivity of the patients towards insulin or with insulin. Among the agents applied to enhance tissue sensitivity towards insulin metformin is a representative example.

Even though sulphonylureas are widely used in the treatment of NIDDM this therapy
20 is, in most instances, not satisfactory: In a large number of NIDDM patients sulphonylureas do not suffice to normalise blood sugar levels and the patients are, therefore, at high risk for acquiring diabetic complications. Also, many patients gradually lose the ability to respond to treatment with sulphonylureas and are thus gradually forced into insulin treatment. This shift of patients from oral hypoglycaemic agents to insulin therapy is usually ascribed to
25 exhaustion of the β -cells in NIDDM patients.

In normal subjects as well as in diabetic subjects, the liver produces glucose in order to avoid hypoglycaemia. This glucose production is derived either from the release of glucose from glycogen stores or from gluconeogenesis, which is a de novo intracellular
30 synthesis of glucose. In type 2 diabetes, however, the regulation of hepatic glucose output is poorly controlled and is increased, and may be doubled after an overnight fast. Moreover, in these patients there exists a strong correlation between the increased fasting plasma glucose levels and the rate of hepatic glucose production. Similarly, hepatic glucose

production will be increased in type 1 diabetes, if the disease is not properly controlled by insulin treatment.

Since existing forms of therapy of diabetes does not lead to sufficient glycaemic control and therefore are unsatisfactory, there is a great demand for novel therapeutic

5 approaches.

Atherosclerosis, a disease of the arteries, is recognized to be the leading cause of death in the United States and Western Europe. The pathological sequence leading to atherosclerosis and occlusive heart disease is well known. The earliest stage in this sequence is the formation of "fatty streaks" in the carotid, coronary and cerebral arteries and in the
10 aorta. These lesions are yellow in colour due to the presence of lipid deposits found principally within smooth-muscle cells and in macrophages of the intima layer of the arteries and aorta. Further, it is postulated that most of the cholesterol found within the fatty streaks, in turn, give rise to development of the "fibrous plaque", which consists of accumulated intimal smooth muscle cells laden with lipid and surrounded by extra-cellular lipid, collagen,
15 elastin and proteoglycans. The cells plus matrix form a fibrous cap that covers a deeper deposit of cell debris and more extracellular lipid. The lipid is primarily free and esterified cholesterol. The fibrous plaque forms slowly, and is likely in time to become calcified and necrotic, advancing to the "complicated lesion" which accounts for the arterial occlusion and tendency toward mural thrombosis and arterial muscle spasm that characterize advanced
20 atherosclerosis.

Epidemiological evidence has firmly established hyperlipidemia as a primary risk factor in causing cardiovascular disease (CVD) due to atherosclerosis. In recent years, leaders of the medical profession have placed renewed emphasis on lowering plasma cholesterol levels, and low density lipoprotein cholesterol in particular, as an essential step in
25 prevention of CVD. The upper limits of "normal" are now known to be significantly lower than heretofore appreciated. As a result, large segments of Western populations are now realized to be at particular high risk. Independent risk factors include glucose intolerance, left ventricular hypertrophy, hypertension, and being of the male sex. Cardiovascular disease is especially prevalent among diabetic subjects, at least in part because of the existence of
30 multiple independent risk factors in this population. Successful treatment of hyperlipidemia in the general population, and in diabetic subjects in particular, is therefore of exceptional medical importance.

Hypertension (or high blood pressure) is a condition, which occurs in the human population as a secondary symptom to various other disorders such as renal artery stenosis,
35 pheochromocytoma, or endocrine disorders. However, hypertension is also evidenced in

many patients in whom the causative agent or disorder is unknown. While such "essential" hypertension is often associated with disorders such as obesity, diabetes, and hypertriglyceridemia, the relationship between these disorders has not been elucidated. Additionally, many patients display the symptoms of high blood pressure in the complete
5 absence of any other signs of disease or disorder.

It is known that hypertension can directly lead to heart failure, renal failure, and stroke (brain haemorrhaging). These conditions are capable of causing short-term death in a patient. Hypertension can also contribute to the development of atherosclerosis and coronary disease. These conditions gradually weaken a patient and can lead to long-term death.

10 The exact cause of essential hypertension is unknown, though a number of factors are believed to contribute to the onset of the disease. Among such factors are stress, uncontrolled emotions, unregulated hormone release (the renin, angiotensin aldosterone system), excessive salt and water due to kidney malfunction, wall thickening and hypertrophy of the vasculature resulting in constricted blood vessels and genetic factors.

15 The treatment of essential hypertension has been undertaken bearing the foregoing factors in mind. Thus a broad range of beta-blockers, vasoconstrictors, angiotensin converting enzyme inhibitors and the like have been developed and marketed as antihypertensives. The treatment of hypertension utilizing these compounds has proven beneficial in the prevention of short-interval deaths such as heart failure, renal failure, and
20 brain haemorrhaging. However, the development of atherosclerosis or heart disease due to hypertension over a long period of time remains a problem. This implies that although high blood pressure is being reduced, the underlying cause of essential hypertension is not responding to this treatment.

Hypertension has been associated with elevated blood insulin levels, a condition
25 known as hyperinsulinemia. Insulin, a peptide hormone whose primary actions are to promote glucose utilization, protein synthesis and the formation and storage of neutral lipids, also acts to promote vascular cell growth and increase renal sodium retention, among other things. These latter functions can be accomplished without affecting glucose levels and are known causes of hypertension. Peripheral vasculature growth, for example, can cause
30 constriction of peripheral capillaries, while sodium retention increases blood volume. Thus, the lowering of insulin levels in hyperinsulinemics can prevent abnormal vascular growth and renal sodium retention caused by high insulin levels and thereby alleviates hypertension.

Cardiac hypertrophy is a significant risk factor in the development of sudden death, myocardial infarction, and congestive heart failure. These cardiac events are due, at least in
35 part, to increased susceptibility to myocardial injury after ischemia and reperfusion, which

can occur in out-patient as well as perioperative settings. There is an unmet medical need to prevent or minimize adverse myocardial perioperative outcomes, particularly perioperative myocardial infarction. Both non-cardiac and cardiac surgery are associated with substantial risks for myocardial infarction or death. Some 7 million patients undergoing non-cardiac surgery are considered to be at risk, with incidences of perioperative death and serious cardiac complications as high as 20-25% in some series. In addition, of the 400,000 patients undergoing coronary by-pass surgery annually, perioperative myocardial infarction is estimated to occur in 5% and death in 1-2%. There is currently no drug therapy in this area, which reduces damage to cardiac tissue from perioperative myocardial ischemia or enhances cardiac resistance to ischemic episodes. Such a therapy is anticipated to be life-saving and reduce hospitalizations, enhance quality of life and reduce overall health care costs of high risk patients.

Obesity is a well-known risk factor for the development of many very common diseases such as atherosclerosis, hypertension, and diabetes. The incidence of obese people and thereby also these diseases is increasing throughout the entire industrialised world. Except for exercise, diet and food restriction no convincing pharmacological treatment for reducing body weight effectively and acceptably currently exist. However, due to its indirect but important effect as a risk factor in mortal and common diseases it will be important to find treatment for obesity and/or means of appetite regulation.

The term obesity implies an excess of adipose tissue. In this context obesity is best viewed as any degree of excess adiposity that imparts a health risk. The cut off between normal and obese individuals can only be approximated, but the health risk imparted by the obesity is probably a continuum with increasing adiposity. The Framingham study demonstrated that a 20% excess over desirable weight clearly imparted a health risk (Mann GV N.Engl.J.Med 291:226, 1974). In the United States a National Institutes of Health consensus panel on obesity agreed that a 20% increase in relative weight or a body mass index (BMI = body weight in kilograms divided by the square of the height in meters) above the 85th percentile for young adults constitutes a health risk. By the use of these criteria 20 to 30 percent of adult men and 30 to 40 percent of adult women in the United States are obese. (NIH, Ann Intern Med 103:147, 1985).

Even mild obesity increases the risk for premature death, diabetes, hypertension, atherosclerosis, gallbladder disease, and certain types of cancer. In the industrialised western world the prevalence of obesity has increased significantly in the past few decades. Because of the high prevalence of obesity and its health consequences, its prevention and treatment should be a high public health priority.

When energy intake exceeds expenditure, the excess calories are stored in adipose tissue, and if this net positive balance is prolonged, obesity results, i.e. there are two components to weight balance, and an abnormality on either side (intake or expenditure) can lead to obesity.

5 The regulation of eating behaviour is incompletely understood. To some extent appetite is controlled by discrete areas in the hypothalamus: a feeding centre in the ventrolateral nucleus of the hypothalamus (VLH) and a satiety centre in the ventromedial hypothalamus (VMH). The cerebral cortex receives positive signals from the feeding centre that stimulate eating, and the satiety centre modulates this process by sending inhibitory
10 impulses to the feeding centre. Several regulatory processes may influence these hypothalamic centres. The satiety centre may be activated by the increases in plasma glucose and/or insulin that follow a meal. Meal-induced gastric distension is another possible inhibitory factor. Additionally the hypothalamic centres are sensitive to catecholamines, and beta-adrenergic stimulation inhibits eating behaviour. Ultimately, the cerebral cortex controls
15 eating behaviour, and impulses from the feeding centre to the cerebral cortex are only one input. Psychological, social, and genetic factors also influence food intake.

At present a variety of techniques are available to effect initial weight loss. Unfortunately, initial weight loss is not an optimal therapeutic goal. Rather, the problem is that most obese patients eventually regain their weight. An effective means to establish
20 and/or sustain weight loss is the major challenge in the treatment of obesity today.

WO 00/58293, WO 01/44216, WO/0183465, WO/0183478, WO/0185706, WO 01/85707, and WO02/08209, to Hoffman-La Roche, discloses compounds as glucokinase activators.

SUMMARY OF THE INVENTION

25 This invention provides amide derivatives of the general formula (I), as described below, as activators of glucokinase. The compounds of the present invention are useful as activators of glucokinase and thus are useful for the management, treatment, control and adjunct treatment of diseases where increasing the activity of glucokinase is beneficial. Such diseases include type I diabetes and type II diabetes. The present invention provides
30 compounds as described below, pharmaceutical compositions comprising the compounds, their use for increasing the activity of glucokinase, their use in preparation of a medicament for treating said diseases and conditions and the use of compounds or pharmaceutical preparations of the present invention for treating said diseases and conditions as well as methods for treating said diseases and conditions, which methods comprise administering to

a subject in need thereof an effective amount of a compound according to the present invention.

The present invention also provides glucokinase activators, for instance of the general formula (I), which are glucose sensitive glucokinase activators, that is, glucokinase
5 activators, the activity of which decreases with increasing glucose concentrations.

The present invention also provides glucokinase activators, for instance of the general formula (I), which are liver specific glucokinase activators, that is, glucokinase activators which increase glucose utilization in the liver (i.e. increase glycogen deposition) without inducing any increase in insulin secretion in response to glucose.

10 The present invention provides the use of a compound or a pharmaceutical preparation according to the present invention for treatment of metabolic disorders.

The present invention provides the use of a compound or a pharmaceutical preparation according to the present invention for blood glucose lowering.

15 The present invention provides the use of a compound or a pharmaceutical preparation according to the present invention for prevention of hyperglycemia.

The present invention provides the use of a compound or a pharmaceutical preparation according to the present invention for treatment of impaired glucose tolerance IGT.

20 The present invention provides the use of a compound or a pharmaceutical preparation according to the present invention for treatment of Syndrome X.

The present invention provides the use of a compound or a pharmaceutical preparation according to the present invention for the treatment of impaired fasting glucose (IFG).

25 The present invention provides the use of a compound or a pharmaceutical preparation according to the present invention for treatment of type 2 diabetes.

The present invention provides the use of a compound or a pharmaceutical preparation according to the present invention for treatment of type 1 diabetes.

30 The present invention provides the use of a compound or a pharmaceutical preparation according to the present invention for delaying the progression of impaired glucose tolerance (IGT) to type 2 diabetes.

The present invention provides the use of a compound or a pharmaceutical preparation according to the present invention for delaying the progression of non-insulin requiring type 2 diabetes to insulin requiring type 2 diabetes.

The present invention provides the use of a compound or a pharmaceutical preparation according to the present invention for treatment of dyslipidemia or hyperlipidemia.

5 The present invention provides the use of a compound or a pharmaceutical preparation according to the present invention for treatment of hypertension.

The present invention provides the use of a compound or a pharmaceutical preparation according to the present invention for the treatment of obesity.

The present invention provides the use of a compound or a pharmaceutical preparation according to the present invention for lowering of food intake

10 The present invention provides the use of a compound or a pharmaceutical preparation according to the present invention for appetite regulation.

The present invention provides the use of a compound or a pharmaceutical preparation according to the present invention for regulating feeding behaviour.

15 The present invention provides the use of a compound or a pharmaceutical preparation according to the present invention for enhancing the secretion of enteroincretins, such as GLP-1.

The present invention provides the use of a compound or a pharmaceutical preparation according to the present invention for the adjuvant treatment of type 1 diabetes for preventing the onset of diabetic complications.

20 The present invention provides the use of a compound or a pharmaceutical preparation according to the present invention for increasing the number and/or the size of beta cells in a mammalian subject.

The present invention provides the use of a compound or a pharmaceutical preparation according to the present invention for treatment of beta cell degeneration, in particular apoptosis of beta cells.

25 The present invention provides the use of a compound or a pharmaceutical preparation according to the present invention for treatment of functional dyspepsia, in particular irritable bowel syndrome.

30 The present invention provides the use of a compound according to the present invention for the preparation of a medicament for treatment of metabolic disorders.

The present invention provides the use of a compound according to the present invention for the preparation of a medicament for blood glucose lowering.

The present invention provides the use of a compound according to the present invention for the preparation of a medicament for the treatment of hyperglycemia.

The present invention provides the use of a compound according to the present invention for the preparation of a medicament for the treatment of IGT.

The present invention provides the use of a compound according to the present invention for the preparation of a medicament for the treatment of Syndrome X.

5 The present invention provides the use of a compound according to the present invention for the preparation of a medicament for the treatment of impaired fasting glucose (IFG).

The present invention provides the use of a compound according to the present invention for the preparation of a medicament for the treatment of type 2 diabetes.

10 The present invention provides the use of a compound according to the present invention for the preparation of a medicament for the treatment of type 1 diabetes.

The present invention provides the use of a compound according to the present invention for the preparation of a medicament for delaying the progression of impaired glucose tolerance (IGT) to type 2 diabetes.

15 The present invention provides the use of a compound according to the present invention for the preparation of a medicament for delaying the progression of non-insulin requiring type 2 diabetes to insulin requiring type 2 diabetes.

The present invention provides the use of a compound according to the present invention for the preparation of a medicament for the treatment of dyslipidemia.

20 The present invention provides the use of a compound according to the present invention for the preparation of a medicament for the treatment of hyperlipidemia.

The present invention provides the use of a compound according to the present invention for the preparation of a medicament for the treatment of hypertension.

25 The present invention provides the use of a compound according to the present invention for the preparation of a medicament for lowering of food intake.

The present invention provides the use of a compound according to the present invention for the preparation of a medicament for appetite regulation.

The present invention provides the use of a compound according to the present invention for the preparation of a medicament for the treatment of obesity.

30 The present invention provides the use of a compound according to the present invention for the preparation of a medicament for regulating feeding behaviour.

The present invention provides the use of a compound according to the present invention for the preparation of a medicament for enhancing the secretion of enteroinsulins, such as GLP-1.

The present invention provides the use of a compound according to the present invention for the preparation of a medicament for the adjuvant treatment of type 1 diabetes for preventing the onset of diabetic complications.

5 The present invention provides the use of a compound according to the present invention for the preparation of a medicament for increasing the number and/or the size of beta cells in a mammalian subject.

The present invention provides the use of a compound according to the present invention for the preparation of a medicament for treatment of beta cell degeneration, in particular apoptosis of beta cells.

10 The present invention provides the use of a compound according to the present invention for the preparation of a medicament for treatment of functional dyspepsia, in particular irritable bowel syndrome.

The present invention provides a method of preventing hypoglycaemia comprising administration of a liver-specific glucokinase activator.

15 The present invention provides the use of a liver-specific glucokinase activator for the preparation of a medicament for the prevention of hypoglycaemia.

Other embodiments and aspects are as defined below and by the appended claims.

DEFINITIONS

20 In the structural formulas given herein and throughout the present specification, the following terms have the indicated meaning:

The term "optionally substituted" as used herein means that the group in question is either unsubstituted or substituted with one or more of the substituents specified. When the group in question are substituted with more than one substituent the substituent may be the
25 same or different.

The term "adjacent" as used herein regards the relative positions of two atoms or variables, these two atoms or variables sharing a bond or one variable preceding or succeeding the other in a variable specification. By way of example, "atom A adjacent to atom B" means that the two atoms A and B share a bond.

30 The term "halogen" or "halo" means fluorine, chlorine, bromine or iodine.

The term "perhalomethyl" means trifluoromethyl, trichloromethyl, tribromomethyl, or triiodomethyl.

The use of prefixes of this structure: C_{x-y} -alkyl, C_{x-y} -alkenyl, C_{x-y} -alkynyl, C_{x-y} -cycloalyl or C_{x-y} -cycloalkyl- C_{x-y} -alkenyl- designates radical of the designated type having from x to y carbon atoms.

The term "alkyl" as used herein, alone or in combination, refers to a straight or
5 branched chain saturated monovalent hydrocarbon radical having from one to ten carbon atoms, for example C_{1-8} -alkyl or C_{1-8} -alkyl. Typical C_{1-8} -alkyl groups and C_{1-8} -alkyl groups include, but are not limited to e.g. methyl, ethyl, n-propyl, isopropyl, n-butyl, sec-butyl, isobutyl, tert-butyl, n-pentyl, 2-methylbutyl, 3-methylbutyl, 4-methylpentyl, neopentyl, n-pentyl, n-hexyl, 1,2-dimethylpropyl, 2,2-dimethylpropyl, 1,2,2-trimethylpropyl and the like.
10 The term " C_{1-8} -alkyl" as used herein also includes secondary C_{3-8} -alkyl and tertiary C_{4-8} -alkyl. The term " C_{1-8} -alkyl" as used herein also includes secondary C_{3-8} -alkyl and tertiary C_{4-8} -alkyl.

The term "alkylene" as used herein, alone or in combination, refers to a straight or branched chain saturated divalent hydrocarbon radical having from one to ten carbon atoms, for example C_{1-8} -alkylene or C_{1-8} -alkylene. Examples of "alkylene" as used herein include, but
15 are not limited to, methylene, ethylene, and the like.

The term "alkenyl" as used herein, alone or in combination, refers to a straight or branched chain monovalent hydrocarbon radical containing from two to ten carbon atoms and at least one carbon-carbon double bond, for example C_{2-8} -alkenyl or C_{2-8} -alkenyl. Typical C_{2-8} -alkenyl groups and C_{2-8} -alkenyl groups include, but are not limited to, vinyl, 1-propenyl,
20 2-propenyl, iso-propenyl, 1,3-butadienyl, 1-butenyl, 2-butenyl, 3-butenyl, 2-methyl-1-propenyl, 1-pentenyl, 2-pentenyl, 3-pentenyl, 4-pentenyl, 3-methyl-2-butenyl, 1-hexenyl, 2-hexenyl, 3-hexenyl, 2,4-hexadienyl, 5-hexenyl and the like.

The term "alkenylene" as used herein, alone or in combination, refers to a straight or branched chain divalent hydrocarbon radical having from two to ten carbon atoms and at
25 least one carbon-carbon double bond, for example C_{2-8} -alkenylene or C_{2-8} -alkenylene. Typical C_{2-8} -alkenylene groups and C_{2-8} -alkenylene groups include, but are not limited to, ethene-1,2-diyl, propene-1,3-diyl, methylene-1,1-diyl, and the like.

The term "alkynyl" as used herein alone or in combination, refers to a straight or branched monovalent hydrocarbon radical containing from two to ten carbon atoms and at
30 least one triple carbon-carbon bond, for example C_{2-8} -alkynyl or C_{2-8} -alkynyl. Typical C_{2-8} -alkynyl groups and C_{2-8} -alkynyl groups include, but are not limited to, ethynyl, 1-propynyl, 2-propynyl, 1-butyne, 2-butyne, 3-butyne, 1-pentyne, 2-pentyne, 3-pentyne, 4-pentyne, 1-hexynyl, 2-hexynyl, 3-hexynyl, 5-hexynyl, 2,4-hexadiynyl and the like.

The term "alkynylene" as used herein alone or in combination, refers to a straight or
35 branched chain divalent hydrocarbon radical having from two to ten carbon atoms and at

least one carbon-carbon triple bond, for example C_{2-8} -alkynylene or C_{2-8} -alkynylene. Typical C_{2-8} -alkynylene groups and C_{2-8} -alkynylene groups include, but are not limited to, ethyne-1,2-diyl, propyne-1,3-diyl, and the like.

5 The term "cycloalkyl" as used herein, alone or in combination, refers to a non-aromatic monovalent hydrocarbon radical having from three to twelve carbon atoms, and optionally with one or more degrees of unsaturation, for example C_{3-8} -cycloalkyl. Such a ring may be optionally fused to one or more benzene rings or to one or more of other cycloalkyl ring(s). Typical C_{3-8} -cycloalkyl groups include, but are not limited to, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, cyclohexenyl, cycloheptyl, cycloheptenyl, cyclooctyl and the like.

10 The term "cycloalkylene" as used herein, alone or in combination, refers to a non-aromatic carbocyclic divalent hydrocarbon radical having from three to twelve carbon atoms and optionally possessing one or more degrees of unsaturation, for example C_{3-8} -cycloalkylene. Such a ring may be optionally fused to one or more benzene rings or to one or more of other cycloalkyl ring(s). Typical C_{3-8} -cycloalkylene groups include, but are not limited to, 15 cyclopropyl-1,1-diyl, cyclopropyl-1,2-diyl, cyclobutyl-1,2-diyl, cyclopentyl-1,3-diyl, cyclohexyl-1,4-diyl, cycloheptyl-1,4-diyl, or cyclooctyl-1,5-diyl, and the like.

The term "heterocyclic" or the term "heterocyclyl" as used herein, alone or in combination, refers to a three to twelve membered heterocyclic ring having one or more degrees of unsaturation containing one or more heteroatomic substitutions selected from S, 20 SO, SO₂, O, or N, for example C_{3-8} -heterocyclyl. Such a ring may be optionally fused to one or more of another "heterocyclic" ring(s) or cycloalkyl ring(s). Typical C_{3-8} -heterocyclyl groups include, but are not limited to, tetrahydrofuran, 1,4-dioxane, 1,3-dioxane, piperidine, pyrrolidine, morpholine, piperazine, and the like.

The term "heterocyclylene" as used herein, alone or in combination, refers to a three 25 to twelve-membered heterocyclic ring diradical optionally having one or more degrees of unsaturation containing one or more heteroatoms selected from S, SO, SO₂, O, or N. Such a ring may be optionally fused to one or more benzene rings or to one or more of another "heterocyclic" rings or cycloalkyl rings. Examples of "heterocyclylene" include, but are not limited to, tetrahydrofuran-2,5-diyl, morpholine-2,3-diyl, pyran-2,4-diyl, 1,4-dioxane-2,3-diyl, 30 1,3-dioxane-2,4-diyl, piperidine-2,4-diyl, piperidine-1,4-diyl, pyrrolidine-1,3-diyl, morpholine-2,4-diyl, piperazine-1,4-diyl, and the like.

The term "alkoxy" as used herein, alone or in combination, refers to the monovalent radical R^aO-, where R^a is alkyl as defined above, for example C_{1-8} -alkyl giving C_{1-8} -alkoxy. Typical C_{1-8} -alkoxy groups include, but are not limited to, methoxy, ethoxy, n-propoxy,

isopropoxy, butoxy, *sec*-butoxy, *tert*-butoxy, pentoxy, isopentoxy, hexoxy, isohexoxy and the like.

The term "alkylthio" as used herein, alone or in combination, refers to a straight or branched monovalent radical comprising an alkyl group as described above linked through a divalent sulphur atom having its free valence bond from the sulphur atom, for example C₁₋₈-alkylthio. Typical C₁₋₈-alkylthio groups include, but are not limited to, methylthio, ethylthio, propylthio, butylthio, pentylthio, hexylthio and the like.

The term "alkoxycarbonyl" as used herein refers to the monovalent radical R^aOC(O)-, where R^a is alkyl as described above, for example C₁₋₈-alkoxycarbonyl. Typical C₁₋₈-alkoxycarbonyl groups include, but are not limited to, methoxycarbonyl, ethoxycarbonyl, propoxycarbonyl, isopropoxycarbonyl, n-butoxycarbonyl, *sec*-butoxycarbonyl, *tert*butoxycarbonyl, 3-methylbutoxycarbonyl, n-hexoxycarbonyl and the like.

The term "carbamoyl" as used herein refers to NH₂C(O)-.

The term "aryl" as used herein refers to a carbocyclic aromatic ring radical or to a aromatic ring system radical. Aryl is also intended to include the partially hydrogenated derivatives of the carbocyclic systems.




The term "heteroaryl", as used herein, alone or in combination, refers to an aromatic ring radical with for instance 5 to 7 member atoms, or to a aromatic ring system radical with for instance from 7 to 18 member atoms, containing one or more heteroatoms selected from nitrogen, oxygen, or sulfur heteroatoms, wherein N-oxides and sulfur monoxides and sulfur dioxides are permissible heteroaromatic substitutions; such as e.g. furanyl, thienyl, thiophenyl, pyrrolyl, imidazolyl, pyrazolyl, triazolyl, tetrazolyl, thiazolyl, oxazolyl, isoxazolyl, oxadiazolyl, thiadiazolyl, isothiazolyl, pyridinyl, pyridazinyl, pyrazinyl, pyrimidinyl, quinolinyl, isoquinolinyl, benzofuranyl, benzothiophenyl, indolyl, and indazolyl, and the like. Heteroaryl is also intended to include the partially hydrogenated derivatives of the heterocyclic systems enumerated below.

Examples of "aryl" and "heteroaryl" includes, but are not limited to phenyl, biphenyl, indene, fluorene, naphthyl (1-naphthyl, 2-naphthyl), anthracene (1-anthracenyl, 2-anthracenyl, 3-anthracenyl), thiophene (2-thienyl, 3-thienyl), furyl (2-furyl, 3-furyl), indolyl, oxadiazolyl, isoxazolyl, thiadiazolyl, oxatriazolyl, thiatriazolyl, quinazolin, fluorenyl, xanthenyl, isoindanyl, benzhydryl, acridinyl, thiazolyl, pyrrolyl (1-pyrrolyl, 2-pyrrolyl, 3-pyrrolyl), pyrazolyl (1-pyrazolyl, 3-pyrazolyl, 4-pyrazolyl, 5-pyrazolyl), imidazolyl (1-imidazolyl, 2-imidazolyl, 4-imidazolyl, 5-imidazolyl), triazolyl (1,2,3-triazol-1-yl, 1,2,3-triazol-4-yl, 1,2,3-triazol-5-yl, 1,2,4-triazol-3-yl, 1,2,4-triazol-5-yl), oxazolyl (2-oxazolyl, 4-oxazolyl, 5-oxazolyl), isooxazolyl (isooxazo-3-yl, isooxazo-4-yl, isooxazo-5-yl), isothiazolyl (isothiazo-3-yl, isothiazo-4-yl,

isothiaz-5-yl) thiazolyl (2-thiazolyl, 4-thiazolyl, 5-thiazolyl), pyridyl (2-pyridyl, 3-pyridyl, 4-pyridyl), pyrimidinyl (2-pyrimidinyl, 4-pyrimidinyl, 5-pyrimidinyl, 6-pyrimidinyl), pyrazinyl, pyridazinyl (3-pyridazinyl, 4-pyridazinyl, 5-pyridazinyl), quinolyl (2-quinolyl, 3-quinolyl, 4-quinolyl, 5-quinolyl, 6-quinolyl, 7-quinolyl, 8-quinolyl), isoquinolyl (1-isoquinolyl, 3-isoquinolyl, 4-isoquinolyl, 5-isoquinolyl, 6-isoquinolyl, 7-isoquinolyl, 8-isoquinolyl), benzo[b]furanyl (2-benzo[b]furanyl, 3-benzo[b]furanyl, 4-benzo[b]furanyl, 5-benzo[b]furanyl, 6-benzo[b]furanyl, 7-benzo[b]furanyl), 2,3-dihydro-benzo[b]furanyl (2-(2,3-dihydro-benzo[b]furanyl), 3-(2,3-dihydro-benzo[b]furanyl), 4-(2,3-dihydro-benzo[b]furanyl), 5-(2,3-dihydro-benzo[b]furanyl), 6-(2,3-dihydro-benzo[b]furanyl), 7-(2,3-dihydro-benzo[b]furanyl)), benzo[b]thiophenyl (benzo[b]thiophen-2-yl, benzo[b]thiophen-3-yl, benzo[b]thiophen-4-yl, benzo[b]thiophen-5-yl, benzo[b]thiophen-6-yl, benzo[b]thiophen-7-yl), 2,3-dihydro-benzo[b]thiophenyl (2,3-dihydro-benzo[b]thiophen-2-yl, 2,3-dihydro-benzo[b]thiophen-3-yl, 2,3-dihydro-benzo[b]thiophen-4-yl, 2,3-dihydro-benzo[b]thiophen-5-yl, 2,3-dihydro-benzo[b]thiophen-6-yl, 2,3-dihydro-benzo[b]thiophen-7-yl), indolyl (1-indolyl, 2-indolyl, 3-indolyl, 4-indolyl, 5-indolyl, 6-indolyl, 7-indolyl), indazole (1-indazolyl, 3-indazolyl, 4-indazolyl, 5-indazolyl, 6-indazolyl, 7-indazolyl), benzimidazolyl (1-benzimidazolyl, 2-benzimidazolyl, 4-benzimidazolyl, 5-benzimidazolyl, 6-benzimidazolyl, 7-benzimidazolyl, 8-benzimidazolyl), benzoxazolyl (2-benzoxazolyl, 3-benzoxazolyl, 4-benzoxazolyl, 5-benzoxazolyl, 6-benzoxazolyl, 7-benzoxazolyl), benzothiazolyl (2-benzothiazolyl, 4-benzothiazolyl, 5-benzothiazolyl, 6-benzothiazolyl, 7-benzothiazolyl), carbazolyl (1-carbazolyl, 2-carbazolyl, 3-carbazolyl, 4-carbazolyl), 5H-dibenz[b,f]azepine (5H-dibenz[b,f]azepine-1-yl, 5H-dibenz[b,f]azepine-2-yl, 5H-dibenz[b,f]azepine-3-yl, 5H-dibenz[b,f]azepine-4-yl, 5H-dibenz[b,f]azepine-5-yl), 10,11-dihydro-5H-dibenz[b,f]azepine (10,11-dihydro-5H-dibenz[b,f]azepine-1-yl, 10,11-dihydro-5H-dibenz[b,f]azepine-2-yl, 10,11-dihydro-5H-dibenz[b,f]azepine-3-yl, 10,11-dihydro-5H-dibenz[b,f]azepine-4-yl, 10,11-dihydro-5H-dibenz[b,f]azepine-5-yl), benzo[1,3]dioxole (2-benzo[1,3]dioxole, 4-benzo[1,3]dioxole, 5-benzo[1,3]dioxole, 6-benzo[1,3]dioxole, 7-benzo[1,3]dioxole), purinyl, and tetrazolyl (5-tetrazolyl, N-tetrazolyl).

The present invention also relates to partly or fully saturated analogues of the ring systems mentioned above.


When two such terms are used in combination, such as in aryl-alkyl-, heteroaryl-alkyl-, cycloalkyl-C₁₋₆-alkyl- and the like, it is to be understood that the first mentioned radical is a substituent on the latter mentioned radical, where the point of substitution is on the latter of the radicals, for example



The term "arylene", as used herein, alone or in combination, refers to carbocyclic aromatic ring diradical or to a aromatic ring system diradical. Examples of "arylene" include, but are not limited to, benzene-1,4-diyl, naphthalene-1,8-diyl, and the like. The term "arylene" alone or in combination also include other divalent radicals of the monovalent radicals mentioned in the definition of aryl.


The term "heteroarylene", as used herein, alone or in combination, refers to a five to seven membered aromatic ring diradical, or to a aromatic ring system diradical, containing one or more heteroatoms selected from nitrogen, oxygen, or sulfur heteroatoms, wherein N-oxides and sulfur monoxides and sulfur dioxides are permissible heteroaromatic substitutions. Examples of "heteroarylene" used herein are furan-2,5-diyl, thiophene-2,4-diyl, 1,3,4-oxadiazole-2,5-diyl, 1,3,4-thiadiazole-2,5-diyl, 1,3-thiazole-2,4-diyl, 1,3-thiazole-2,5-diyl, pyridine-2,4-diyl, pyridine-2,3-diyl, pyridine-2,5-diyl, pyrimidine-2,4-diyl, quinoline-2,3-diyl, and the like. The term "heteroarylene" alone or in combination also include other divalent radicals of the monovalent radicals mentioned in the definition of heteroaryl.

As used herein, the term "fused cycloalkylaryl" refers to a cycloalkyl group fused to an aryl group, the two having two atoms in common, and wherein the aryl group is the point of substitution. Examples of "fused cycloalkylaryl" used herein include 5-indanyl, 5,6,7,8-tetrahydro-2-naphthyl,




, and the like.

As used herein, the term "fused cycloalkylarylene" refers to a fused cycloalkylaryl, wherein the aryl group is divalent. Examples include




, and the like.

As used herein, the term "fused arylcycloalkyl" refers to an aryl group fused to a
5 cycloalkyl group, the two having two atoms in common, and wherein the cycloalkyl group is the point of substitution. Examples of "fused arylcycloalkyl" used herein include 1-indanyl, 2-indanyl, 1-(1,2,3,4-tetrahydronaphthyl),




, and the like.

As used herein, the term "fused arylcycloalkylene" refers to a fused arylcycloalkyl,
10 wherein the cycloalkyl group is divalent. Examples include




, and the like.

As used herein, the term "fused heterocyclaryl" refers to a heterocyclyl group
fused to an aryl group, the two having two atoms in common, and wherein the aryl group is
the point of substitution. Examples of "fused heterocyclaryl" used herein include 3,4-
15 methylenedioxy-1-phenyl,




, and the like

As used herein, the term "fused heterocyclarylene" refers to a fused
heterocyclaryl, wherein the aryl group is divalent. Examples include




, and the like.




As used herein, the term "fused arylheterocyclyl" refers to an aryl group fused to a heterocyclyl group, the two having two atoms in common, and wherein the heterocyclyl group is the point of substitution. Examples of "fused arylheterocyclyl" used herein include 2-(1,3-benzodioxolyl),




As used herein, the term "fused arylheterocyclene" refers to a fused arylheterocyclyl, wherein the heterocyclyl group is divalent. Examples include



As used herein, the term "fused cycloalkylheteroaryl" refers to a cycloalkyl group fused to a heteroaryl group, the two having two atoms in common, and wherein the heteroaryl group is the point of substitution. Examples of "fused cycloalkylheteroaryl" used herein include 5-aza-6-indanyl,




As used herein, the term "fused heteroarylcycloalkylene" refers to a fused heteroarylcycloalkyl, wherein the cycloalkyl group is divalent. Examples include




, and the like.

As used herein, the term "fused heterocyclylheteroaryl" refers to a heterocyclyl group fused to a heteroaryl group, the two having two atoms in common, and wherein the heteroaryl group is the point of substitution. Examples of "fused heterocyclylheteroaryl" used herein include 1,2,3,4-tetrahydro-beta-carbolin-8-yl,




and the like.

As used herein, the term "fused heterocyclylheteroarylene" refers to a fused heterocyclylheteroaryl, wherein the heteroaryl group is divalent. Examples include




, and the like.

As used herein, the term "fused heteroarylheterocyclyl" refers to a heteroaryl group fused to a heterocyclyl group, the two having two atoms in common, and wherein the heterocyclyl group is the point of substitution. Examples of "fused heteroarylheterocyclyl" used herein include 5-aza-2,3-dihydrobenzofuran-2-yl,



, and the like.

As used herein, the term "fused heteroarylheterocyclylene" refers to a fused heteroarylheterocyclyl, wherein the heterocyclyl group is divalent. Examples include



, and the like.

5 The term "alkylsulfanyl", as used herein, refers to the group R^aS- , where R^a is alkyl as described above.

The term "alkylsulfenyl", as used herein, refers to the group $R^aS(O)-$, where R^a is alkyl as described above.

The term "alkylsulfonyl", as used herein, refers to the group R^aSO_2- , where R^a is alkyl as described above.

10 The term "acyl", as used herein, refers to the group $R^aC(O)-$, where R^a is alkyl, alkenyl, alkynyl, cycloalkyl, cycloalkenyl, or heterocyclyl as described above.

The term "aroyl", as used herein, refers to the group $R^aC(O)-$, where R^a is aryl as described above.

15 The term "heteroaroyl", as used herein, refers to the group $R^aC(O)-$, where R^a is heteroaryl as described above.

The term "aryloxycarbonyl", as used herein, refers to the group $R^a-O-C(O)-$, where R^a is aryl as described above.

The term "acyloxy", as used herein, refers to the group $R^aC(O)O-$, where R^a is alkyl, alkenyl, alkynyl, cycloalkyl, cycloalkenyl, or heterocyclyl as described above.

20 The term "aryloxy", as used herein refers to the group R^a-O- , where R^a is aryl as described above.

The term "aroyloxy", as used herein, refers to the group $R^aC(O)O-$, where R^a is aryl as described above.

25 The term "heteroaroyloxy", as used herein, refers to the group $R^aC(O)O-$, where R^a is heteroaryl as described above.

Whenever the terms "alkyl", "cycloalkyl", "aryl", "heteroaryl" or the like or either of their prefix roots appear in a name of a substituent (e.g. arylalkoxyaryloxy) they shall be interpreted as including those limitations given above for "alkyl" and "aryl".

As used herein, the term "oxo" shall refer to the substituent $=O$.

30 As used herein, the term "mercapto" shall refer to the substituent $-SH$.

As used herein, the term "carboxy" shall refer to the substituent $-COOH$.

As used herein, the term "cyano" shall refer to the substituent -CN.

As used herein, the term "aminosulfonyl" shall refer to the substituent -SO₂NH₂.

As used herein, the term "sulfanyl" shall refer to the substituent -S-.

As used herein, the term "sulfenyl" shall refer to the substituent -S(O)-.

5 As used herein, the term "sulfonyl" shall refer to the substituent -S(O)₂-.

As used herein, the term "direct bond", where part of a structural variable specification, refers to the direct joining of the substituents flanking (preceding and succeeding) the variable taken as a "direct bond".

10 The term "lower", as used herein, refers to an group having between one and six carbons, and may be indicated with the prefix C_{x-6}-. Lower alkyl may thus be indicated as C₁₋₆-alkyl, while lower alkylene may be indicated as C₂₋₆-alkylene.

A radical such as C_{x-y}-cycloalkyl-C_{a-b}-alkenyl shall designate that the radical's point of attachment is in part of the radical mentioned last.

15 As used herein, the term "optionally" means that the subsequently described event(s) may or may not occur, and includes both event(s) which occur and events that do not occur.

As used herein, the term "substituted" refers to substitution with the named substituent or substituents, multiple degrees of substitution being allowed unless otherwise stated.

20 As used herein, the terms "contain" or "containing" can refer to in-line substitutions at any position along the above defined alkyl, alkenyl, alkynyl or cycloalkyl substituents with one or more of any of O, S, SO, SO₂, N, or N-alkyl, including, for example, -CH₂-O-CH₂-, -CH₂-SO₂-CH₂-, -CH₂-NH-CH₃ and so forth.

25 Certain of the above defined terms may occur more than once in the structural formulae, and upon such occurrence each term shall be defined independently of the other.

As used herein, the term "solvate" is a complex of variable stoichiometry formed by a solute (in this invention, a compound of formula (I)) and a solvent. Such solvents for the purpose of the present invention may not interfere with the biological activity of the solute. Solvents may be, by way of example, water, ethanol, or acetic acid.

30 As used herein, the term "biohydrolyzable ester" is an ester of a drug substance (in this invention, a compound of formula (I)) which either a) does not interfere with the biological activity of the parent substance but confers on that substance advantageous properties *in vivo* such as duration of action, onset of action, and the like, or b) is biologically inactive but is readily converted *in vivo* by the subject to the biologically active principle. The advantage is that, for example, the biohydrolyzable ester is orally absorbed from the gut and

35

is transformed to (I) in plasma. Many examples of such are known in the art and include by way of example lower alkyl esters (e.g., C₁₋₄), lower acyloxyalkyl esters, lower alkoxyacyloxyalkyl esters, alkoxyacyloxy esters, alkyl acylamino alkyl esters, and choline esters.

5 As used herein, the term "biohydrolyzable amide" is an amide of a drug substance (in this invention, a compound of general formula (I)) which either a) does not interfere with the biological activity of the parent substance but confers on that substance advantageous properties in vivo such as duration of action, onset of action, and the like, or b) is biologically inactive but is readily converted in vivo by the subject to the biologically active principle. The
10 advantage is that, for example, the biohydrolyzable amide is orally absorbed from the gut and is transformed to (I) in plasma. Many examples of such are known in the art and include by way of example lower alkyl amides, α -amino acid amides, alkoxyacyl amides, and alkylaminoalkylcarbonyl amides.

 As used herein, the term "prodrug" includes biohydrolyzable amides and
15 biohydrolyzable esters and also encompasses a) compounds in which the biohydrolyzable functionality in such a prodrug is encompassed in the compound of formula (I) and b) compounds which may be oxidized or reduced biologically at a given functional group to yield drug substances of formula (I). Examples of these functional groups include, but are not limited to, 1,4-dihydropyridine, N-alkylcarbonyl-1,4-dihydropyridine, 1,4-cyclohexadiene, tert-
20 butyl, and the like.

 The term "pharmacologically effective amount" or shall mean that amount of a drug or pharmaceutical agent that will elicit the biological or medical response of a tissue, animal or human that is being sought by a researcher or clinician. This amount can be a therapeutically effective amount. The term "therapeutically effective amount" shall mean that
25 amount of a drug or pharmaceutical agent that will elicit the therapeutic response of an animal or human that is being sought.

 The term "treatment" and "treating" as used herein means the management and care of a patient for the purpose of combating a disease, disorder or condition. The term is intended to include the full spectrum of treatments for a given disorder from which the patient is suffering,
30 such as the delaying of the progression of the disease, disorder or condition, the alleviation or relief of symptoms and complications, the prevention of the disease and/or the cure or elimination of the disease, disorder or condition. The patient to be treated is preferably a mammal, in particular a human being.

The term "human insulin" as used herein refers to naturally produced insulin or recombinantly produced insulin. Recombinant human insulin may be produced in any suitable host cell, for example the host cells may be bacterial, fungal (including yeast), insect, animal or plant cells.

The expression "insulin derivative" as used herein (and related expressions) refers to human insulin or an analogue thereof in which at least one organic substituent is bound to one or more of the amino acids.

By "analogue of human insulin" as used herein (and related expressions) is meant human insulin in which one or more amino acids have been deleted and/or replaced by other amino acids, including non-codeable amino acids, or human insulin comprising additional amino acids, i.e. more than 51 amino acids, such that the resulting analogue possesses insulin activity.

DETAILED DESCRIPTION OF THE INVENTION

Glucokinase (GK) plays an essential role in blood glucose homeostasis. GK catalyses glucose phosphorylation, and is the rate-limiting reaction for glycolysis in hepatocytes and pancreatic β -cells. In liver GK determine the rates of both glucose uptake and glycogen synthesis, and it is also thought to be essential for the regulation of various glucose-responsive genes (Girard, J. et al., *Annu Rev Nutr* 17, 325-352 (1997)). In the β -cells, GK determines glucose utilization and thus is necessary for glucose-stimulated insulin secretion. GK is also expressed in a population of neurones in the hypothalamus where it might be involved in feeding behaviour, and in the gut where it might contribute to the secretion of enteroincretins.

GK has two main distinctive characteristics: its expression, which is limited to tissues that require glucose-sensing (mainly liver and pancreatic β -cells), and its $S_{0.5}$ for glucose, which is much higher (8-12 mM) than that of the other members of the hexokinase family. Due to these kinetic characteristics, changes in serum glucose levels are paralleled by changes in glucose metabolism in liver which in turn regulate the balance between hepatic glucose output and glucose consumption.

Activators of glucokinase may thus be useful for treating diseases where increasing the activity of glucokinase is beneficial. Thus, there is a need for agents which activate glucokinase and increase glucokinase enzymatic activity. Such agents would be useful for the treatment of type I diabetes and type II diabetes.

Activators of glucokinase may also play a role in sensing low glucose levels and generating neurohumoral responses to hypoglycemia and may thus be useful for treating

those patients with type 1 diabetes, which have a higher tendency to suffer from hypoglycemia.


Type I diabetes mellitus is a complex disease characterized by an elevated blood glucose concentration and polyuria. Secondary to the persistent elevation in blood glucose, patients develop devastating complications such as retinopathy, nephropathy, neuropathy, and cardiovascular disease. A major goal to improve the diabetic phenotype is to reduce fasting and postprandial hyperglycemia and, thus, avoid or delay the onset of diabetic complications. The Diabetes Control and Complications Trial has indicated that tight glycemic control through administration of daily multiple insulin injections delays the onset of complications. However, such intensive therapy is associated with an increase in body weight and higher risk for development of hypoglycaemic events. Alternative treatments to achieve glucose control without these side effects are, therefore, being developed. The combination of GK overexpression in the liver and subcutaneous insulin injections provides better glycemic control in type 1 diabetic animals than treatment with insulin alone (Morral, N., et al. *Human Gene Therapy* 13, 1561-1570 (2002)). Moreover, overexpression of hepatic GK can compensate, in part, for the metabolic disorders developed by insulin receptor-deficient mice (Jackerott, M. et al. *Diabetologia* 45, 1292-1297 (2002)).

The present invention also relates to the use of a GK activator for the combined treatment of diabetes and obesity. GK, the GK regulatory protein and the KATP channel are expressed in neurones of the hypothalamus, a region of the brain that is important in the regulation of energy balance and the control of food intake. These neurones have been shown to express orectic and anorectic neuropeptides and have been assumed to be the glucose-sensing neurones within the hypothalamus that are either inhibited or excited by changes in ambient glucose concentrations (Mobbs, C. V. et al, *American Journal of Physiology, Endocrinology & Metabolism* 281, E649-54 (2001)). The ability of these neurones to sense changes in glucose levels is defective in a variety of genetic and experimentally induced models of obesity (Spanswick, D. et al, *Nature Neuroscience* 3, 757-8 (2000), Levin, B. E. et al, *Brain Research* 808, 317-9 (1998)). Intracerebroventricular (icv) infusion of glucose analogues, which are competitive inhibitors of glucokinase, stimulate food intake in lean rats (Kurata, K. et al, *Metabolism: Clinical & Experimental* 38, 46-51 (1989)). In contrast, icv infusion of glucose suppresses feeding (Kurata, K. et al, *Physiology & Behavior* 37, 615-20 (1986)). Small molecule activators of GK may thus decrease food intake and weight gain through central effects on GK. Therefore, GK activators may be of therapeutic use in treating eating disorders, including obesity, in addition to diabetes. The hypothalamic effects will be additive or synergistic to the effects of the same compounds acting in the liver

and/or pancreas in normalising glucose homeostasis, for the treatment of type 2 diabetes. Thus the GK/GK regulatory protein system can be described as a potential target of benefit in both diabetes and obesity.

The amplitude of glucose-induced insulin release is highly dependent on the action of the gastrointestinal hormones GLP-1 (glucagon-like peptide 1) and GIP. Unlike sulfonylureas, which stimulate insulin release at low as well as high glucose levels, the action of GLP-1 on β -cells is glucose dependent (Gromada, J. et al., *Pflügers Arch* 435, 583-594 (1998)). GLP-1 receptor agonist and drugs that slow the degradation of active GLP-1 are therefore under development as a novel treatments for type 2 diabetes. An alternative strategy would be to enhance endogenous GLP-1 levels. Of potential interest is the possibility that the release of GLP-1 and GIP might be regulated by glucokinase-expressing endocrine cells (Jetton, T.L. et al., *J. Biol. Chem.* 269, 3641-3654 (1994)) and glucose-responsive neurons (Liu, M. et al., *J. Neurosci.* 19, 10305-10317 (1999)). It has been reported that the release of GIP by intestinal K-cells is directly controlled by glucose (Kieffer, T.J. et al., *Am J Physiol* 267, E489-E496 (1994)), and GLP-1 secretion from GLUTag cells is triggered by glucose through a mechanism similar to that found in β -cells for insulin secretion (Reimann, F. et al, *Diabetes* 51, 2757-2763 (2002)). Small molecule activators of glucokinase may thus be used to increase GLP-1 and /or GIP secretion and thus for treatment, modulation, inhibition, decrease, reduction, arrest or prevention of beta cell degeneration, such as necrosis or apoptosis of β -cells.

In a first embodiment, the present invention provides compounds of the general formula (I).



(I)

5 wherein

A¹ is selected from the group consisting of arylene, heteroarylene, fused cycloalkylarylene, fused heterocycliarylene, fused cycloalkylheteroarylene, or fused heterocycliheteroarylene; optionally substituted with one or more substituents R²³, R²⁴, R²⁵, R²⁶, and R²⁷, wherein

10 R²³, R²⁴, R²⁵, R²⁶, and R²⁷ independently of each other are selected from the group consisting of

halogen, -C(O)OR², -CN, -CF₃, -OCF₃, -NO₂, -OR², -NR²R³, C₁₋₆-alkyl-Z-,

C₂₋₆-alkenyl-Z-, C₂₋₆-alkynyl-Z-, cycloalkyl-Z-, heterocyclyl-Z-, aryl-Z-, heteroaryl-Z-,

aryl-C₁₋₆-alkylene-Z-, heteroaryl-C₁₋₆-alkylene-Z-, heterocyclyl-C₁₋₆-alkylene-Z-,

cycloalkyl-C₁₋₆-alkylene-Z-, N(R⁴R⁵)-C₁₋₆-alkylene-Z-, R⁶-W¹-Z-, R⁶-W¹-N(R⁴)-Z-, R⁶-

15 N(R⁴)-Z, and R⁶-W¹-C₁₋₆-alkylene-Z-, wherein

R² and R³ independently of each other are hydrogen, C₁₋₆-alkyl,

aryl-C₁₋₆-alkylene-, heteroaryl-C₁₋₆-alkylene-, C₁₋₆-alkyl-arylene-,

C₁₋₆-alkyl-heteroarylene-, heteroaryl, or aryl;

or

20 R² and R³, when attached to the same nitrogen atom, together with said

nitrogen atom may form a 3 to 8 membered heterocyclic ring optionally

containing one or two further heteroatoms selected from nitrogen, oxygen

and sulfur, and optionally containing one or two double bonds;

Z and W¹ independently of each other are a direct bond, -O-, -N(R⁷)-, -S-,

25 -SO₂-, -C(O)N(R⁷)-, -N(R⁷)C(O)-, -N(R⁷)CON(R⁸)-, -N(R⁷)SO₂-, -SO₂N(R⁷)-,

-C(O)-O-, -N(R⁷)SO₂N(R⁸)-, or -O-C(O)-, wherein

R⁷ and R⁸ in each individual case independently of each other are

hydrogen or alkyl; and

R⁴, R⁵, and R⁶ independently of each other are selected from the group

30 consisting of hydrogen, aryl, alkyl, heteroaryl-alkylene-, and aryl-alkylene-;


or

R^4 and R^5 may be taken together to form a ring having the formula $-(CH_2)_j-Q-(CH_2)_k-$ bonded to the nitrogen atom to which R^4 and R^5 are attached, wherein

5

j and k independently of each other is 1, 2, 3, or 4; and

Q is a direct bond, $-CH_2-$, $-O-$, $-S-$, $-S(O_2)-$, $-C(O)-$, $-C(O)NH-$, $-NHC(O)-$, $-NHC(O)NH-$, $-NHSO_2-$, $-SO_2NH-$, $-C(O)-O-$, $-O-C(O)-$, $-NHSO_2NH-$,



10

wherein

R^9 and R^{10} independently of each other are selected from the group consisting of hydrogen, aryl, alkyl, and aryl-alkylene;

L^1 is $-D$ -alkylene- E -, $-D$ -alkenylene- E -, $-D$ -alkynylene- E -, $-D$ -cycloalkylene- E -,

15 $-D$ -heterocyclylene- E -, $-O-$, $-S-$, $-S(O)-$, $-S(O)_2-$, $-C(O)-$, $-N(R^{11})-$, or $-C(=N-OR^{12})-$, wherein

D and E independently of each other are a direct bond, $-O-$ or $-S-$;

R^{11} is selected from hydrogen, alkyl, aryl, carbamoyl, aryl-alkylene-,

heteroaryl-alkylene-, alkyl- $O-C(O)-$, aryl-alkylene- $O-C(O)-$,

heteroaryl-alkylene- $O-C(O)-$, alkyl- $NH-C(O)-$, aryl-alkylene- $NH-C(O)-$,

20

heteroaryl-alkylene- $NH-C(O)-$, alkyl- SO_2 , aryl-alkylene- SO_2 ,

heteroaryl-alkylene- SO_2 , aryl- SO_2 , heteroaryl- SO_2 , alkyl- $NH-SO_2$,

aryl-alkylene- $NH-SO_2$, heteroaryl-alkylene- $NH-SO_2$, alkyl- $C(O)-$,

aryl-alkylene- $C(O)-$, heteroaryl-alkylene- $C(O)-$, alkyl- Y -, aryl- Y -, heteroaryl- Y -,

aryl-alkylene- Y -, heteroaryl-alkylene- Y -, $N(R^{13})(R^{14})$ -alkylene- Y -, and

25

R^{15} - W^2 -alkylene- Y -, wherein

Y and W^2 independently of each other are a direct bond, $-CH_2-$, $-SO_2-$,

$-N(H)CO-$, $-N(H)SO_2-$, or $-O-C(O)-$;

R^{13} and R^{14} independently of each other are selected from hydrogen, aryl,

heteroaryl, C_{1-6} -alkyl, C_{1-6} -alkoxy, aryl- C_{1-6} -alkylene-,

30

heteroaryl- C_{1-6} -alkylene-, aryl- C_{1-6} -alkoxy-, heteroaryl- C_{1-6} -alkoxy-,




C₁₋₆-alkyl-arylene-, C₁₋₆-alkyl-heteroarylene-, C₁₋₆-alkoxy-heteroarylene-, or C₁₋₆-alkoxy-arylene-;

or

R₁₃ and R₁₄ may be taken together to form a ring having the formula
 5 -(CH₂)_o-X-(CH₂)_p- bonded to the nitrogen atom to which R₁₃ and R₁₄ are attached, wherein

o and p are independently of each other are 1, 2, 3, or 4; and

X is a direct bond, -CH₂-, -O-, -S-, -S(O₂)-, -C(O)-, -CON(H)-, -NHC(O)-, -NHCON(H)-, -NHSO₂-, -SO₂N(H)-, -C(O)-O-, -O-C(O)-, -NHSO₂NH-,



wherein

R¹⁸ and R¹⁷ are selected from hydrogen, aryl, heteroaryl, C₁₋₆-alkyl, C₁₋₆-alkoxy, aryl-C₁₋₆-alkylene-,

heteroaryl-C₁₋₆-alkylene-, C₁₋₆-alkyl-arylene-,

C₁₋₆-alkyl-heteroarylene-, C₁₋₆-alkoxy-arylene-,

C₁₋₆-alkoxy-heteroarylene-, heteroarylaryl-C₁₋₆-alkoxy-, or aryl-C₁₋₆-alkoxy-; and

R¹⁵ is selected from the group consisting of aryl, heteroaryl, cycloalkyl, heterocyclyl, alkyl, heteroaryl-alkylene-, or aryl-alkylene-; and

R¹² is selected from hydrogen, aryl, heteroaryl, alkyl, aryl-alkylene-, heteroaryl-alkylene-, alkyl-arylene-, alkyl-heteroarylene-, alkoxy-heteroarylene-, or alkoxy-arylene-;

G¹ is alkyl, alkenyl, alkynyl, cycloalkyl or heterocyclyl,

optionally substituted with one or more substituents selected from the group consisting of -CN, -CF₃, -OCF₃, -OR¹⁸, -NR¹⁸R¹⁹, C₃₋₁₀-cycloalkyl and C₁₋₆-alkyl, wherein

R¹⁸ and R¹⁹ independently of each other are hydrogen, C₁₋₆-alkyl,

heteroaryl-C₁₋₆-alkylene-, aryl-C₁₋₆-alkylene-, C₁₋₆-alkyl-arylene-,

5 C₁₋₆-alkyl-heteroarylene-, heteroaryl, or aryl;

or

R¹⁸ and R¹⁹, when attached to the same nitrogen atom, together with the said nitrogen atom may form a 3 to 8 membered heterocyclic ring optionally containing one or two further heteroatoms selected from nitrogen, oxygen and sulfur, and

10 optionally containing one or two double bonds;

or

G¹ is aryl, heteroaryl, fused cycloalkylheteroaryl, fused heterocyclaryl, or fused cycloalkylaryl,

optionally substituted with one or more substituents R⁴⁰, R⁴¹, and R⁴²;

15 L² is a direct bond, alkylene, alkenylene, alkynylene, -N(R²⁰)-, -alkylene-N(R²⁰)-, -alkenylene-N(R²⁰)-, -alkynylene-N(R²⁰)-, wherein

R²⁰ is hydrogen, or

R²⁰ is alkyl, alkenyl, alkynyl, cycloalkyl-W³-, heterocyclyl-W³-, aryl-W³-,

heteroaryl-W³-, optionally substituted with one or more substituents R³⁰, R³¹, and

20 R³², wherein

W³ is alkylene or a direct bond;

wherein L¹ and L² are attached to adjacent atoms in A¹;

L³ is -C(O)-, or -S(O)₂-;

R¹ is hydrogen, or

25 R¹ is alkyl, alkenyl, alkynyl, cycloalkyl-W⁴-, heterocyclyl-W⁴-, aryl-W⁴-, or heteroaryl-W⁴-, optionally substituted with one or more substituents R³³, R³⁴, and R³⁵, wherein


W⁴ is alkylene or a direct bond;

G² is heteroaryl, fused heterocyclylheteroaryl, or fused cycloalkylheteroaryl,

optionally substituted with one or more substituents R⁴³, R⁴⁴, and R⁴⁵, wherein said heteroaryl

30 group possesses a nitrogen atom adjacent to the atom joining said heteroaryl group to -N(R¹)-;

or a group of the formula



wherein

G^3 and G^5 independently of each other are alkyl, alkenyl, alkynyl, cycloalkyl- R^{22} -, heterocyclyl- R^{22} -, aryl- R^{22} -, heteroaryl- R^{22} -,

5 optionally substituted with one or more substituents R^{48} , R^{47} , and R^{48} , wherein

R^{22} is alkylene or a direct bond; and

R^{21} is hydrogen, alkyl, alkenyl, alkynyl, cycloalkyl- W^5 -, or heterocyclyl- W^5 -,

optionally substituted with one or more substituents R^{38} , R^{37} , and R^{38} , or

10 R^{21} is aryl- W^5 -, or heteroaryl- W^5 -, optionally substituted with one or more substituents R^{49} , R^{50} , and R^{51} , wherein

W^5 is alkylene or a direct bond;

wherein

R^{30} , R^{31} , R^{32} , R^{33} , R^{34} , R^{35} , R^{36} , R^{37} , and R^{38} independently of each other are selected from
 15 - CHF_2 -, - CF_3 -, - OCF_3 -, - $OCHF_2$ -, - OCH_2CF_3 -, - OCF_2CHF_2 -, - $S(O)_2CF_3$ -, - SCF_3 -, - OR^{52} -, - $NR^{52}R^{53}$ -,
 - SR^{52} -, - $NR^{52}S(O)_2R^{53}$ -, - $S(O)_2NR^{52}R^{53}$ -, - $S(O)NR^{52}R^{53}$ -, - $S(O)R^{52}$ -, - $S(O)_2R^{52}$ -, - $C(O)NR^{52}R^{53}$ -,
 - $OC(O)NR^{52}R^{53}$ -, - $NR^{52}C(O)R^{53}$ -, - $CH_2C(O)NR^{52}R^{53}$ -, - $OCH_2C(O)NR^{52}R^{53}$ -, - CH_2OR^{52} -,
 - $CH_2NR^{52}R^{53}$ -, - $OC(O)R^{52}$ -, - $C(O)R^{52}$ and - $C(O)OR^{52}$; or

C_{2-6} -alkenyl and C_{2-6} -alkynyl, which may optionally be substituted with one or more substituents selected from -CN, - CF_3 -, - OCF_3 -, - OR^{52} -, - $NR^{52}R^{53}$ and C_{1-6} -alkyl; or

20 C_{3-10} -cycloalkyl, C_{4-8} -cycloalkenyl, heterocyclyl, C_{3-10} -cycloalkyl- C_{1-6} -alkylene-,
 C_{3-10} -cycloalkyl- C_{1-6} -alkoxy-, C_{3-10} -cycloalkyloxy, C_{3-10} -cycloalkyl- C_{1-6} -alkylthio-,
 C_{3-10} -cycloalkylthio, C_{3-10} -cycloalkyl- C_{2-6} -alkenylene-, C_{3-10} -cycloalkyl- C_{2-6} -alkynylene-,
 C_{4-8} -cycloalkenyl- C_{1-6} -alkylene-, C_{4-8} -cycloalkenyl- C_{2-6} -alkenylene-,
 C_{4-8} -cycloalkenyl- C_{2-6} -alkynylene-, heterocyclyl- C_{1-6} -alkylene-, heterocyclyl- C_{2-6} -alkenylene-,

25 heterocyclyl- C_{2-6} -alkynylene-, aryl, aryloxy, aryloxycarbonyl, aroyl, aryl- C_{1-6} -alkoxy-,
 aryl- C_{1-6} -alkylene-, aryl- C_{2-6} -alkenylene-, aryl- C_{2-6} -alkynylene-, heteroaryl,
 heteroaryl- C_{1-6} -alkylene-, heteroaryl- C_{2-6} -alkenylene- and heteroaryl- C_{2-6} -alkynylene-, of

which the aryl and heteroaryl moieties optionally may be substituted with one or more substituents selected from halogen, - $C(O)OR^{52}$ -, -CN, - CF_3 -, - OCF_3 -, - NO_2 -, - OR^{52} -, - $NR^{52}R^{53}$ and

30 C_{1-6} -alkyl, wherein

R^{52} and R^{53} independently of each other are hydrogen, C_{1-6} -alkyl, aryl- C_{1-6} -alkylene-, heteroaryl- C_{1-6} -alkylene-, heteroaryl, or aryl;

or

R^{52} and R^{53} , when attached to the same nitrogen atom, together with the said nitrogen atom may form a 3 to 8 membered heterocyclic ring optionally containing one or two further heteroatoms selected from nitrogen, oxygen and sulfur, and

5 optionally containing one or two double bonds;

R^{40} , R^{41} , R^{42} , R^{43} , R^{44} , R^{45} , R^{46} , R^{47} , R^{48} , R^{49} , R^{50} and R^{51} independently of each other are halogen, -CN, -NO₂, C₁₋₆-alkyl, -CHF₂, -CF₃, -OCF₃, -OCHF₂, -OCH₂CF₃, -OCF₂CHF₂, -S(O)₂CF₃, -SCF₃, -OR⁵⁴, -NR⁵⁴R⁵⁵, -SR⁵⁴, -NR⁵⁴S(O)₂R⁵⁵, -S(O)₂NR⁵⁴R⁵⁵, -S(O)NR⁵⁴R⁵⁵, -S(O)R⁵⁴, -S(O)₂R⁵⁴, -C(O)NR⁵⁴R⁵⁵, -OC(O)NR⁵⁴R⁵⁵, -NR⁵⁴C(O)R⁵⁵, -CH₂C(O)NR⁵⁴R⁵⁵, -CH₂C(O)OR⁵⁴, -OCH₂C(O)NR⁵⁴R⁵⁵, -CH₂OR⁵⁴, -CH₂NR⁵⁴R⁵⁵, -OC(O)R⁵⁴, -C(O)R⁵⁴ and -C(O)OR⁵⁴; or

C₂₋₆-alkenyl and C₂₋₆-alkynyl, which may optionally be substituted with one or more substituents selected from -CN, -CF₃, -OCF₃, -OR⁵⁴, -NR⁵⁴R⁵⁵ and C₁₋₆-alkyl; C₃₋₁₀-cycloalkyl, C₄₋₈-cycloalkenyl, heterocyclyl, C₃₋₁₀-cycloalkyl-C₁₋₆-alkylene-, C₃₋₁₀-cycloalkyl-C₁₋₆-alkoxy-, C₃₋₁₀-cycloalkoxy, C₃₋₁₀-cycloalkyl-C₁₋₆-alkylthio-, C₃₋₁₀-cycloalkylthio, C₃₋₁₀-cycloalkyl-C₂₋₆-alkenylene-, C₃₋₁₀-cycloalkyl-C₂₋₆-alkynylene-, C₄₋₈-cycloalkenyl-C₁₋₆-alkylene-, C₄₋₈-cycloalkenyl-C₂₋₆-alkenylene-, C₄₋₈-cycloalkenyl-C₂₋₆-alkynylene-, heterocyclyl-C₁₋₆-alkylene-, heterocyclyl-C₂₋₆-alkenylene-, heterocyclyl-C₂₋₆-alkynylene-; or

15 aryl, aryloxy, aryloxycarbonyl, aroyl, aryl-C₁₋₆-alkoxy-, aryl-C₁₋₆-alkylene-, aryl-C₂₋₆-alkenylene-, aryl-C₂₋₆-alkynylene-, heteroaryl, heteroaryl-C₁₋₆-alkylene-, heteroaryl-C₂₋₆-alkenylene- and heteroaryl-C₂₋₆-alkynylene-, of which the aryl and heteroaryl moieties optionally may be substituted with one or more substituents selected from halogen, -C(O)OR⁵⁴, -CN, -CF₃, -OCF₃, -NO₂, -OR⁵⁴, -NR⁵⁴R⁵⁵ or C₁₋₆-alkyl, wherein

25 R^{54} and R^{55} independently of each other are hydrogen, C₁₋₆-alkyl, C₁₋₆-alkyl-arylene-, C₁₋₆-alkyl-heteroarylene-, aryl-C₁₋₆-alkylene-, heteroaryl-C₁₋₆-alkylene-, heteroaryl, or aryl;

or

30 R^{54} and R^{55} independently of each other are hydrogen or -(CHR⁶³)_u-(CHR⁶⁴)_v-Z, wherein

u is 1 or 2;

v is 0, 1 or 2;

R^{63} and R^{64} independently of each other are hydrogen, C₁₋₆-alkyl,

C₁₋₆-alkyl-arylene- aryl, hydroxy, hydroxyalkyl, amino, or aminoalkyl;

Z is hydrogen, $-O-R^{65}$, $-C(O)O-R^{65}$, $-CONR^{65}R^{66}$, alkylamino, or dialkylamino, wherein

R^{65} and R^{66} independently of each other is hydrogen or C_{1-6} -alkyl;

or

5 Z is a five or six membered ring wherein at least one ring atom is nitrogen and the remaining ring atoms are either carbon or oxygen;

or

R^{54} and R^{55} , when attached to the same nitrogen atom, together with the said nitrogen atom may form a 3 to 8 membered heterocyclic ring optionally containing
10 one or two further heteroatoms selected from nitrogen, oxygen and sulfur, and optionally containing one or two double bonds, and optionally substituted with one or more C_{1-6} -alkyl groups;

or a pharmaceutically acceptable salt, solvate, or prodrug thereof.

15 Other embodiments of the present invention are clear from the following list of embodiments.

Embodiment 2: A compound according to embodiment 1, wherein

A^1 is arylene or heteroarylene, optionally substituted with one or more substituents R^{23} , R^{24} , R^{25} , R^{26} , and R^{27} , wherein

20 R^{23} , R^{24} , R^{25} , R^{26} , and R^{27} independently of each other are selected from the group consisting of
halogen, $-C(O)OR^2$, $-CN$, $-CF_3$, $-OCF_3$, $-NO_2$, $-OR^2$, $-NR^2R^3$, C_{1-6} -alkyl-Z-,
 C_{2-6} -alkenyl-Z-, C_{2-6} -alkynyl-Z-, cycloalkyl-Z-, heterocyclyl-Z-, aryl-Z-, heteroaryl-Z-,
aryl- C_{1-6} -alkylene-Z-, heteroaryl- C_{1-6} -alkylene-Z-, heterocyclyl- C_{1-6} -alkylene-Z-,
cycloalkyl- C_{1-6} -alkylene-Z-, $N(R^4R^5)-C_{1-6}$ -alkylene-Z-, R^6-W^1-Z -, $R^6-W^1-N(R^4)-Z$ -, R^6-
25 $N(R^4)-Z$, and $R^6-W^1-C_{1-6}$ -alkylene-Z-, wherein

R^2 , R^3 , R^4 , R^5 , R^6 , Z, and W^1 are as defined for embodiment 1.

Embodiment 3: A compound according to embodiment 2, wherein

A^1 is C_{6-10} -arylene or C_{4-10} -heteroarylene, optionally substituted with one or more substituents R^{23} , R^{24} , R^{25} , R^{26} , and R^{27} , wherein

30 R^{23} , R^{24} , R^{25} , R^{26} , and R^{27} independently of each other are selected from the group consisting of
halogen, $-C(O)OR^2$, $-CN$, $-CF_3$, $-OCF_3$, $-NO_2$, $-OR^2$, $-NR^2R^3$, C_{1-6} -alkyl-Z-,
 C_{2-6} -alkenyl-Z-, C_{2-6} -alkynyl-Z-, cycloalkyl-Z-, heterocyclyl-Z-, aryl-Z-, heteroaryl-Z-,
aryl- C_{1-6} -alkylene-Z-, heteroaryl- C_{1-6} -alkylene-Z-, heterocyclyl- C_{1-6} -alkylene-Z-,

cycloalkyl-C₁₋₆-alkylene-Z-, N(R⁴R⁵)-C₁₋₆-alkylene-Z-, R⁶-W¹-Z-, R⁶-W¹-N(R⁴)-Z-, R⁶-N(R⁴)-Z-, and R⁶-W¹-C₁₋₆-alkylene-Z-, wherein

R², R³, R⁴, R⁵, R⁶, Z, and W¹ are as defined for embodiment 1.

Embodiment 4: A compound according to any of embodiments 1 to 3, wherein

5 R²³, R²⁴, R²⁵, R²⁶, and R²⁷ independently of each other are selected from the group consisting of

halogen, -C(O)OR², -CN, -CF₃, -OCF₃, -NO₂, -OR², -NR²R³, C₁₋₆-alkyl-Z-, cycloalkyl-Z-, heterocyclyl-Z-, aryl-Z-, or heteroaryl-Z-, N(R⁴R⁵)-C₁₋₆-alkylene-Z-, R⁶-W¹-Z-, R⁶-W¹-N(R⁴)-Z-, R⁶-N(R⁴)-Z-, and R⁶-W¹-C₁₋₆-alkylene-Z-, wherein

10 R², R³, R⁴, R⁵, R⁶, Z, and W¹ are as defined for embodiment 1.

Embodiment 5: A compound according to any of embodiments 1 to 4, wherein

R⁴, R⁵, and R⁶ independently of each other are selected from the group consisting of hydrogen, aryl, alkyl, heteroaryl-alkylene-, and aryl-alkylene-.

Embodiment 6: A compound according to embodiment 5, wherein

15 R⁴, R⁵, and R⁶ independently of each other are hydrogen or alkyl.

Embodiment 7: A compound according to embodiment 6, wherein


R⁴, R⁵, and R⁶ are hydrogen.

Embodiment 8: A compound according to any of embodiments 1 to 4, wherein

20 R⁴ and R⁵ is taken together to form a ring having the formula -(CH₂)_j-Q-(CH₂)_k- bonded to the nitrogen atom to which R⁴ and R⁵ are attached, wherein

j and k independently of each other is 1, 2, 3, or 4;

Q is a direct bond, -CH₂-, -O-, -S-, -S(O₂)-, -C(O)-, -C(O)NH-, -NHC(O)-, -NHC(O)NH-, -NHSO₂-, -SO₂NH-, -C(O)-O-, -O-C(O)-, -NHSO₂NH-,




25 wherein

R⁹ and R¹⁰ independently of each other are selected from the group consisting of hydrogen, aryl, alkyl, and arylalkyl-.

Embodiment 9: A compound according to embodiment 8, wherein

Q is a direct bond, $-\text{CH}_2-$, $-\text{O}-$, $-\text{S}-$, $-\text{S}(\text{O}_2)-$, $-\text{C}(\text{O})-$, $-\text{C}(\text{O})\text{NH}-$, $-\text{NHC}(\text{O})-$, $-\text{NHC}(\text{O})\text{NH}-$, $-\text{NHSO}_2-$, $-\text{SO}_2\text{NH}-$, $-\text{C}(\text{O})-\text{O}-$, $-\text{O}-\text{C}(\text{O})-$, $-\text{NHSO}_2\text{NH}-$,




wherein

5 R^9 and R^{10} independently of each other are hydrogen or alkyl.

Embodiment 10: A compound according to embodiment 9, wherein

Q is a direct bond, $-\text{CH}_2-$, $-\text{O}-$, $-\text{S}-$, $-\text{S}(\text{O}_2)-$, $-\text{C}(\text{O})-$, $-\text{C}(\text{O})\text{NH}-$, $-\text{NHC}(\text{O})-$, $-\text{NHC}(\text{O})\text{NH}-$, $-\text{NH}\text{SO}_2-$, $-\text{SO}_2\text{NH}-$, $-\text{C}(\text{O})-\text{O}-$, $-\text{O}-\text{C}(\text{O})-$, $-\text{NH}\text{SO}_2\text{NH}-$,



10 Embodiment 11: A compound according to embodiment 10, wherein Q is a direct bond.

Embodiment 12: A compound according to any of embodiments 1 to 11, wherein

W¹ is a direct bond, -O-, -NH-, -S-, -SO₂-, -C(O)NH-, -NHC(O)-, -N(H)CON(H)-, -N(H)SO₂-, -SO₂N(H)-, -C(O)-O-, -N(H)SO₂N(H)-, or -O-C(O)-.

Embodiment 13: A compound according to embodiment 12, wherein W^1 is a direct bond, -O-,
15 -S-, or -SO₂.

Embodiment 14: A compound according to embodiment 13, wherein W^1 is a direct bond.

Embodiment 15: A compound according to embodiment 4, wherein

at least one of R^{23} , R^{24} , R^{25} , R^{26} , and R^{27} is halogen, $-C(O)OR^2$, $-CN$, $-CF_3$, $-OCF_3$, $-NO_2$, $-OR^2$, $-NR^2R^3$, C_{1-6} -alkyl-Z-, cycloalkyl-Z-, heterocyclyl-Z-, aryl-Z-, or heteroaryl-Z-, wherein

R^2 , R^3 , and Z are as defined for embodiment 1.

Embodiment 16: A compound according to embodiment 15, wherein

at least one of R²³, R²⁴, R²⁵, R²⁶, and R²⁷ is halogen, -C(O)OR², -CN, -CF₃, -OCF₃, -NO₂, -OR², -NR²R³, C₁₋₈-alkyl-Z-, C₃₋₁₀-cycloalkyl-Z-, C₃₋₁₀-heterocyclyl-Z-, C₃₋₁₀-aryl-Z-, or C₃₋₁₀-heteroaryl-Z-, wherein

R^2 , R^3 , and Z are as defined for embodiment 1.

Embodiment 17: A compound according to embodiment 16, wherein

at least one of R^{23} , R^{24} , R^{25} , R^{26} , and R^{27} is halogen, $-C(O)OR^2$, $-CN$, $-NO_2$, $-OR^2$, $-NR^2R^3$, C_{1-6} -alkyl-Z-, C_{3-10} -cycloalkyl-Z-, C_{3-10} -heterocyclyl-Z-, C_{3-10} -aryl-Z-, or C_{3-10} -heteroaryl-Z-, wherein

R^2 , R^3 , and Z are as defined for embodiment 1.

Embodiment 18: A compound according to embodiment 17, wherein

at least one of R^{23} , R^{24} , R^{25} , R^{26} , and R^{27} is halogen, $-C(O)OR^2$, $-CN$, $-NO_2$, $-OR^2$, or $-NR^2R^3$, wherein

R^2 , R^3 , and Z are as defined for embodiment 1.

Embodiment 19: A compound according to any of embodiments 1 to 18, wherein

R^2 and R^3 independently of each other are hydrogen, C_{1-6} -alkyl, aryl- C_{1-6} -alkylene-, heteroaryl- C_{1-6} -alkylene-, C_{1-6} -alkyl-arylene-, C_{1-6} -alkyl-heteroarylene-, heteroaryl, or aryl.

Embodiment 20: A compound according to embodiment 19, wherein

R^2 and R^3 independently of each other, are hydrogen, C_{1-6} -alkyl, aryl- C_{1-6} -alkylene- or aryl.

Embodiment 21: A compound according to embodiment 20, wherein

R^2 is hydrogen or C_{1-6} -alkyl.

Embodiment 22: A compound according to embodiment 21, wherein

R^2 is hydrogen.

Embodiment 23: A compound according to any of embodiments 1 to 22, wherein

R^3 is hydrogen or C_{1-6} -alkyl.

Embodiment 24: A compound according to embodiment 23, wherein

R^3 is hydrogen.

Embodiment 25: A compound according to any of embodiments 1 to 15, wherein

R^2 and R^3 , when attached to the same nitrogen atom, together with said nitrogen atom may form a 3 to 8 membered heterocyclic ring optionally containing one or two further heteroatoms selected from nitrogen, oxygen and sulfur, and optionally containing one or two double bonds.

Embodiment 26: A compound according to any of embodiments 1 to 25, wherein


Z is a direct bond, $-O-$, $-NH-$, $-S-$, $-SO_2-$, $-C(O)NH-$, $-NHC(O)-$, $-N(H)CON(H)-$, $-N(H)SO_2-$, $-SO_2N(H)-$, $-C(O)-O-$, $-N(H)SO_2N(H)-$, or $-O-C(O)-$.

Embodiment 27: A compound according to embodiment 26, wherein

Z is a direct bond, $-O-$, $-S-$, or $-SO_2-$.

Embodiment 28: A compound according to embodiment 27, wherein
Z is a direct bond.

Embodiment 29: A compound according to embodiment 2, wherein
A¹ is



5 , wherein

R²³, R²⁴, R²⁵, and R²⁶, independently of each other, are hydrogen or as defined for
embodiment 1.

Embodiment 30: A compound according to embodiment 29, wherein

10 R²³, R²⁴, R²⁵, and R²⁶ independently of each other are selected from the group
consisting of
halogen, -C(O)OR², -CN, -CF₃, -OCF₃, -NO₂, -OR², -NR²R³, C₁₋₆-alkyl-Z-,
cycloalkyl-Z-, heterocyclyl-Z-, aryl-Z-, or heteroaryl-Z-, N(R⁴R⁵)-C₁₋₆-alkylene-Z-,
R⁶-W¹-Z-, R⁶-W¹-N(R⁴)-Z-, R⁶-N(R⁴)-Z-, and R⁶-W¹-C₁₋₆-alkylene-Z-, wherein
R², R³, R⁴, R⁵, R⁶, Z, and W¹ are as defined for embodiment 1.

15 Embodiment 31: A compound according to embodiment 29 or 30, wherein
R⁴, R⁵, and R⁶ independently of each other are selected from the group consisting of
hydrogen, aryl, alkyl, heteroaryl-alkylene-, and aryl-alkylene-.

Embodiment 32: A compound according to embodiment 31, wherein

R⁴, R⁵, and R⁶ independently of each other are hydrogen or alkyl.

20 Embodiment 33: A compound according to embodiment 32, wherein
R⁴, R⁵, and R⁶ are hydrogen.


Embodiment 34: A compound according to embodiment 29 or 30, wherein

R⁴ and R⁵ is taken together to form a ring having the formula -(CH₂)_j-Q-(CH₂)_k-
bonded to the nitrogen atom to which R⁴ and R⁵ are attached, wherein

25 j and k independently of each other is 1, 2, 3, or 4;

Q is a direct bond, -CH₂-, -O-, -S-, -S(O₂)-, -C(O)-, -C(O)NH-, -NHC(O)-,
-NHC(O)NH-, -NHSO₂-, -SO₂NH-, -C(O)-O-, -O-C(O)-, -NHSO₂NH-,

35




wherein

R^9 and R^{10} independently of each other are selected from the group consisting of hydrogen, aryl, alkyl, and arylalkyl-.

5 Embodiment 35: A compound according to embodiment 34, wherein

Q is a direct bond, $-\text{CH}_2-$, $-\text{O}-$, $-\text{S}-$, $-\text{S}(\text{O}_2)-$, $-\text{C}(\text{O})-$, $-\text{C}(\text{O})\text{NH}-$, $-\text{NHC}(\text{O})-$, $-\text{NHC}(\text{O})\text{NH}-$, $-\text{NH}\text{SO}_2-$, $-\text{SO}_2\text{NH}-$, $-\text{C}(\text{O})-\text{O}-$, $-\text{O}-\text{C}(\text{O})-$, $-\text{NH}\text{SO}_2\text{NH}-$,




wherein

10 R^9 and R^{10} independently of each other are hydrogen or alkyl.

Embodiment 36: A compound according to embodiment 35, wherein

Q is a direct bond, $-\text{CH}_2-$, $-\text{O}-$, $-\text{S}-$, $-\text{S}(\text{O}_2)-$, $-\text{C}(\text{O})-$, $-\text{C}(\text{O})\text{NH}-$, $-\text{NHC}(\text{O})-$, $-\text{NHC}(\text{O})\text{NH}-$, $-\text{NH}\text{SO}_2-$, $-\text{SO}_2\text{NH}-$, $-\text{C}(\text{O})-\text{O}-$, $-\text{O}-\text{C}(\text{O})-$, $-\text{NH}\text{SO}_2\text{NH}-$,



15 Embodiment 37: A compound according to embodiment 36, wherein

Q is a direct bond.

Embodiment 38: A compound according to any of embodiments 29 to 37, wherein

W^1 is a direct bond, $-\text{O}-$, $-\text{NH}-$, $-\text{S}-$, $-\text{SO}_2-$, $-\text{C}(\text{O})\text{NH}-$, $-\text{NHC}(\text{O})-$, $-\text{N}(\text{H})\text{CON}(\text{H})-$, $-\text{N}(\text{H})\text{SO}_2-$, $-\text{SO}_2\text{N}(\text{H})-$, $-\text{C}(\text{O})-\text{O}-$, $-\text{N}(\text{H})\text{SO}_2\text{N}(\text{H})-$, or $-\text{O}-\text{C}(\text{O})-$.

Embodiment 39: A compound according to embodiment 38, wherein

W^1 is a direct bond, -O-, -S-, or -SO₂-.

Embodiment 40: A compound according to embodiment 39, wherein

W^1 is a direct bond.

5 Embodiment 41: A compound according to embodiment 29, wherein

at least one of R^{23} , R^{24} , R^{25} , and R^{26} is halogen, -C(O)OR², -CN, -CF₃, -OCF₃, -NO₂, -OR², -NR²R³, C₁₋₆-alkyl-Z-, cycloalkyl-Z-, heterocyclyl-Z-, aryl-Z-, or heteroaryl-Z-, wherein

R^2 , R^3 , and Z are as defined for embodiment 29.

10 Embodiment 42: A compound according to embodiment 41, wherein

at least one of R^{23} , R^{24} , R^{25} , and R^{26} is halogen, -C(O)OR², -CN, -CF₃, -OCF₃, -NO₂, -OR², -NR²R³, C₁₋₆-alkyl-Z-, C₃₋₁₀-cycloalkyl-Z-, C₃₋₁₀-heterocyclyl-Z-, C₃₋₁₀-aryl-Z-, or C₃₋₁₀-heteroaryl-Z-, wherein

R^2 , R^3 , and Z are as defined for embodiment 29.

15 Embodiment 43: A compound according to embodiment 42, wherein

at least one of R^{23} , R^{24} , R^{25} , and R^{26} is halogen, -C(O)OR², -CN, -NO₂, -OR², -NR²R³, C₁₋₆-alkyl-Z-, C₃₋₁₀-cycloalkyl-Z-, C₃₋₁₀-heterocyclyl-Z-, C₃₋₁₀-aryl-Z-, or C₃₋₁₀-heteroaryl-Z-, wherein

R^2 , R^3 , and Z are as defined for embodiment 29.

20 Embodiment 44: A compound according to any of embodiments 29 to 43, wherein

Z is a direct bond, -O-, -NH-, -S-, -SO₂-, -C(O)NH-, -NHC(O)-, -N(H)CON(H)-, -N(H)SO₂-, -SO₂N(H)-, -C(O)-O-, -N(H)SO₂N(H)-, or -O-C(O)-.

Embodiment 45: A compound according to embodiment 44, wherein

Z is a direct bond, -O-, -S-, or -SO₂-.

25 Embodiment 46: A compound according to embodiment 45, wherein

Z is a direct bond.

Embodiment 47: A compound according to embodiment 43, wherein

at least one of R^{23} , R^{24} , R^{25} , and R^{26} is halogen, -C(O)OR², -CN, -NO₂, -OR², -NR²R³, or C₁₋₆-alkyl, wherein

30 R^2 , and R^3 are as defined for embodiment 29.

Embodiment 48: A compound according to any of embodiments 29 to 47, wherein

R^2 and R^3 independently of each other are hydrogen, C₁₋₆-alkyl, aryl-C₁₋₆-alkylene-, heteroaryl-C₁₋₆-alkylene-, C₁₋₆-alkyl-arylene-, C₁₋₆-alkyl-heteroarylene-, heteroaryl, or aryl.

35 Embodiment 49: A compound according to embodiment 48, wherein

R^2 and R^3 independently of each other, are hydrogen, C_{1-6} -alkyl, aryl- C_{1-6} -alkylene- or aryl.

Embodiment 50: A compound according to embodiment 49, wherein

R^2 is hydrogen or C_{1-6} -alkyl.

5 Embodiment 51: A compound according to embodiment 50, wherein

R^2 is hydrogen.

Embodiment 52: A compound according to any of embodiments 29 to 51, wherein

R^3 is hydrogen or C_{1-6} -alkyl.

Embodiment 53: A compound according to embodiment 52, wherein

10 R^3 is hydrogen.

Embodiment 54: A compound according to any of embodiments 29 to 41, wherein

R^2 and R^3 , when attached to the same nitrogen atom, together with said nitrogen atom may form a 3 to 8 membered heterocyclic ring optionally containing one or two further heteroatoms selected from nitrogen, oxygen and sulfur, and optionally
15 containing one or two double bonds.

Embodiment 55: A compound according to any of embodiments 29 to 46, wherein

at least one of R^{23} , R^{24} , R^{25} , and R^{26} are hydrogen.

Embodiment 56: A compound according to embodiment 55, wherein

at least two of R^{23} , R^{24} , R^{25} , and R^{26} are hydrogen.

20 Embodiment 57: A compound according to claim 56, wherein

R^{23} and R^{26} are hydrogen.

Embodiment 58: A compound according to embodiment 56 or claim 57, wherein

at least three of R^{23} , R^{24} , R^{25} , and R^{26} are hydrogen.

Embodiment 59: A compound according to any of embodiments 29 to 58, wherein R^{24} or R^{25}

25 is halogen.

Embodiment 60: A compound according to embodiment 59, wherein R^{24} or R^{25} is fluoro.

Embodiment 61: A compound according to any of embodiments 29 to 58, wherein R^{24} or R^{25} is C_{1-6} -alkyl.

Embodiment 62: A compound according to embodiment 59, wherein R^{24} or R^{25} is methyl.

30 Embodiment 63: A compound according to any of embodiments 29 to 58, wherein

R^{24} is hydrogen.

Embodiment 64: A compound according to any of embodiments 29 to 63, wherein

R^{25} is hydrogen.

Embodiment 65: A compound according to embodiment 29, wherein

35 R^{23} , R^{24} , R^{25} , and R^{26} are hydrogen.

Embodiment 66: A compound according to any of embodiments 1 to 65, wherein

L¹ is -D-alkylene-E-, -O-, -S-, -S(O)-, -S(O)₂-, -C(O)-, -N(R¹¹)-, or -C(=N-OR¹²),
wherein

D, E, R¹¹ and R¹² are as defined for embodiment 1.

5 Embodiment 67: A compound according to embodiment 66, wherein

L¹ is -D-alkylene-E-, -O-, -C(O)-, -N(R¹¹)-, or -C(=N-OR¹²)-, wherein

D, E, R¹¹ and R¹² are as defined for embodiment 1.

Embodiment 68: A compound according to embodiment 66, wherein

L¹ is -O-.

10 Embodiment 69: A compound according to embodiment 66, wherein

L¹ is -S-.

Embodiment 70: A compound according to embodiment 66, wherein

L¹ is -C(O)-.

Embodiment 71: A compound according to any of embodiments 1 to 67, wherein

15 D is a direct bond or -O-;

E is a direct bond or -O-; and

R¹¹ and R¹² are as defined for embodiment 1.

Embodiment 72: A compound according to embodiment 71, wherein

D is a direct bond.

20 Embodiment 73: A compound according to embodiment 71, wherein

D is -O-.

Embodiment 74: A compound according to any of embodiments 71 to 73, wherein

E is a direct bond.

Embodiment 75: A compound according to any of embodiments 71 to 73, wherein

25 E is -O-.

Embodiment 76: A compound according to any of embodiments 1 to 75, wherein

R¹¹ is selected from hydrogen, alkyl, aryl, carbamoyl, aryl-alkylene-, alkyl-NH-C(O)-, aryl-alkylene-NH-C(O)-, alkyl-SO₂-, aryl-alkylene-SO₂-, aryl-SO₂-, SO₂-, alkyl-C(O)-, aryl-alkylene-C(O)-, N(R¹³)(R¹⁴)-alkylene-Y-, and R¹⁵-W²-alkylene-Y-, wherein

30 Y and W² independently of each other are a direct bond, -CH₂-, -SO₂-, -N(H)CO-, -N(H)SO₂-, or -O-C(O)-;




R¹³ and R¹⁴ independently of each other are selected from hydrogen, aryl, C₁₋₆-alkyl, or aryl-C₁₋₆-alkylene-;

or

R^{13} and R^{14} may be taken together to form a ring having the formula $-(CH_2)_o-X-(CH_2)_p-$ bonded to the nitrogen atom to which R^{13} and R^{14} are attached, wherein

5

o and p are independently of each other are 1, 2, 3, or 4; and X is a direct bond, $-CH_2-$, $-O-$, $-S-$, $-S(O_2)-$, $-C(O)-$, $-CON(H)-$, $-NHC(O)-$, $-NHCON(H)-$, $-NHSO_2-$, $-SO_2N(H)-$, $-C(O)-O-$, $-O-C(O)-$, $-NHSO_2NH-$,



wherein

10

R^{16} and R^{17} are selected from hydrogen, aryl, heteroaryl, C_{1-6} -alkyl, C_{1-6} -alkoxy, aryl- C_{1-6} -alkylene-, heteroaryl- C_{1-6} -alkylene-, C_{1-6} -alkyl-arylene-, C_{1-6} -alkyl-heteroarylene-, C_{1-6} -alkoxy-arylene-, C_{1-6} -alkoxy-heteroarylene-, heteroarylaryl- C_{1-6} -alkoxy-, or aryl- C_{1-6} -alkoxy-; and

15

R^{15} is selected from the group consisting of aryl, heteroaryl, cycloalkyl, heterocyclyl, alkyl, or aryl-alkylene-.

Embodiment 77: A compound according to embodiment 76, wherein

20

R^{11} is selected from hydrogen, alkyl, aryl, carbamoyl, aryl-alkylene-, alkyl-NH-C(O)-, aryl-alkylene-NH-C(O)-, alkyl-SO₂-, aryl-alkylene-SO₂-, aryl-SO₂-, SO₂-, alkyl-C(O)-, aryl-alkylene-C(O)-, $N(R^{13})(R^{14})$ -alkylene-Y-, and R^{15} -W²-alkylene-Y-, wherein

Y and W^2 independently of each other are a direct bond, $-CH_2-$, $-SO_2-$, $-N(H)CO-$, $-N(H)SO_2-$, or $-O-C(O)-$;




25

R^{13} and R^{14} independently of each other are selected from hydrogen, aryl, C_{1-6} -alkyl, or aryl- C_{1-6} -alkylene-;

or

R^{13} and R^{14} may be taken together to form a ring having the formula $-(CH_2)_o-X-(CH_2)_p-$ bonded to the nitrogen atom to which R^{13} and R^{14} are attached, wherein

5 o and p are independently of each other are 1, 2, 3, or 4; and
 X is a direct bond, $-CH_2-$, $-O-$, $-S-$, $-S(O_2)-$, $-C(O)-$, $-CON(H)-$,
 $-NHC(O)-$, $-NHCON(H)-$, $-NHSO_2-$, $-SO_2N(H)-$, $-C(O)-O-$, $-O-C(O)-$,
 $-NHSO_2NH-$,



10 wherein

R^{16} and R^{17} are hydrogen; and

R^{15} is selected from the group consisting of aryl, heteroaryl, cycloalkyl, heterocyclyl, alkyl, or aryl-alkylene-.

Embodiment 78: A compound according to embodiment 77, wherein

15 R^{11} is selected from hydrogen, alkyl, aryl, carbamoyl, aryl-alkylene-, alkyl-NH-C(O)-, aryl-alkylene-NH-C(O)-, alkyl-SO₂-, aryl-alkylene-SO₂-, aryl-SO₂-, SO₂-, alkyl-C(O)-, aryl-alkylene-C(O)-, $N(R^{13})(R^{14})$ -alkylene-Y-, and R^{15} -W²-alkylene-Y-, wherein

Y and W^2 independently of each other are a direct bond, $-CH_2-$, $-SO_2-$, $-N(H)CO-$, $-N(H)SO_2-$, or $-O-C(O)-$;

20 R^{13} and R^{14} independently of each other are selected from hydrogen, aryl, C₁₋₈-alkyl, or aryl-C₁₋₈-alkylene-;

or

R^{13} and R^{14} may be taken together to form a ring having the formula $-(CH_2)_o-X-(CH_2)_p-$ bonded to the nitrogen atom to which R^{13} and R^{14} are attached, wherein

25

o and p are independently of each other are 1, 2, 3, or 4;

X is a direct bond; and

R¹⁵ is selected from the group consisting of aryl, heteroaryl, cycloalkyl, heterocyclyl, alkyl, or aryl-alkylene-.

5 Embodiment 79: A compound according to embodiment 78, wherein

R¹¹ is selected from hydrogen, alkyl, aryl, carbamoyl, aryl-alkylene-, alkyl-NH-C(O)-, aryl-alkylene-NH-C(O)-, alkyl-SO₂-, aryl-alkylene-SO₂-, aryl-SO₂-, SO₂-, alkyl-C(O)-, aryl-alkylene-C(O)-, N(R¹³)(R¹⁴)-alkylene-Y-, and R¹⁵-W²-alkylene-Y-, wherein

10 Y and W² independently of each other are a direct bond, -CH₂-, -SO₂-, -N(H)CO-, -N(H)SO₂-, or -O-C(O)-;

R¹³ and R¹⁴ independently of each other are selected from hydrogen, aryl, C₁₋₆-alkyl, or aryl-C₁₋₆-alkylene-; and

R¹⁵ is selected from the group consisting of aryl, heteroaryl, cycloalkyl, heterocyclyl, alkyl, or aryl-alkylene-.

15 Embodiment 80: A compound according to embodiment 79, wherein

R¹¹ is selected from hydrogen, alkyl, aryl, carbamoyl, aryl-alkylene-, alkyl-NH-C(O)-, aryl-alkylene-NH-C(O)-, alkyl-SO₂-, aryl-alkylene-SO₂-, aryl-SO₂-, SO₂-, alkyl-C(O)-, aryl-alkylene-C(O)-, N(R¹³)(R¹⁴)-alkylene-Y-, and R¹⁵-W²-alkylene-Y-, wherein

20 Y and W² independently of each other are a direct bond, -CH₂-, -SO₂-, -N(H)CO-, -N(H)SO₂-, or -O-C(O)-;

R¹³ and R¹⁴ are hydrogen; and

R¹⁵ is selected from the group consisting of aryl, heteroaryl, cycloalkyl, heterocyclyl, alkyl, or aryl-alkylene-.

Embodiment 81: A compound according to embodiment 80, wherein

25 R¹¹ is selected from hydrogen, alkyl, aryl, carbamoyl, aryl-alkylene-, alkyl-NH-C(O)-, aryl-alkylene-NH-C(O)-, alkyl-SO₂-, aryl-alkylene-SO₂-, aryl-SO₂-, SO₂-, alkyl-C(O)-, aryl-alkylene-C(O)-, N(R¹³)(R¹⁴)-alkylene-Y-, and R¹⁵-W²-alkylene-Y-, wherein

Y is a direct bond;

W² is a direct bond, -CH₂-, -SO₂-, -N(H)CO-, -N(H)SO₂-, or -O-C(O)-;

30 R¹³ and R¹⁴ are hydrogen; and

R¹⁵ is selected from the group consisting of aryl, heteroaryl, cycloalkyl, heterocyclyl, alkyl, or aryl-alkylene-.

Embodiment 82: A compound according to embodiment 80, wherein

R¹¹ is selected from hydrogen, alkyl, aryl, carbamoyl, aryl-alkylene-, alkyl-NH-C(O)-, aryl-alkylene-NH-C(O)-, alkyl-SO₂-, aryl-alkylene-SO₂-, aryl-SO₂-, SO₂-, alkyl-C(O)-, aryl-alkylene-C(O)-, N(R¹³)(R¹⁴)-alkylene-Y-, and R¹⁵-W²-alkylene-Y-, wherein

Y is a direct bond, -CH₂-, -SO₂-, -N(H)CO-, -N(H)SO₂-, or -O-C(O)-;

5 W² is a direct bond;

R¹³ and R¹⁴ are hydrogen; and

R¹⁵ is selected from the group consisting of aryl, heteroaryl, cycloalkyl, heterocyclyl, alkyl, or aryl-alkylene-.

Embodiment 83: A compound according to any of embodiments 80 to 82, wherein

10 R¹¹ is selected from hydrogen, alkyl, aryl, carbamoyl, aryl-alkylene-, alkyl-NH-C(O)-, aryl-alkylene-NH-C(O)-, alkyl-SO₂-, aryl-alkylene-SO₂-, aryl-SO₂-, SO₂-, alkyl-C(O)-, aryl-alkylene-C(O)-, NH₂-alkylene-, and R¹⁵-alkylene-, wherein

R¹⁵ is selected from the group consisting of aryl, heteroaryl, cycloalkyl, heterocyclyl, alkyl, or aryl-alkylene-.

15 Embodiment 84: A compound according to embodiment 83, wherein

R¹¹ is selected from hydrogen, alkyl, aryl, carbamoyl, aryl-alkylene-, alkyl-NH-C(O)-, aryl-alkylene-NH-C(O)-, alkyl-SO₂-, aryl-alkylene-SO₂-, aryl-SO₂-, SO₂-, alkyl-C(O)-, aryl-alkylene-C(O)-, and NH₂-alkylene-.

Embodiment 85: A compound according to embodiment 84, wherein

20 R¹¹ is hydrogen or alkyl.

Embodiment 86: A compound according to embodiment 85, wherein

R¹¹ is hydrogen.

Embodiment 87: A compound according to any of embodiments 1 to 86, wherein

R¹² is hydrogen or alkyl.

25 Embodiment 88: A compound according to embodiment 87, wherein

R¹² is hydrogen.

Embodiment 89: A compound according to any of embodiments 1 to 88, wherein

G¹ is alkyl, alkenyl, alkynyl, cycloalkyl or heterocyclyl, optionally substituted with one or more substituents selected from the group consisting of -CN, -CF₃, -OCF₃, -OR¹⁸, -NR¹⁸R¹⁹, C₃₋₁₀-cycloalkyl and C₁₋₆-alkyl, wherein

30

R¹⁸ and R¹⁹ are as defined for embodiment 1.

Embodiment 90: A compound according to embodiment 89, wherein

G¹ is alkyl, alkenyl, alkynyl, cycloalkyl or heterocyclyl, optionally substituted with one or more substituents selected from the group consisting of -CN, -CF₃, -OCF₃, -OR¹⁸,

35 -NR¹⁸R¹⁹ and C₁₋₆-alkyl, wherein

R^{18} and R^{19} are as defined for embodiment 1.

Embodiment 91: A compound according to embodiment 89, wherein

5 G^1 is C_{1-6} -alkyl, C_{2-6} -alkenyl, C_{2-6} -alkynyl, C_{3-10} -cycloalkyl or C_{3-10} -heterocyclyl, optionally substituted with one or more substituents selected from the group consisting of -CN, -CF₃, -OCF₃, -OR¹⁸, -NR¹⁸R¹⁹, C_{3-10} -cycloalkyl and C_{1-6} -alkyl, wherein

R^{18} and R^{19} are as defined for embodiment 1.

Embodiment 92: A compound according to embodiment 90 or embodiment 91, wherein

10 G^1 is C_{1-6} -alkyl, C_{2-6} -alkenyl, C_{2-6} -alkynyl, C_{3-10} -cycloalkyl or C_{3-10} -heterocyclyl, optionally substituted with one or more substituents selected from the group consisting of -CN, -CF₃, -OCF₃, -OR¹⁸, -NR¹⁸R¹⁹ and C_{1-6} -alkyl, wherein

R^{18} and R^{19} are as defined for embodiment 1.

Embodiment 93: A compound according to any of embodiments 89 to 92, wherein

15 R^{18} and R^{19} , independently of each other, are hydrogen, C_{1-6} -alkyl, heteroaryl- C_{1-6} -alkylene-, aryl- C_{1-6} -alkylene-, C_{1-6} -alkyl-arylene-, C_{1-6} -alkyl-heteroarylene-, heteroaryl, or aryl.

Embodiment 94: A compound according to embodiment 93, wherein

20 R^{18} and R^{19} , independently of each other, are hydrogen, C_{1-6} -alkyl, C_{3-10} -heteroaryl- C_{1-6} -alkylene-, C_{3-10} -aryl- C_{1-6} -alkylene-, C_{1-6} -alkyl- C_{3-10} -arylene-, C_{1-6} -alkyl- C_{3-10} -heteroarylene-, C_{3-10} -heteroaryl, or C_{3-10} -aryl.

Embodiment 95: A compound according to embodiment 94, wherein

R^{18} and R^{19} , independently of each other, are hydrogen or C_{1-6} -alkyl.

Embodiment 96: A compound according to embodiment 95, wherein

R^{18} is hydrogen.

25 Embodiment 97: A compound according to embodiment 95 or 96, wherein

R^{19} is hydrogen.

Embodiment 98: A compound according to embodiment 90 or embodiment 92, wherein

30 R^{18} and R^{19} , when attached to the same nitrogen atom, together with the said nitrogen atom forms a 3 to 8 membered heterocyclic ring optionally containing one or two further heteroatoms selected from nitrogen, oxygen and sulfur, and optionally containing one or two double bonds.

Embodiment 99: A compound according to any of embodiments 1 to 88, wherein

35 G^1 is alkyl or cycloalkyl, optionally substituted with one or more substituents selected from the group consisting of -CN, -CF₃, -OCF₃, -OR¹⁸, -NR¹⁸R¹⁹ and C_{1-6} -alkyl,

or G¹ is aryl optionally substituted with one or more substituents R⁴⁰, R⁴¹, and R⁴², wherein R¹⁸, R¹⁹, R⁴⁰, R⁴¹, and R⁴² are as defined for embodiment 1.

Embodiment 100: A compound according to embodiment 99, wherein

5 G¹ is C₁₋₆-alkyl or C₃₋₁₀-cycloalkyl, optionally substituted with one or more substituents selected from the group consisting of -CN, -CF₃, -OCF₃, -OR¹⁸, -NR¹⁸R¹⁹ and C₁₋₆-alkyl, or G¹ is C₃₋₁₀-aryl optionally substituted with one or more substituents R⁴⁰, R⁴¹, and R⁴², wherein R¹⁸, R¹⁹, R⁴⁰, R⁴¹, and R⁴² are as defined for embodiment 1.

Embodiment 101: A compound according to embodiment 99 or embodiment 100, wherein

10 R¹⁸ and R¹⁹, independently of each other, are hydrogen, C₁₋₆-alkyl, heteroaryl-C₁₋₆-alkylene-, aryl-C₁₋₆-alkylene-, C₁₋₆-alkyl-arylene-, C₁₋₆-alkyl-heteroarylene-, heteroaryl, or aryl.

Embodiment 102: A compound according to embodiment 101, wherein

15 R¹⁸ and R¹⁹, independently of each other, are hydrogen, C₁₋₆-alkyl, C₃₋₁₀-heteroaryl-C₁₋₆-alkylene-, C₃₋₁₀-aryl-C₁₋₆-alkylene-, C₁₋₆-alkyl-C₃₋₁₀-arylene-, C₁₋₆-alkyl-C₃₋₁₀-heteroarylene-, C₃₋₁₀-heteroaryl, or C₃₋₁₀-aryl.

Embodiment 103: A compound according to embodiment 102, wherein

R¹⁸ and R¹⁹, independently of each other, are hydrogen or C₁₋₆-alkyl.

Embodiment 104: A compound according to embodiment 103, wherein

20 R¹⁸ is hydrogen.

Embodiment 105: A compound according to embodiment 103 or 104, wherein

R¹⁹ is hydrogen.

Embodiment 106: A compound according to embodiment 99 or embodiment 100, wherein

25 R¹⁸ and R¹⁹, when attached to the same nitrogen atom, together with the said nitrogen atom forms a 3 to 8 membered heterocyclic ring optionally containing one or two further heteroatoms selected from nitrogen, oxygen and sulfur, and optionally containing one or two double bonds.

Embodiment 107: A compound according to any of embodiments 99 to 106, wherein

30 R⁴⁰, R⁴¹, and R⁴² independently of each other are halogen, -CN, -NO₂, C₁₋₆-alkyl, -CHF₂, -CF₃, -OR⁵⁴, -NR⁵⁴R⁵⁵, -SR⁵⁴, -NR⁵⁴S(O)₂R⁵⁵, -S(O)₂NR⁵⁴R⁵⁵, -S(O)NR⁵⁴R⁵⁵, -S(O)R⁵⁴, -S(O)₂R⁵⁴, -C(O)NR⁵⁴R⁵⁵, -OC(O)NR⁵⁴R⁵⁵, -NR⁵⁴C(O)R⁵⁵, -CH₂C(O)NR⁵⁴R⁵⁵, -CH₂C(O)OR⁵⁴, -OCH₂C(O)NR⁵⁴R⁵⁵, -CH₂OR⁵⁴, -CH₂NR⁵⁴R⁵⁵, and -C(O)OR⁵⁴; or

C₂₋₆-alkenyl and C₂₋₆-alkynyl, which may optionally be substituted with one or more substituents selected from -CN, -CF₃, -OCF₃, -OR⁵⁴, -NR⁵⁴R⁵⁵ and C₁₋₆-alkyl, wherein

- 5 R⁵⁴ and R⁵⁵ independently of each other are hydrogen, C₁₋₆-alkyl, C₁₋₆-alkyl-arylene-, C₁₋₆-alkyl-heteroarylene-, aryl-C₁₋₆-alkylene-, heteroaryl-C₁₋₆-alkylene-, heteroaryl, or aryl;
or
R⁵⁴ and R⁵⁵, when attached to the same nitrogen atom, together with the said nitrogen atom may form a 3 to 8 membered heterocyclic ring optionally
10 containing one or two further heteroatoms selected from nitrogen, oxygen and sulfur, and optionally containing one or two double bonds.

Embodiment 108: A compound according to embodiment 107, wherein

- R⁵⁴ and R⁵⁵ independently of each other are hydrogen, or C₁₋₆-alkyl, C₁₋₆-alkyl-C₃₋₁₀-arylene-, C₁₋₆-alkyl-C₃₋₁₀-heteroarylene-, C₃₋₁₀-aryl-C₁₋₆-alkylene-,
15 C₃₋₁₀-heteroaryl-C₁₋₆-alkylene-, C₃₋₁₀-heteroaryl, or C₃₋₁₀-aryl.

Embodiment 109: A compound according to embodiment 108, wherein

 R⁵⁴ and R⁵⁵ independently of each other are hydrogen or C₁₋₆-alkyl.

Embodiment 110: A compound according to embodiment 109, wherein

 R⁵⁴ is hydrogen.

- 20 Embodiment 111: A compound according to embodiment 109 or embodiment 110, wherein R⁵⁵ is hydrogen.

Embodiment 112: A compound according to any of embodiments 1 to 88, wherein

 G¹ is aryl or heteroaryl, optionally substituted with one or more substituents R⁴⁰, R⁴¹, and R⁴², wherein

- 25 R⁴⁰, R⁴¹, and R⁴² are as defined for embodiment 1.

Embodiment 113: A compound according to embodiment 112, wherein

 G¹ is C₃₋₁₀-aryl or C₃₋₁₀-heteroaryl, optionally substituted with one or more substituents R⁴⁰, R⁴¹, and R⁴², wherein

 R⁴⁰, R⁴¹, and R⁴² are as defined for embodiment 1.

- 30 Embodiment 114: A compound according to embodiment 112, wherein

 G¹ is aryl, optionally substituted with one or more substituents R⁴⁰, R⁴¹, and R⁴², wherein

 R⁴⁰, R⁴¹, and R⁴² are as defined for embodiment 1.

Embodiment 115: A compound according to embodiment 114, wherein

G¹ is C₃₋₁₀-aryl, optionally substituted with one or more substituents R⁴⁰, R⁴¹, and R⁴², wherein

R⁴⁰, R⁴¹, and R⁴² are as defined for embodiment 1.

Embodiment 116: A compound according to any of embodiments 112 to 115, wherein

- 5 R⁴⁰, R⁴¹, and R⁴² independently of each other are
 halogen, -CN, -NO₂, C₁₋₆-alkyl, -CHF₂, -CF₃, -OR⁵⁴, -NR⁵⁴R⁵⁵, -SR⁵⁴, -NR⁵⁴S(O)₂R⁵⁵,
 -S(O)₂NR⁵⁴R⁵⁵, -S(O)NR⁵⁴R⁵⁵, -S(O)R⁵⁴, -S(O)₂R⁵⁴, -C(O)NR⁵⁴R⁵⁵, -OC(O)NR⁵⁴R⁵⁵,
 -NR⁵⁴C(O)R⁵⁵, -CH₂C(O)NR⁵⁴R⁵⁵, -CH₂C(O)OR⁵⁴, -OCH₂C(O)NR⁵⁴R⁵⁵, -CH₂OR⁵⁴,
 -CH₂NR⁵⁴R⁵⁵, and -C(O)OR⁵⁴; or
 10 C₂₋₈-alkenyl and C₂₋₈-alkynyl, which may optionally be substituted with one or more
 substituents selected from -CN, -CF₃, -OCF₃, -OR⁵⁴, -NR⁵⁴R⁵⁵ and C₁₋₆-alkyl;
 wherein
 R⁵⁴ and R⁵⁵ independently of each other are hydrogen, C₁₋₆-alkyl,
 C₁₋₆-alkyl-arylene-, C₁₋₆-alkyl-heteroarylene-, aryl-C₁₋₆-alkylene-,
 15 heteroaryl-C₁₋₆-alkylene-, heteroaryl, or aryl;
 or
 R⁵⁴ and R⁵⁵, when attached to the same nitrogen atom, together with the
 said nitrogen atom may form a 3 to 8 membered heterocyclic ring optionally
 containing one or two further heteroatoms selected from nitrogen, oxygen
 20 and sulfur, and optionally containing one or two double bonds.

Embodiment 117: A compound according to embodiment 116, wherein

R⁵⁴ and R⁵⁵ independently of each other are hydrogen, or C₁₋₆-alkyl,
 C₁₋₆-alkyl-C₃₋₁₀-arylene-, C₁₋₆-alkyl-C₃₋₁₀-heteroarylene-, C₃₋₁₀-aryl-C₁₋₆-alkylene-,
 C₃₋₁₀-heteroaryl-C₁₋₆-alkylene-, C₃₋₁₀-heteroaryl, or C₃₋₁₀-aryl.

25 Embodiment 118: A compound according to embodiment 117, wherein

R⁵⁴ and R⁵⁵ independently of each other are hydrogen or C₁₋₆-alkyl.

Embodiment 119: A compound according to embodiment 118, wherein


R⁵⁴ is hydrogen.

Embodiment 120: A compound according to embodiment 118 or embodiment 119, wherein

30 R⁵⁵ is hydrogen.

Embodiment 121: A compound according to embodiment 114, wherein

G¹ is



, wherein

- R⁵⁶, R⁵⁷, R⁵⁸, R⁵⁹, and R⁶⁰, independently of each other, are hydrogen or halogen, -CN, -NO₂, C₁₋₆-alkyl, -CHF₂, -CF₃, -OCF₃, -OCHF₂, -OCH₂CF₃, -OCF₂CHF₂, -S(O)₂CF₃, -SCF₃, -OR⁶¹, -NR⁶¹R⁶², -SR⁶¹, -NR⁶¹S(O)₂R⁶², -S(O)₂NR⁶¹R⁶², -S(O)NR⁶¹R⁶², -S(O)R⁶¹, -S(O)₂R⁶¹, -C(O)NR⁶¹R⁶², -OC(O)NR⁶¹R⁶², -NR⁶¹C(O)R⁶², -CH₂C(O)NR⁶¹R⁶², -CH₂C(O)OR⁶¹, -OCH₂C(O)NR⁶¹R⁶², -CH₂OR⁶¹, -CH₂NR⁶¹R⁶², -OC(O)R⁶¹, -C(O)R⁶¹ and -C(O)OR⁶¹; or C₂₋₆-alkenyl and C₂₋₆-alkynyl, which may optionally be substituted with one or more substituents selected from -CN, -CF₃, -OCF₃, -OR⁶¹, -NR⁶¹R⁶² and C₁₋₆-alkyl; C₃₋₁₀-cycloalkyl, C₄₋₈-cycloalkenyl, heterocyclyl, C₃₋₁₀-cycloalkyl-C₁₋₆-alkylene-, C₃₋₁₀-cycloalkyl-C₁₋₆-alkoxy-, C₃₋₁₀-cycloalkyloxy, C₃₋₁₀-cycloalkyl-C₁₋₆-alkylthio-, C₃₋₁₀-cycloalkylthio-, C₃₋₁₀-cycloalkyl-C₂₋₆-alkenylene-, C₃₋₁₀-cycloalkyl-C₂₋₆-alkynylene-, C₄₋₈-cycloalkenyl-C₁₋₆-alkylene-, C₄₋₈-cycloalkenyl-C₂₋₆-alkenylene-, C₄₋₈-cycloalkenyl-C₂₋₆-alkynylene-, heterocyclyl-C₁₋₆-alkylene-, heterocyclyl-C₂₋₆-alkenylene-, heterocyclyl-C₂₋₆-alkynylene-; or aryl, aryloxy, aryloxycarbonyl, aroyl, aryl-C₁₋₆-alkoxy-, aryl-C₁₋₆-alkylene-, aryl-C₂₋₆-alkenylene-, aryl-C₂₋₆-alkynylene-, heteroaryl, heteroaryl-C₁₋₆-alkylene-, heteroaryl-C₂₋₆-alkenylene- and heteroaryl-C₂₋₆-alkynylene-, of which the aryl and heteroaryl moieties optionally may be substituted with one or more substituents selected from halogen, -C(O)OR⁶¹, -CN, -CF₃, -OCF₃, -NO₂, -OR⁶¹, -NR⁶¹R⁶² or C₁₋₆-alkyl, wherein
- R⁶¹ and R⁶² independently of each other are hydrogen, C₁₋₆-alkyl, C₁₋₆-alkyl-arylene-, C₁₋₆-alkyl-heteroarylene-, aryl-C₁₋₆-alkylene-, heteroaryl-C₁₋₆-alkylene-, heteroaryl, or aryl;
- or
- R⁶¹ and R⁶², when attached to the same nitrogen atom, together with the said nitrogen atom may form a 3 to 8 membered heterocyclic ring optionally containing one or two further heteroatoms selected from nitrogen, oxygen and sulfur, and optionally containing one or two double bonds.

Embodiment 122: A compound according to embodiment 121, wherein

- R⁵⁶, R⁵⁷, R⁵⁸, R⁵⁹, and R⁶⁰ independently of each other are halogen, -CN, -NO₂, C₁₋₆-alkyl, -CHF₂, -CF₃, -OR⁶¹, -NR⁶¹R⁶², -SR⁶¹, -NR⁶¹S(O)₂R⁶², -S(O)₂NR⁶¹R⁶², -S(O)NR⁶¹R⁶², -S(O)R⁶¹, -S(O)₂R⁶¹, -C(O)NR⁶¹R⁶², -OC(O)NR⁶¹R⁶²,

$-\text{NR}^{61}\text{C}(\text{O})\text{R}^{62}$, $-\text{CH}_2\text{C}(\text{O})\text{NR}^{61}\text{R}^{62}$, $-\text{CH}_2\text{C}(\text{O})\text{OR}^{61}$, $-\text{OCH}_2\text{C}(\text{O})\text{NR}^{61}\text{R}^{62}$, $-\text{CH}_2\text{OR}^{61}$, $-\text{CH}_2\text{NR}^{61}\text{R}^{62}$, and $-\text{C}(\text{O})\text{OR}^{61}$; or

C_{2-6} -alkenyl and C_{2-6} -alkynyl, which may optionally be substituted with one or more substituents selected from $-\text{CN}$, $-\text{CF}_3$, $-\text{OCF}_3$, $-\text{OR}^{61}$, $-\text{NR}^{61}\text{R}^{62}$ and C_{1-6} -alkyl,

5 wherein

R^{61} and R^{62} independently of each other are hydrogen, C_{1-6} -alkyl, C_{1-6} -alkyl-arylene-, C_{1-6} -alkyl-heteroarylene-, aryl- C_{1-6} -alkylene-, heteroaryl- C_{1-6} -alkylene-, heteroaryl, or aryl;

or

10 R^{61} and R^{62} , when attached to the same nitrogen atom, together with the said nitrogen atom may form a 3 to 8 membered heterocyclic ring optionally containing one or two further heteroatoms selected from nitrogen, oxygen and sulfur, and optionally containing one or two double bonds.

Embodiment 123: A compound according to embodiment 121 or embodiment 122, wherein

15 R^{24} or R^{25} is $-\text{OR}^{61}$.

Embodiment 124: A compound according to embodiment 122, wherein

R^{61} and R^{62} independently of each other are hydrogen, or C_{1-6} -alkyl, C_{1-6} -alkyl- C_{3-10} -arylene-, C_{1-6} -alkyl- C_{3-10} -heteroarylene-, C_{3-10} -aryl- C_{1-6} -alkylene-, C_{3-10} -heteroaryl- C_{1-6} -alkylene-, C_{3-10} -heteroaryl, or C_{3-10} -aryl.

20 Embodiment 125: A compound according to embodiment 124, wherein

R^{61} and R^{62} independently of each other are hydrogen or C_{1-6} -alkyl.

Embodiment 126: A compound according to embodiment 125, wherein

R^{61} is hydrogen.

Embodiment 127: A compound according to embodiment 125 or embodiment 126; wherein

25 R^{62} is hydrogen.

Embodiment 128: A compound according to any of embodiments 121 to 127, wherein at least one of R^{56} , R^{57} , R^{58} , R^{59} , and R^{60} are hydrogen.

Embodiment 129: A compound according to embodiment 128, wherein

at least two of R^{23} , R^{24} , R^{25} , and R^{26} are hydrogen.

30 Embodiment 130: A compound according to embodiment 129, wherein

at least three of R^{23} , R^{24} , R^{25} , and R^{26} are hydrogen.

Embodiment 131: A compound according to any of embodiments 1 to 130, wherein

L^2 is a direct bond, C_{1-6} -alkylene, C_{2-6} -alkenylene, C_{2-6} -alkynylene, $-\text{N}-\text{R}^{20}$ -, $-\text{C}_{1-6}$ -alkylene- $\text{N}(\text{R}^{20})$ -, $-\text{C}_{2-6}$ -alkenylene- $\text{N}(\text{R}^{20})$ -, or $-\text{C}_{2-6}$ -alkynylene- $\text{N}(\text{R}^{20})$ -, wherein

35 R^{20} is as defined for embodiment 1.

Embodiment 132: A compound according to any of embodiments 1 to 127, wherein

L^2 is $-N-R^{20}-$, $-alkylene-N(R^{20})-$, $-alkenylene-N(R^{20})-$, or $-alkynylene-N(R^{20})-$, wherein R^{20} is as defined for embodiment 1.

Embodiment 133: A compound according to embodiment 131 or embodiment 132, wherein

5 L^2 is $-N-R^{20}-$, $-C_{1-6}-alkylene-N(R^{20})-$, $-C_{2-6}-alkenylene-N(R^{20})-$, or $-C_{2-6}-alkynylene-N(R^{20})-$, wherein R^{20} is as defined for embodiment 1.

Embodiment 134: A compound according to embodiment 133, wherein

10 L^2 is $-N-R^{20}-$, wherein R^{20} is as defined for embodiment 1.

Embodiment 135: A compound according to embodiment 131, wherein

L^2 is a direct bond.

Embodiment 136: A compound according to any of embodiments 1 to 134, wherein

15 R^{20} is hydrogen, or R^{20} is $C_{1-6}-alkyl$, $C_{2-6}-alkenyl$, $C_{2-6}-alkynyl$, $C_{3-10}-cycloalkyl-W^3-$, $C_{3-10}-heterocyclyl-W^3-$, $C_{3-10}-aryl-W^3-$, or $C_{4-10}-heteroaryl-W^3-$, optionally substituted with one or more substituents R^{30} , R^{31} , and R^{32} , wherein W^3 , R^{30} , R^{31} , and R^{32} are as defined for embodiment 1.

Embodiment 137: A compound according to embodiment 136, wherein

20 W^3 is alkylene.

Embodiment 138: A compound according to embodiment 137, wherein

W^3 is $C_{2-6}-alkylene$.

Embodiment 139: A compound according to embodiment 136, wherein

W^3 is a direct bond.

25 Embodiment 140: A compound according to any of embodiments 1 to 134, wherein

R^{20} is hydrogen, alkyl, alkenyl, or alkynyl, optionally substituted with one or more substituents R^{30} , R^{31} , and R^{32} , wherein R^{30} , R^{31} , and R^{32} are as defined for embodiment 1.

Embodiment 141: A compound according to any of embodiments 136 to 140, wherein

30 R^{20} is hydrogen, or R^{20} is $C_{1-6}-alkyl$, $C_{1-6}-alkenyl$, or $C_{1-6}-alkynyl$, optionally substituted with one or more substituents R^{30} , R^{31} , and R^{32} , wherein R^{30} , R^{31} , and R^{32} are as defined for embodiment 1.

Embodiment 142: A compound according to any of embodiments 1 to 141, wherein

R^{30} , R^{31} , and R^{32} independently of each other are selected from $-\text{CHF}_2$, $-\text{CF}_3$, $-\text{OCF}_3$, $-\text{OCHF}_2$, $-\text{OCH}_2\text{CF}_3$, $-\text{OCF}_2\text{CHF}_2$, $-\text{S}(\text{O})_2\text{CF}_3$, $-\text{SCF}_3$, $-\text{OR}^{52}$, $-\text{NR}^{52}\text{R}^{53}$, $-\text{SR}^{52}$, $-\text{NR}^{52}\text{S}(\text{O})_2\text{R}^{53}$, $-\text{S}(\text{O})_2\text{NR}^{52}\text{R}^{53}$, $-\text{S}(\text{O})\text{NR}^{52}\text{R}^{53}$, $-\text{S}(\text{O})\text{R}^{52}$, $-\text{S}(\text{O})_2\text{R}^{52}$, $-\text{C}(\text{O})\text{NR}^{52}\text{R}^{53}$, $-\text{OC}(\text{O})\text{NR}^{52}\text{R}^{53}$, $-\text{NR}^{52}\text{C}(\text{O})\text{R}^{53}$, $-\text{CH}_2\text{C}(\text{O})\text{NR}^{52}\text{R}^{53}$, $-\text{OCH}_2\text{C}(\text{O})\text{NR}^{52}\text{R}^{53}$, $-\text{CH}_2\text{OR}^{52}$, $-\text{CH}_2\text{NR}^{52}\text{R}^{53}$, $-\text{OC}(\text{O})\text{R}^{52}$, $-\text{C}(\text{O})\text{R}^{52}$ and $-\text{C}(\text{O})\text{OR}^{52}$, wherein

R^{52} and R^{53} are as defined for embodiment 1.

Embodiment 143: A compound according to embodiment 142, wherein

R^{52} and R^{53} , when attached to the same nitrogen atom, together with the said nitrogen atom may form a 3 to 8 membered heterocyclic ring optionally containing one or two further heteroatoms selected from nitrogen, oxygen and sulfur, and optionally containing one or two double bonds.

Embodiment 144: A compound according to any of embodiments 1 to 142, wherein

R^{52} and R^{53} independently of each other are hydrogen, C_{1-6} -alkyl, aryl- C_{1-6} -alkylene-heteroaryl- C_{1-6} -alkylene-heteroaryl, or aryl.

Embodiment 145: A compound according to embodiment 144, wherein

R^{52} and R^{53} independently of each other are hydrogen, C_{1-6} -alkyl, aryl- C_{1-6} -alkylene-or aryl.

Embodiment 146: A compound according to embodiment 145, wherein

R^{52} and R^{53} independently of each other are hydrogen or C_{1-6} -alkyl.

Embodiment 147: A compound according to embodiment 146, wherein

R^{52} is hydrogen.

Embodiment 148: A compound according to embodiment 146 or embodiment 147, wherein

R^{53} is hydrogen.

Embodiment 149: A compound according to embodiment 141, wherein

R^{20} is hydrogen.

Embodiment 150: A compound according to any of embodiments 1. to 149, wherein

L^3 is $-\text{C}(\text{O})-$.

Embodiment 151: A compound according to any of embodiments 1 to 150, wherein

R^1 is hydrogen, or

R^1 is C_{1-6} -alkyl, C_{2-6} -alkenyl, C_{2-6} -alkynyl, C_{3-10} -cycloalkyl- W^4 -, C_{3-10} -heterocyclyl- W^4 -, C_{3-10} -aryl- W^4 -, or C_{4-10} -heteroaryl- W^4 -, optionally substituted with one or more substituents R^{33} , R^{34} , and R^{35} , wherein

W^4 , R^{33} , R^{34} , and R^{35} are as defined for embodiment 1.

Embodiment 152: A compound according to embodiment 151, wherein

W^4 is alkylene.

Embodiment 153: A compound according to embodiment 152, wherein

W^4 is C_{2-6} -alkylene.

Embodiment 154: A compound according to embodiment 151, wherein

W^4 is a direct bond.

5 Embodiment 155: A compound according to any of embodiments 1 to 150, wherein

R^1 is hydrogen, alkyl, alkenyl, or alkynyl, optionally substituted with one or more substituents R^{33} , R^{34} , and R^{35} , wherein

R^{33} , R^{34} , and R^{35} are as defined for embodiment 1.

Embodiment 156: A compound according to any of embodiments 151 to 155, wherein

10 R^1 is hydrogen, C_{1-6} -alkyl, C_{2-6} -alkenyl, or C_{2-6} -alkynyl, optionally substituted with one or more substituents R^{33} , R^{34} , and R^{35} , wherein

R^{33} , R^{34} , and R^{35} are as defined for embodiment 1.

Embodiment 157: A compound according to embodiment 156, wherein

15 R^1 is hydrogen or C_{1-6} -alkyl optionally substituted with one or more substituents R^{33} , R^{34} , and R^{35} , wherein

R^{33} , R^{34} , and R^{35} are as defined for embodiment 1.

Embodiment 158: A compound according to any of embodiments 1 to 157, wherein

20 R^{33} , R^{34} , and R^{35} independently of each other are selected from $-CHF_2$, $-CF_3$, $-OCF_3$, $-OCHF_2$, $-OCH_2CF_3$, $-OCF_2CHF_2$, $-S(O)_2CF_3$, $-SCF_3$, $-OR^{52}$, $-NR^{52}R^{53}$, $-SR_{52}$, $-NR^{52}S(O)_2R^{53}$, $-S(O)_2NR^{52}R^{53}$, $-S(O)NR^{52}R^{53}$, $-S(O)R^{52}$, $-S(O)_2R^{52}$, $-C(O)NR^{52}R^{53}$, $-OC(O)NR^{52}R^{53}$, $-NR^{52}C(O)R^{53}$, $-CH_2C(O)NR^{52}R^{53}$, $-OCH_2C(O)NR^{52}R^{53}$, $-CH_2OR^{52}$, $-CH_2NR^{52}R^{53}$, $-OC(O)R^{52}$, $-C(O)R^{52}$ and $-C(O)OR^{52}$, wherein

R^{52} and R^{53} are as defined for embodiment 1.

Embodiment 159: A compound according to embodiment 157, wherein

25 R^{52} and R^{53} , when attached to the same nitrogen atom, together with the said nitrogen atom may form a 3 to 8 membered heterocyclic ring optionally containing one or two further heteroatoms selected from nitrogen, oxygen and sulfur, and optionally containing one or two double bonds.

Embodiment 160: A compound according to embodiment 157, wherein

30 R^{52} and R^{53} independently of each other are hydrogen, C_{1-6} -alkyl, aryl- C_{1-6} -alkylene-heteroaryl- C_{1-6} -alkylene- heteroaryl, or aryl.

Embodiment 161: A compound according to embodiment 160, wherein

R^{52} and R^{53} independently of each other are hydrogen, C_{1-6} -alkyl, aryl- C_{1-6} -alkylene-or aryl.

35 Embodiment 162: A compound according to embodiment 161, wherein

R^{52} and R^{53} independently of each other are hydrogen or C_{1-6} -alkyl.

Embodiment 163: A compound according to embodiment 162, wherein

R^{52} is hydrogen.

Embodiment 164: A compound according to embodiment 162 or embodiment 163, wherein

5 R^{53} is hydrogen.

Embodiment 165: A compound according to embodiment 157, wherein

R^1 is hydrogen.

Embodiment 166: A compound according to any of embodiments 1 to 165, wherein

10 G^2 is heteroaryl, fused heterocyclylheteroaryl, or fused cycloalkylheteroaryl, optionally substituted with one or more substituents R^{43} , R^{44} , and R^{45} , wherein said heteroaryl group possesses a nitrogen atom adjacent to the atom joining G^2 with $-N(R^1)-$, and wherein

R^{43} , R^{44} , and R^{45} are as defined for embodiment 1.

Embodiment 167: A compound according to embodiment 166, wherein

15 G^2 is heteroaryl optionally substituted with one or more substituents R^{43} , R^{44} , and R^{45} , wherein said heteroaryl group possesses a nitrogen atom adjacent to the atom joining G^2 with $-N(R^1)-$, and wherein

R^{43} , R^{44} , and R^{45} are as defined for embodiment 1.

Embodiment 168: A compound according to embodiment 167, wherein

20 G^2 is furanyl, thienyl, thiophenyl, pyrrolyl, imidazolyl, pyrazolyl, triazolyl, tetrazolyl, thiazolyl, oxazolyl, isoxazolyl, oxadiazolyl, thiadiazolyl, isothiazolyl, pyridinyl, pyridazinyl, pyrazinyl, pyrimidinyl, quinoliny, isoquinoliny, benzofuranyl, benzothiophenyl, indolyl, or indazolyl, optionally substituted with one or more substituents R^{43} , R^{44} , and R^{45} , wherein

25 R^{43} , R^{44} , and R^{45} are as defined for embodiment 1.

Embodiment 169: A compound according to any of embodiments 1 to 168, wherein

30 R^{43} , R^{44} , and R^{45} independently of each other are selected from halogen, $-CN$, $-NO_2$, C_{1-6} -alkyl, $-CHF_2$, $-CF_3$, $-OCF_3$, $-OCHF_2$, $-OCH_2CF_3$, $-OCF_2CHF_2$, $-S(O)_2CF_3$, $-SCF_3$, $-OR^{54}$, $-NR^{54}R^{55}$, $-SR^{54}$, $-NR^{54}S(O)_2R^{55}$, $-S(O)_2NR^{54}R^{55}$, $-S(O)NR^{54}R^{55}$, $-S(O)R^{54}$, $-S(O)_2R^{54}$, $-C(O)NR^{54}R^{55}$, $-OC(O)NR^{54}R^{55}$, $-NR^{54}C(O)R^{55}$, $-CH_2C(O)NR^{54}R^{55}$, $-CH_2C(O)OR^{54}$, $-OCH_2C(O)NR^{54}R^{55}$, $-CH_2OR^{54}$, $-CH_2NR^{54}R^{55}$, $-OC(O)R^{54}$, $-C(O)R^{54}$ and $-C(O)OR^{54}$, wherein

R^{54} and R^{55} are as defined for embodiment 1.

Embodiment 170: A compound according to embodiment 169, wherein

R⁵⁴ and R⁵⁵, when attached to the same nitrogen atom, together with the said nitrogen atom may form a 3 to 8 membered heterocyclic ring optionally containing one or two further heteroatoms selected from nitrogen, oxygen and sulfur, and optionally containing one or two double bonds, and optionally substituted with one or more C₁₋₆-alkyl groups.

Embodiment 171: A compound according to embodiment 170, wherein

R⁵⁴ and R⁵⁵, when attached to the same nitrogen atom, together with the said nitrogen atom may form a 3 to 8 membered heterocyclic ring optionally containing one or two further heteroatoms selected from nitrogen, oxygen and sulfur, and optionally containing one or two double bonds.

Embodiment 172: A compound according to embodiment 169, wherein

R⁵⁴ and R⁵⁵ independently of each other are hydrogen, C₁₋₆-alkyl, aryl-C₁₋₆-alkylene- or aryl.

Embodiment 173: A compound according to embodiment 172, wherein

R⁵⁴ and R⁵⁵ independently of each other are hydrogen, C₁₋₆-alkyl, C₃₋₁₀-aryl-C₁₋₆-alkylene- or C₃₋₁₀-aryl.

Embodiment 174: A compound according to embodiment 173, wherein

R⁵⁴ and R⁵⁵ independently of each other are hydrogen or C₁₋₆-alkyl.

Embodiment 175: A compound according to embodiment 174, wherein

R⁵⁴ is hydrogen.

Embodiment 176: A compound according to embodiment 174 or embodiment 175, wherein

R⁵⁵ is hydrogen.

Embodiment 177: A compound according to embodiment 169, wherein

R⁵⁴ and R⁵⁵ independently of each other are hydrogen or -(CHR⁶³)_u-(CHR⁶⁴)_v-Z, wherein

u is 1 or 2;

v is 0, 1 or 2;

R⁶³ and R⁶⁴ independently of each other are hydrogen, C₁₋₆-alkyl, C₁₋₆-alkyl-C₃₋₁₀-arylene- C₃₋₁₀-aryl, hydroxy, hydroxy-C₁₋₆-alkyl, amino, or amino-C₁₋₆-alkyl;

Z is hydrogen, -C-O-R⁶⁵, -C(O)O-R⁶⁵, -CONR⁶⁵R⁶⁶, C₁₋₆-alkylamino, or di(C₁₋₆-alkyl)-amino, wherein

R⁶⁵ and R⁶⁶ independently of each other is hydrogen or C₁₋₆-alkyl;

or

Z is a five or six membered ring wherein at least one ring atom is nitrogen and the remaining ring atoms are either carbon or oxygen.

Embodiment 178: A compound according to embodiment 177, wherein

u is 1; and

5

v is 0, or 1.

Embodiment 179: A compound according to embodiment 177 or embodiment 178, wherein

R⁶³ and R⁶⁴ independently of each other are hydrogen, C₁₋₆-alkyl, hydroxy, hydroxy-C₁₋₆-alkyl, amino, or amino-C₁₋₆-alkyl.

Embodiment 180: A compound according to any of embodiments 177 to 179, wherein

10

Z is -C-O-R⁶⁵, or -C(O)O-R⁶⁵, wherein

R⁶⁵ and R⁶⁶ independently of each other is hydrogen or C₁₋₆-alkyl.

Embodiment 181: A compound according to any of embodiments 177 to 179, wherein

Z is a five or six membered ring where at least one ring atom is nitrogen and the remaining ring atoms are either carbon or oxygen.

15

Embodiment 182: A compound according to embodiment 181, wherein

Z is a five or six membered ring wherein at least one ring atom is nitrogen, one ring atom is oxygen and the remaining ring atoms are carbon.

Embodiment 183: A compound according to embodiment 181, wherein

20

Z is a five or six membered ring wherein at least one ring atom is nitrogen, and the remaining ring atoms are carbon.

Embodiment 184: A compound according to any of embodiments 181 to 183, wherein

Z is a five or six membered ring wherein one ring atom is nitrogen.

Embodiment 185: A compound according to any of embodiments 181 to 184, wherein

25

a nitrogen atom is the point of attachment of the Z group to the -(CHR⁶³)_u-(CHR⁶⁴)_v-group.

Embodiment 186: A compound according to any of embodiments 177 to 185, wherein

R⁵⁵ is hydrogen.

Embodiment 187: A compound according to any of embodiments 166 to 169, wherein

G² is substituted with a substituent R⁴³, wherein

30

R⁴³ is halogen, -CN, -NO₂, C₁₋₆-alkyl, -CHF₂, -CF₃, -OCF₃, -OCHF₂, -OCH₂CF₃, -OCF₂CHF₂, -S(O)₂CF₃, -SCF₃, -OR⁵⁴, -NR⁵⁴R⁵⁵, -SR⁵⁴, -NR⁵⁴S(O)₂R⁵⁵, -S(O)₂NR⁵⁴R⁵⁵, -S(O)NR⁵⁴R⁵⁵, -S(O)R⁵⁴, -S(O)₂R⁵⁴, -C(O)NR⁵⁴R⁵⁵, -OC(O)NR⁵⁴R⁵⁵, -NR⁵⁴C(O)R⁵⁵, -CH₂C(O)NR⁵⁴R⁵⁵, -CH₂C(O)OR⁵⁴, -OCH₂C(O)NR⁵⁴R⁵⁵, -CH₂OR⁵⁴, -CH₂NR⁵⁴R⁵⁵, -OC(O)R⁵⁴, -C(O)R⁵⁴ or -C(O)OR⁵⁴, wherein

35

R^{54} and R^{55} are as defined for embodiment 1.

Embodiment 188: A compound according to embodiment 187, wherein

5 R^{54} and R^{55} , when attached to the same nitrogen atom, together with the said nitrogen atom may form a 3 to 8 membered heterocyclic ring optionally containing one or two further heteroatoms selected from nitrogen, oxygen and sulfur, and optionally containing one or two double bonds, and optionally substituted with one or more C_{1-6} -alkyl groups.

Embodiment 189: A compound according to embodiment 188, wherein

10 R^{54} and R^{55} , when attached to the same nitrogen atom, together with the said nitrogen atom may form a 3 to 8 membered heterocyclic ring optionally containing one or two further heteroatoms selected from nitrogen, oxygen and sulfur, and optionally containing one or two double bonds.

Embodiment 190: A compound according to embodiment 187, wherein

15 R^{54} and R^{55} independently of each other are hydrogen, C_{1-6} -alkyl, aryl- C_{1-6} -alkylene- or aryl.

Embodiment 191: A compound according to embodiment 190, wherein

R^{54} and R^{55} independently of each other are hydrogen, C_{1-6} -alkyl, C_{3-10} -aryl- C_{1-6} -alkylene- or C_{3-10} -aryl.

Embodiment 192: A compound according to embodiment 191, wherein

20 R^{54} and R^{55} independently of each other are hydrogen or C_{1-6} -alkyl.

Embodiment 193: A compound according to embodiment 192, wherein

R^{54} is hydrogen.

Embodiment 194: A compound according to embodiment 192 or embodiment 193, wherein

R^{55} is hydrogen.

25 Embodiment 195: A compound according to embodiment 187, wherein

R^{54} and R^{55} independently of each other are hydrogen or $-(CHR^{63})_u-(CHR^{64})_v-Z$, wherein

u is 1 or 2;

v is 0, 1 or 2;

30 R^{63} and R^{64} independently of each other are hydrogen, C_{1-6} -alkyl, C_{1-6} -alkyl- C_{3-10} -arylene- C_{3-10} -aryl, hydroxy, hydroxy- C_{1-6} -alkyl, amino, or amino- C_{1-6} -alkyl;

Z is hydrogen, $-C-O-R^{65}$, $-C(O)O-R^{65}$, $-CONR^{65}R^{66}$, C_{1-6} -alkylamino, or di(C_{1-6} -alkyl)-amino, wherein

35 R^{65} and R^{66} independently of each other is hydrogen or C_{1-6} -alkyl;

or

Z is a five or six membered ring wherein at least one ring atom is nitrogen and the remaining ring atoms are either carbon or oxygen.

Embodiment 196: A compound according to embodiment 195, wherein

5 u is 1; and

v is 0, or 1.

Embodiment 197: A compound according to embodiment 195 or embodiment 196, wherein

R⁶³ and R⁶⁴ independently of each other are hydrogen, C₁₋₆-alkyl, hydroxy, hydroxy-C₁₋₆-alkyl, amino, or amino-C₁₋₆-alkyl.

10 Embodiment 198: A compound according to any of embodiments 195 to 197, wherein

Z is -C-O-R⁶⁵, or -C(O)O-R⁶⁵, wherein

R⁶⁵ and R⁶⁶ independently of each other is hydrogen or C₁₋₆-alkyl.

Embodiment 199: A compound according to any of embodiments 195 to 197, wherein

15 Z is a five or six membered ring where at least one ring atom is nitrogen and the remaining ring atoms are either carbon or oxygen.

Embodiment 200: A compound according to embodiment 199, wherein

Z is a five or six membered ring wherein at least one ring atom is nitrogen, one ring atom is oxygen and the remaining ring atoms are carbon.

Embodiment 201: A compound according to embodiment 199, wherein

20 Z is a five or six membered ring wherein at least one ring atom is nitrogen, and the remaining ring atoms are carbon.

Embodiment 202: A compound according to any of embodiments 199 to 201, wherein

Z is a five or six membered ring wherein one ring atom is nitrogen.

Embodiment 203: A compound according to any of embodiments 199 to 202, wherein

25 a nitrogen atom is the point of attachment of the Z group to the -(CHR⁶³)_u-(CHR⁶⁴)_v-group.

Embodiment 204: A compound according to any of embodiments 195 to 203, wherein

R⁵⁵ is hydrogen.

Embodiment 205: A compound according to embodiment 187, wherein

30 R⁴³ is -CH₂C(O)OR⁵⁴, wherein

R⁵⁴ is as defined for embodiment 1.

Embodiment 206: A compound according to embodiment 205, wherein

R⁵⁴ is hydrogen, C₁₋₆-alkyl, aryl-C₁₋₆-alkylene- or aryl.

Embodiment 207: A compound according to embodiment 206, wherein

35 R⁵⁴ is hydrogen, C₁₋₆-alkyl, C₃₋₁₀-aryl-C₁₋₆-alkylene- or C₃₋₁₀-aryl.

Embodiment 208: A compound according to embodiment 207, wherein


R^{54} is hydrogen or C_{1-6} -alkyl.

Embodiment 209: A compound according to embodiment 208, wherein

R^{54} is hydrogen.

- 5 Embodiment 210: A compound according to any of embodiments 187 to 209, wherein R^{43} is attached to the atom adjacent to the nitrogen atom adjacent to to the atom joining G^2 with $-N(R^1)-$.


Embodiment 211: A compound according to any of embodiments 166 to 210, wherein G^2 is



R^{43} , R^{44} , and R^{45} independently of each other are hydrogen or as defined for embodiment 1.

Embodiment 212: A compound according to embodiment 211, wherein

G^2 is



R^{43} , R^{44} , and R^{45} independently of each other are hydrogen or as defined for embodiment 1.

Embodiment 213: A compound according to embodiment 211 or embodiment 212, wherein

- 20 R^{43} , R^{44} , and R^{45} independently of each other are selected from hydrogen, halogen, -CN, -NO₂, C_{1-6} -alkyl, -CHF₂, -CF₃, -OCF₃, -OCHF₂, -OCH₂CF₃, -OCF₂CHF₂, -S(O)₂CF₃, -SCF₃, -OR⁵⁴, -NR⁵⁴R⁵⁵, -SR⁵⁴, -NR⁵⁴S(O)₂R⁵⁵, -S(O)₂NR⁵⁴R⁵⁵, -S(O)NR⁵⁴R⁵⁵, -S(O)R⁵⁴, -S(O)₂R⁵⁴, -C(O)NR⁵⁴R⁵⁵, -OC(O)NR⁵⁴R⁵⁵, -NR⁵⁴C(O)R⁵⁵, -CH₂C(O)NR⁵⁴R⁵⁵, -CH₂C(O)OR⁵⁴, -OCH₂C(O)NR⁵⁴R⁵⁵, -CH₂OR⁵⁴, -CH₂NR⁵⁴R⁵⁵, -OC(O)R⁵⁴, -C(O)R⁵⁴ and -C(O)OR⁵⁴, wherein

- 25 R^{54} and R^{55} are as defined for embodiment 1.

Embodiment 214: A compound according to embodiment 213, wherein

R^{43} is halogen, -CN, -NO₂, C_{1-6} -alkyl, -CHF₂, -CF₃, -OCF₃, -OCHF₂, -OCH₂CF₃, -OCF₂CHF₂, -S(O)₂CF₃, -SCF₃, -OR⁵⁴, -NR⁵⁴R⁵⁵, -SR⁵⁴, -NR⁵⁴S(O)₂R⁵⁵,

-S(O)₂NR⁵⁴R⁵⁵, -S(O)NR⁵⁴R⁵⁵, -S(O)R⁵⁴, -S(O)₂R⁵⁴, -C(O)NR⁵⁴R⁵⁵, -OC(O)NR⁵⁴R⁵⁵,
 -NR⁵⁴C(O)R⁵⁵, -CH₂C(O)NR⁵⁴R⁵⁵, -CH₂C(O)OR⁵⁴, -OCH₂C(O)NR⁵⁴R⁵⁵, -CH₂OR⁵⁴,
 -CH₂NR⁵⁴R⁵⁵, -OC(O)R⁵⁴, -C(O)R⁵⁴ or -C(O)OR⁵⁴, wherein

R⁵⁴ and R⁵⁵ are as defined for embodiment 1.

5 Embodiment 215: A compound according to embodiment 213 or 214, wherein

R⁴³ is -C(O)OR⁵⁴, -CH₂C(O)OR⁵⁴, -C(O)NR⁵⁴R⁵⁵, or -CH₂C(O)NR⁵⁴R⁵⁵, wherein

R⁵⁴ and R⁵⁵ are as defined for embodiment 1.

Embodiment 216: A compound according to embodiment 214 or 215, wherein

R⁴⁴ is alkyl or hydrogen.

10 Embodiment 217: A compound according to embodiment 216, wherein R⁴⁴ is

C₁₋₆-alkyl or hydrogen.

Embodiment 218: A compound according to embodiment 217, wherein

R⁴⁴ is hydrogen.

Embodiment 219: A compound according to embodiment 213, wherein

15 R⁴⁴ is halogen, -CN, -NO₂, C₁₋₆-alkyl, -CHF₂, -CF₃, -OCF₃, -OCHF₂, -OCH₂CF₃,
 -OCF₂CHF₂, -S(O)₂CF₃, -SCF₃, -OR⁵⁴, -NR⁵⁴R⁵⁵, -SR⁵⁴, -NR⁵⁴S(O)₂R⁵⁵,
 -S(O)₂NR⁵⁴R⁵⁵, -S(O)NR⁵⁴R⁵⁵, -S(O)R⁵⁴, -S(O)₂R⁵⁴, -C(O)NR⁵⁴R⁵⁵, -OC(O)NR⁵⁴R⁵⁵,
 -NR⁵⁴C(O)R⁵⁵, -CH₂C(O)NR⁵⁴R⁵⁵, -CH₂C(O)OR⁵⁴, -OCH₂C(O)NR⁵⁴R⁵⁵, -CH₂OR⁵⁴,
 -CH₂NR⁵⁴R⁵⁵, -OC(O)R⁵⁴, -C(O)R⁵⁴ or -C(O)OR⁵⁴, wherein

20 R⁵⁴ and R⁵⁵ are as defined for embodiment 1.

Embodiment 220: A compound according to embodiment 213 or embodiment 219, wherein

R⁴⁴ is -C(O)OR⁵⁴, -CH₂C(O)OR⁵⁴, -C(O)NR⁵⁴R⁵⁵, or -CH₂C(O)NR⁵⁴R⁵⁵, wherein

R⁵⁴ and R⁵⁵ are as defined for embodiment 1.

Embodiment 221: A compound according to embodiment 219 or 220, wherein

25 R⁴³ is alkyl or hydrogen.

Embodiment 222: A compound according to embodiment 221, wherein

R⁴³ is C₁₋₆-alkyl or hydrogen.

Embodiment 223: A compound according to embodiment 222, wherein

R⁴³ is hydrogen.

30 Embodiment 224: A compound according to any of embodiments 211 to 223, wherein

R⁴⁵ is hydrogen.

Embodiment 225: A compound according to any of embodiments 211 to 224, wherein

R⁵⁴ and R⁵⁵, when attached to the same nitrogen atom, together with the said
 nitrogen atom may form a 3 to 8 membered heterocyclic ring optionally containing
 35 one or two further heteroatoms selected from nitrogen, oxygen and sulfur, and

optionally containing one or two double bonds, and optionally substituted with one or more C₁₋₆-alkyl groups.

Embodiment 226: A compound according to embodiment 225, wherein

5 R⁵⁴ and R⁵⁵, when attached to the same nitrogen atom, together with the said nitrogen atom may form a 3 to 8 membered heterocyclic ring optionally containing one or two further heteroatoms selected from nitrogen, oxygen and sulfur, and optionally containing one or two double bonds.

Embodiment 227: A compound according to any of embodiments 211 to 224, wherein

10 R⁵⁴ and R⁵⁵ independently of each other are hydrogen, C₁₋₆-alkyl, aryl-C₁₋₆-alkylene- or aryl.

Embodiment 228: A compound according to any of embodiments 211 to 224, wherein

R⁵⁴ and R⁵⁵ independently of each other are hydrogen, C₁₋₆-alkyl, aryl-C₁₋₆-alkylene- or aryl.

Embodiment 229: A compound according to embodiment 228, wherein

15 R⁵⁴ and R⁵⁵ independently of each other are hydrogen or C₁₋₆-alkyl.

Embodiment 230: A compound according to embodiment 229, wherein

R⁵⁴ is hydrogen.

Embodiment 231: A compound according to any of embodiments 211 to 230, wherein

R⁵⁵ is hydrogen.

20 Embodiment 232: A compound according to any of embodiments 211 to 224, wherein

R⁵⁴ and R⁵⁵ independently of each other are hydrogen or -(CHR⁶³)_u-(CHR⁶⁴)_v-Z, wherein

u is 1 or 2;

v is 0, 1 or 2;

25 R⁶³ and R⁶⁴ independently of each other are hydrogen, C₁₋₆-alkyl, C₁₋₆-alkyl-C₃₋₁₀-arylene- C₃₋₁₀-aryl, hydroxy, hydroxy-C₁₋₆-alkyl, amino, or amino-C₁₋₆-alkyl;

Z is hydrogen, -C-O-R⁶⁵, -C(O)O-R⁶⁵, -CONR⁶⁵R⁶⁶, C₁₋₆-alkylamino, or di(C₁₋₆-alkyl)-amino, wherein

30 R⁶⁵ and R⁶⁶ independently of each other is hydrogen or C₁₋₆-alkyl; or

Z is a five or six membered ring wherein at least one ring atom is nitrogen and the remaining ring atoms are either carbon or oxygen.

Embodiment 233: A compound according to embodiment 232, wherein

35 u is 1; and

v is 0, or 1.

Embodiment 234: A compound according to embodiment 232 or embodiment 233, wherein R^{63} and R^{64} independently of each other are hydrogen, C_{1-6} -alkyl, hydroxy, hydroxy- C_{1-6} -alkyl, amino, or amino- C_{1-6} -alkyl.

- 5 Embodiment 235: A compound according to any of embodiments 232 to 234, wherein Z is $-C(O)R^{65}$, or $-C(O)O-R^{65}$, wherein R^{65} and R^{66} independently of each other is hydrogen or C_{1-6} -alkyl.

- Embodiment 236: A compound according to any of embodiments 232 to 234, wherein Z is a five or six membered ring where at least one ring atom is nitrogen and the
10 remaining ring atoms are either carbon or oxygen.

Embodiment 237: A compound according to embodiment 236, wherein Z is a five or six membered ring wherein at least one ring atom is nitrogen, one ring atom is oxygen and the remaining ring atoms are carbon.

- Embodiment 238: A compound according to embodiment 236, wherein
15 Z is a five or six membered ring wherein at least one ring atom is nitrogen, and the remaining ring atoms are carbon.

Embodiment 239: A compound according to any of embodiments 236 to 238, wherein Z is a five or six membered ring wherein one ring atom is nitrogen.

- Embodiment 240: A compound according to any of embodiments 236 to 239, wherein
20 a nitrogen atom is the point of attachment of the Z group to the $-(CHR^{63})_v-(CHR^{64})_v$ -group.

Embodiment 241: A compound according to any of embodiments 195 to 240, wherein R^{65} is hydrogen.

- Embodiment 242: A compound according to embodiment 213, wherein
25 R^{64} is hydrogen.

Embodiment 243: A compound according to embodiment 213 or embodiment 242, wherein R^{63} is hydrogen.

Embodiment 244: A compound according to embodiment 242 or embodiment 243, wherein R^{65} is hydrogen.

- 30 Embodiment 245: A compound according to embodiment 1, which is,
N-(2-phenoxyphenyl)-N'-(thiazol-2-yl)urea,
N-[2-(2,3-dimethoxyphenoxy)-5-fluorophenyl]-N'-(thiazol-2-yl)sulfamide,
1-(2-phenoxyphenyl)-5-(thiazol-2-yl)biuret,
2-[[[(2-phenoxyanilino)sulfonyl]amino]carbonyl]amino]thiazole,
35 N-(2-phenylsulfanylphenyl)-N'-(thiazol-2-yl)urea,

- N-(2-phenylsulfonylphenyl)-N'-(thiazol-2-yl)urea,
N-(2-benzylphenyl)-N'-(thiazol-2-yl)urea,
N-(2-benzoylphenyl)-N'-(thiazol-2-yl)urea,
N-[2-(phenylamino)phenyl]-N'-(thiazol-2-yl)urea,
5 N-[2-fluoro-6-(4-methoxyphenoxy)benzyl]-N'-(thiazol-2-yl)urea,
N-(2-benzyloxyphenyl)-N'-(thiazol-2-yl)urea,
N-[2-(2,3,4-trimethoxybenzyloxy)phenyl]-N'-(thiazol-2-yl)urea,
N-(2-ethoxyphenyl)-N'-(thiazol-2-yl)urea,
N-(2-phenoxyphenyl)-N'-(pyridin-2-yl)urea,
10 N-(2-phenoxyphenyl)-N'-[(4-methoxycarbonylmethyl)thiazol-2-yl]urea,
N-methyl-N-(2-phenoxyphenyl)-N'-(thiazol-2-yl)urea,
N-isopropyl-N-(2-phenoxyphenyl)-N'-(thiazol-2-yl)urea,
N-[2-(4-methoxyphenoxy)phenyl]-N'-(thiazol-2-yl)urea,
N-[2-(4-fluorophenoxy)phenyl]-N'-(thiazol-2-yl)urea,
15 N-[2-(4-chlorophenoxy)phenyl]-N'-(thiazol-2-yl)urea,
N-[2-(4-cyanophenoxy)phenyl]-N'-thiazolylurea,
N-[2-(4-methoxycarbonylphenoxy)phenyl]-N'-(thiazol-2-yl)urea,
N-[2-(4-isopropylphenoxy)phenyl]-N'-(thiazol-2-yl)urea,
N-[2-(3,4-difluorophenoxy)phenyl]-N'-(thiazol-2-yl)urea,
20 N-[2-(3,4-dichlorophenoxy)phenyl]-N'-(thiazol-2-yl)urea,
N-[2-(4-chloro-3-methylphenoxy)phenyl]-N'-(thiazol-2-yl)urea,
N-[2-(3,4-dimethoxyphenoxy)phenyl]-N'-(thiazol-2-yl)urea,
N-[2-(3,4-methylenedioxyphenoxy)phenyl]-N'-(thiazol-2-yl)urea,
N-[2-(2,4-dichlorophenoxy)phenyl]-N'-(thiazol-2-yl)urea,
25 N-[2-(2,4-difluorophenoxy)phenyl]-N'-(thiazol-2-yl)urea,
N-[2-(4-fluoro-2-methoxyphenoxy)phenyl]-N'-(thiazol-2-yl)urea,
N-[2-(4-methoxy-2-methoxycarbonylphenoxy)phenyl]-N'-(thiazol-2-yl)urea,
N-[2-(3-methoxyphenoxy)phenyl]-N'-(thiazol-2-yl)urea,
N-[2-(3-fluorophenoxy)phenyl]-N'-(thiazol-2-yl)urea,
30 N-[2-(3-trifluoromethylphenoxy)phenyl]-N'-(thiazol-2-yl)urea,
N-[2-(2-methylphenoxy)phenyl]-N'-(thiazol-2-yl)urea,
N-[2-(2-methoxyphenoxy)phenyl]-N'-(thiazol-2-yl)urea,
N-[2-(2-isopropoxyphenoxy)phenyl]-N'-(thiazol-2-yl)urea,
N-[2-(2-fluorophenoxy)phenyl]-N'-(thiazol-2-yl)urea,
35 N-[2-(2-methylsulfanylphenoxy)phenyl]-N'-(thiazol-2-yl)urea,

- N-[2-(2-methylsulfonylphenoxy)phenyl]-N'-thiazolylurea,
N-[2-(2-trifluoromethylphenoxy)phenyl]-N'-(thiazol-2-yl)urea,
N-[2-(2,6-dimethoxyphenoxy)phenyl]-N'-(thiazol-2-yl)urea,
N-[2-(2,6-difluorophenoxy)phenyl]-N'-(thiazol-2-yl)urea,
5 N-[2-(2-fluoro-6-methoxyphenoxy)phenyl]-N'-(thiazol-2-yl)urea,
N-[2-(2-methoxy-6-methoxycarbonylphenoxy)phenyl]-N'-(thiazol-2-yl)urea,
N-[(3-methoxy-2-methoxycarbonylphenoxy)phenyl]-N'-(thiazol-2-yl)urea,
N-[2-(2,3-dimethoxyphenoxy)phenyl]-N'-(thiazol-2-yl)urea,
N-[2-(2,3,4-trichlorophenoxy)phenyl]-N'-(thiazol-2-yl)urea,
10 N-[2-(2,4,6-trifluorophenoxy)phenyl]-N'-(thiazol-2-yl)urea,
N-[2-(2,4-dichloronaphth-1-oxy)phenyl]-N'-(thiazol-2-yl)urea,
N-[2-(2-methoxyphenoxy)-5-(methylsulfonyl)phenyl]-N'-(thiazol-2-yl)urea,
N-[5-cyano-2-(2-methoxyphenoxy)phenyl]-N'-(thiazol-2-yl)urea,
N-[5-fluoro-2-(2-methylsulfonylphenoxy)phenyl]-N'-thiazolylurea,
15 N-[5-fluoro-2-(2-fluoro-6-methoxyphenoxy)phenyl]-N'-(thiazol-2-yl)urea,
N-[2-(2,3-dimethoxyphenoxy)-5-fluorophenyl]-N'-(thiazol-2-yl)urea,
N-[2-(3,4-difluorophenoxy)-5-fluorophenyl]-N'-(thiazol-2-yl)urea,
N-[5-fluoro-2-(4-fluorophenoxy)phenyl]-N'-(thiazol-2-yl)urea,
N-[2-(2,4-dichlorophenoxy)-5-fluorophenyl]-N'-(thiazol-2-yl)urea,
20 N-[5-fluoro-2-(4-methoxyphenoxy)phenyl]-N'-(thiazol-2-yl)urea,
N-[5-fluoro-2-(2-methoxyphenoxy)phenyl]-N'-(thiazol-2-yl)urea,
N-[5-fluoro-2-(2-trifluoromethylphenoxy)phenyl]-N'-(thiazol-2-yl)urea,
N-[5-fluoro-2-(naphth-2-oxy)phenyl]-N'-(thiazol-2-yl)urea,
N-[2-(2,3-dimethoxyphenoxy)-6-fluorophenyl]-N'-(thiazol-2-yl)urea,
25 N-[2-(4-fluoro-2-methoxyphenoxy)phenyl]-N'-(thiazol-2-yl)urea,
N-[2-(2,3-dimethoxyphenoxy)-4-fluorophenyl]-N'-(thiazol-2-yl)urea,
N-[4-fluoro-2-(2-fluoro-6-methoxyphenoxy)phenyl]-N'-(thiazol-2-yl)urea,
N-[2-(2-fluoro-6-methoxyphenoxy)-4-methoxyphenyl]-N'-(thiazol-2-yl)urea,
N-[3-fluoro-2-(2-fluoro-6-methoxyphenoxy)phenyl]-N'-(thiazol-2-yl)urea,
30 N-[2-(2,3-dimethoxyphenoxy)-5-methoxyphenyl]-N'-thiazol-2-ylurea,
N-[2-(2,3-dimethoxyphenoxy)-4-methylphenyl]-N'-(thiazol-2-yl)urea,
N-[2-(2,3-dimethoxyphenoxy)-3-methylphenyl]-N'-(thiazol-2-yl)urea,
N-[2-(2-chlorophenoxy)-5-chlorophenyl]-N'-(thiazol-2-yl)urea,
N-[5-chloro-2-(4-chloro-3-methylphenoxy)phenyl]-N'-(thiazol-2-yl)urea,
35 N-[2-(4-chlorophenoxy)-5-(trifluoromethyl)phenyl]-N'-(thiazol-2-yl)urea,

- N-[4,5-difluoro-2-(2,3-dimethoxyphenoxy)phenyl]-N'-(thiazol-2-yl)urea,
N-[4,5-dichloro-2-(2,3-dimethoxyphenoxy)phenyl]-N'-(thiazol-2-yl)urea,
N-[5-chloro-2-(2,3-dimethoxyphenoxy)-4-dimethylaminophenyl]-N'-(thiazol-2-yl)urea,
N-[5-chloro-2-(2,3-dimethoxyphenoxy)-4-(4-morpholino)phenyl]-N'-(thiazol-2-yl)urea,
5 N-[2-(2,4-difluorophenoxy)pyridin-3-yl]-N'-(thiazol-2-yl)urea,
N-[2-(2-fluorophenoxy)pyridin-3-yl]-N'-(thiazol-2-yl)urea,
N-[2-(2-methoxyphenoxy)pyridin-3-yl]-N'-(thiazol-2-yl)urea,
N-[2-(2,3-dimethoxyphenoxy)pyridin-3-yl]-N'-(thiazol-2-yl)urea,
N-[4-chloro-2-(2-chlorobenzoyl)phenyl]-N'-(thiazol-2-yl)urea,
10 N-[4-chloro-2-(2-fluorobenzoyl)phenyl]-N'-(thiazol-2-yl)urea,
N-[2-(2,4-difluorophenylsulfanyl)phenyl]-N'-(thiazol-2-yl)urea,
1-[2-(2-fluorophenylsulfanyl)phenyl]-3-(thiazol-2-yl)urea,
N-[2-(2-chloro-4-fluorophenylsulfanyl)phenyl]-N'-(thiazol-2-yl)urea,
N-[2-(2,3-dichlorophenylsulfanyl)phenyl]-N'-(thiazol-2-yl)urea,
15 N-[2-(3,5-dimethylphenylsulfanyl)phenyl]-N'-(thiazol-2-yl)urea,
N-[2-(2-methoxycarbonylphenylsulfanyl)phenyl]-N'-(thiazol-2-yl)urea,
N-[2-(2-methoxyphenylsulfanyl)phenyl]-N'-(thiazol-2-yl)urea,
N-[2-(2-pyridinylsulfanyl)phenyl]-N'-(thiazol-2-yl)urea,
N-(2-propyloxyphenyl)-N'-(thiazol-2-yl)urea,
20 N-(2-butyloxyphenyl)-N'-(thiazol-2-yl)urea,
N-(2-(cyclopentyloxy)phenyl)-N'-(thiazol-2-yl)urea,
N-(2-isopropoxyphenyl)-N'-(thiazol-2-yl)urea,
N-[2-(2-methylpropoxy)phenyl]-N'-(thiazol-2-yl)urea,
N-[2-(cyclopentylmethoxy)phenyl]-N'-(thiazol-2-yl)urea,
25 N-[2-(3-pentoxy)phenyl]-N'-(thiazol-2-yl)urea,
N-[2-(2-pentoxy)phenyl]-N'-(thiazol-2-yl)urea,
N-[2-(2-methoxyethoxy)phenyl]-N'-(thiazol-2-yl)urea,
(2-[3-(2-benzylphenyl)ureido]thiazol-4-yl)acetic acid,
(2-[3-(2-benzoyl-4-chlorophenyl)ureido]thiazol-4-yl)acetic acid,
30 (2-[3-(2-(2-methylphenoxy)phenyl)ureido]thiazol-4-yl)acetic acid,
(2-[3-(2-(4-methoxyphenoxy)-5-(trifluoromethyl)phenyl)ureido]thiazol-4-yl)acetic acid,
{2-[3-(2-phenoxyphenyl)ureido]thiazol-4-yl}acetic acid,
2-{3-[5-fluoro-2-(2-fluoro-6-methoxyphenoxy)phenyl]ureido}thiazole-4-carboxylic acid,
2-{3-[2-(2,3-dimethoxyphenoxy)-5-fluorophenyl]ureido}thiazole-4-carboxylic acid ethyl ester,
35 (2-{3-[2-(2,3-dimethoxyphenoxy)-5-fluorophenyl]ureido}thiazol-4-yl)acetic acid ethyl ester,

- (2-{3-[5-fluoro-2-(2-fluoro-6-methoxyphenoxy)phenyl]ureido}thiazol-4-yl)acetic acid,
2-{3-[2-(2,3-dimethoxyphenoxy)-5-fluorophenyl]ureido}thiazole-4-carboxylic acid,
(2-{3-[2-(2,3-dimethoxyphenoxy)-5-fluorophenyl]ureido}thiazol-4-yl)acetic acid,
2-{3-[5-fluoro-2-(2-fluoro-6-methoxyphenoxy)phenyl]ureido}-4-methylthiazole-5-carboxylic
5 acid ethyl ester,
2-{3-[5-fluoro-2-(2-fluoro-6-methoxyphenoxy)phenyl]ureido}-4-methylthiazole-5-carboxylic
acid,
N-ethyl-2-(2-{3-[5-fluoro-2-(2-fluoro-6-methoxyphenoxy)phenyl]ureido}thiazol-4-yl)acetamide,
2-(2-{3-[5-fluoro-2-(2-fluoro-6-methoxyphenoxy)phenyl]ureido}thiazol-4-yl)-N-(2-methoxy-
10 ethyl)acetamide,
2-(2-{3-[5-fluoro-2-(2-fluoro-6-methoxyphenoxy)phenyl]ureido}thiazol-4-yl)-N-(2-morpholin-4-
ylethyl)acetamide,
[2-(2-{3-[5-fluoro-2-(2-fluoro-6-methoxyphenoxy)phenyl]ureido}thiazol-4-yl)acetylamino]acetic
acid methyl ester,
15 2-{3-[2-(2,3-dimethoxyphenoxy)-5-fluorophenyl]ureido}thiazole-4-carboxylic acid (2-
methoxyethyl)amide,
[2-(2-{3-[5-fluoro-2-(2-fluoro-6-methoxyphenoxy)phenyl]ureido}thiazol-4-yl)acetylamino]acetic
acid,
2-{3-[2-(2,3-dimethoxyphenoxy)-5-fluorophenyl]ureido}thiazole-4-carboxylic acid ethylamide,
20 [(2-{3-[2-(2,3-dimethoxyphenoxy)-5-fluorophenyl]ureido}thiazole-4-carbonyl)amino]acetic acid
methyl ester,
(5-{3-[5-fluoro-2-(2-fluoro-6-methoxyphenoxy)phenyl]ureido}-[1,3,4]thiadiazol-2-yl)acetic acid
ethyl ester,
[(2-{3-[2-(2,3-dimethoxyphenoxy)-5-fluorophenyl]ureido}thiazole-4-carbonyl)amino]acetic
25 acid,
(5-{3-[5-fluoro-2-(2-fluoro-6-methoxyphenoxy)phenyl]ureido}-[1,3,4]thiadiazol-2-yl)acetic acid,
5-{3-[2-(2,3-dimethoxyphenoxy)-5-fluorophenyl]ureido}-[1,3,4]thiadiazole-2-carboxylic acid
ethyl ester,
(5-{3-[2-(2,3-dimethoxyphenoxy)-5-fluorophenyl]ureido}-[1,3,4]thiadiazol-2-yl) acetic acid
30 ethyl ester,
3-[2-(2-{3-[2-(2,3-dimethoxyphenoxy)-5-fluorophenyl]ureido}thiazol-4-
yl)acetylamino]propionic acid methyl ester,
2-(2-{3-[2-(2,3-dimethoxyphenoxy)-5-fluorophenyl]ureido}thiazol-4-yl)-N-(2-morpholin-4-
ylethyl)acetamide,

[(2-{3-[5-fluoro-2-(2-fluoro-6-methoxyphenoxy)phenyl]ureido}thiazole-4-carbonyl)amino]acetic acid methyl ester,

3-[2-(2-{3-[2-(2,3-dimethoxyphenoxy)-5-fluorophenyl]ureido}thiazol-4-yl)acetilamino]propionic acid,

5 [(2-{3-[5-fluoro-2-(2-fluoro-6-methoxyphenoxy)phenyl]ureido}thiazole-4-carbonyl)amino]acetic acid,

(R) 3-[(2-{3-[5-fluoro-2-(2-fluoro-6-methoxyphenoxy)phenyl]ureido}thiazole-4-carbonyl)amino]-2-hydroxy-propionic acid,

2-[(2-{3-[5-fluoro-2-(2-fluoro-6-methoxyphenoxy)phenyl]ureido}thiazole-4-carbonyl)amino]-3-hydroxy-propionic acid,

1-(2-cyclopentanecarbonyl-4-methylphenyl)-3-thiazol-2-yl-urea,

1-(2-isobutyryl-4-methylphenyl)-3-thiazol-2-yl-urea,

1-[5-fluoro-2-(3-methylbutyryl)phenyl]-3-thiazol-2-yl-urea,

1-[5-methyl-2-(3-methylbutyryl)phenyl]-3-thiazol-2-yl-urea,

15 [3-cyclopentanecarbonyl-4-(3-thiazol-2-yl-ureido)phenyl]acetic acid ethyl ester,

[3-cyclopentanecarbonyl-4-(3-thiazol-2-yl-ureido)phenyl]acetic acid,

2-[3-cyclopentanecarbonyl-4-(3-thiazol-2-yl-ureido)phenyl]-N-methylacetamide,

{2-[3-(2-cyclopentanecarbonyl-4-methyl-phenyl)-ureido]-thiazol-4-yl}-acetic acid ethyl ester,

{2-[3-(2-cyclopentanecarbonyl-4-methyl-phenyl)-ureido]-thiazol-4-yl}acetic acid,

20 {3-cyclopentanecarbonyl-4-{3-(4-ethoxycarbonylmethylthiazol-2-yl)-ureido}phenyl}acetic acid ethyl ester,

1-(2-cyclopentanecarbonyl-4-methyl-phenyl)-3-(5-methanesulfonyl-thiazol-2-yl)-urea,

2-[3-(2-cyclopentanecarbonyl-4-methylphenyl)-ureido]-thiazole-4-carboxylic acid ethyl ester,

2-[3-(2-cyclopentanecarbonyl-4-methylphenyl)-ureido]-thiazole-4-carboxylic acid,

25 2-[3-(2-cyclopentanecarbonyl-4-methylphenyl)-ureido]-thiazole-4-carboxamide,

2-{2-[3-(2-cyclopentanecarbonyl-4-methylphenyl)-ureido]-thiazole-4-yl}-acetamide,

2-{2-[3-(2-cyclopentanecarbonyl-4-methylphenyl)-ureido]-thiazol-4-yl}-N-methyl-acetamide,

4-(2-{2-[3-(2-cyclopentanecarbonyl-4-methylphenyl)-ureido]-thiazol-4-yl}-acetyl)-1-methyl-piperazinium chloride,

30 1-[4-methyl-2-(2-methylpropoxy)phenyl]-3-thiazol-2-yl-urea,

{2-[3-(4-methyl-2-[2-methylpropoxy]phenyl)-ureido]-thiazol-4-yl}-acetic acid, or

{2-[3-(4-methyl-2-[2-methylpropoxy]phenyl)-ureido]-thiazol-4-yl}-N-methyl-acetamide.

Embodiment 246: A compound according to any of embodiments 1 to 245, which compound is an activator of glucokinase, when tested in the Glucokinase Activation Assay (I) disclosed

35 herein at a glucose concentration of 2 mM.

Embodiment 247: A compound according to any of embodiments 1 to 246, which compound is an activator of glucokinase, when tested in the Glucokinase Activation Assay (I) disclosed herein at a glucose concentration of from 10 to 15 mM.

5 Embodiment 248: A compound according to any of embodiments 1 to 247, which compound, at a concentration of 30 μ M, is capable of providing an at least 1.5, such as at least 1.7, for instance at least 2.0 fold activation of glucokinase in the Glucokinase Activation Assay (I) disclosed herein at a glucose concentration of 2 mM.

10 Embodiment 249: A compound according to any of embodiments 1 to 248, which compound, at a concentration of 30 μ M, is capable of providing an at least 1.5, such as at least 1.7, for instance at least 2.0 fold activation of glucokinase in the Glucokinase Activation Assay (I) disclosed herein at a glucose concentration of from 10 to 15 mM.

15 Embodiment 250: A compound according to any of embodiments 1 to 249, which at a concentration of 5 μ M is capable of providing an at least 1.5, such as at least 1.7, for instance at least 2.0 fold activation of glucokinase in the Glucokinase Activation Assay (I) disclosed herein at a glucose concentration of 2 mM.

Embodiment 251: A compound according to any of embodiments 1 to 250, which at a concentration of 5 μ M is capable of providing an at least 1.5, such as at least 1.7, for instance at least 2.0 fold activation of glucokinase in the Glucokinase Activation Assay (I) disclosed herein at a glucose concentration of from 10 to 15 mM.

20 Embodiment 252: A glucose kinase activator compound defined as a compound which at a compound concentration of at or below 30 μ M at a glucose concentration of 2 mM in the Glucokinase Activation Assay (I) disclosed herein gives 1.5 - fold higher glucokinase activity than measured at a glucose concentration of 2 mM in the Glucokinase Activation Assay (I) without compound, where the increase in glucokinase activity provided by the compound increases with decreasing concentrations of glucose.

Embodiment 253: A compound according to any of embodiments 1 to 251, which compound provides an increase in glucokinase activity, where the increase in glucokinase activity provided by the compound increases with decreasing concentrations of glucose.

30 Embodiment 254: A compound according to embodiment 252 or embodiment 253, which provides an increase in glucokinase activity in Glucokinase Activation Assay (I) disclosed herein at a glucose concentration of 5 mM, which increase is significantly higher than the increase in glucokinase activity provided by the compound in Glucokinase Activation Assay (I) disclosed herein at a glucose concentration of 15 mM.

35 Embodiment 255: A compound according to any of embodiments 252 to 254, which at a compound concentration of 10 μ M provides an increase in glucokinase activity in

Glucokinase Activation Assay (I) disclosed herein at a glucose concentration of 5 mM, which increase is significantly higher than the increase in glucokinase activity provided by the compound at a compound concentration of 10 μ M in Glucokinase Activation Assay (I) disclosed herein at a glucose concentration of 15 mM.

- 5 Embodiment 256: A compound according to any of embodiments 252 to 255, which at a compound concentration of 10 μ M provides an increase in glucokinase activity in Glucokinase Activation Assay (I) disclosed herein at a glucose concentration of 5 mM, which increase is at least 1.1 fold higher, such as at least 1.2 fold higher, for instance at least 1.3 fold higher, such as at least 1.4 fold higher, for instance 1.5 fold higher, such as at least 1.6 fold higher, for instance at least 1.7 fold higher, such as at least 1.8 fold higher, for instance at least 1.9 fold higher, such as at least 2.0 fold higher than the increase in glucokinase activity provided by the compound at a compound concentration of 10 μ M in Glucokinase Activation Assay (I) disclosed herein at a glucose concentration of 15 mM.
- 10 Embodiment 257: A glucose kinase activator compound defined as a compound which at a compound concentration of at or below 30 μ M at a glucose concentration of 2 mM in the Glucokinase Activation Assay (I) disclosed herein gives 1.5 - fold higher glucokinase activity than measured at a glucose concentration of 2 mM in the Glucokinase Activation Assay (I) without compound, which glucose kinase activator compound increases glucose utilization in the liver without inducing any increase in insulin secretion in response to glucose.
- 15 Embodiment 258: A glucose kinase activator compound defined as a compound which at a compound concentration of at or below 30 μ M at a glucose concentration of 2 mM in the Glucokinase Activation Assay (I) disclosed herein gives 1.5 - fold higher glucokinase activity than measured at a glucose concentration of 2 mM in the Glucokinase Activation Assay (I) without compound, which glucokinase activator compound shows a significantly higher activity in isolated hepatocytes compared to the activity of the glucokinase compound in Ins-1 cells.
- 20 Embodiment 259: A compound according to any of embodiments 1 to 256, which compound increases glucose utilization in the liver without inducing any increase in insulin secretion in response to glucose.
- 25 Embodiment 260: A compound according to any of embodiments 1 to 256, which compound shows a significantly higher activity in isolated hepatocytes compared to the activity of the compound in Ins-1 cells.
- 30 Embodiment 261: A compound according to any of embodiments 257 to 260, which compound shows a significantly higher activity in isolated hepatocytes measured as

described in the Glucokinase Activity Assay (II) compared to the activity of the compound in Ins-1 cells measured as described in the Glucokinase Activity Assay (III).

Embodiment 262: A compound according to embodiment 261, which compound shows an activity in isolated hepatocytes measured as described in the Glucokinase Activity Assay (II)

- 5 which activity is at least 1.1 fold higher, such as at least 1.2 fold higher, for instance at least 1.3 fold higher, such as at least 1.4 fold higher, for instance 1.5 fold higher, such as at least 1.6 fold higher, for instance at least 1.7 fold higher, such as at least 1.8 fold higher, for instance at least 1.9 fold higher, such as at least 2.0 fold higher, for instance at least a 3.0 fold higher, such as at least a 4.0 fold higher, for instance at least 5.0 fold higher, such as at least 10 fold higher than the activity of the compound in Ins-1 cells measured as described in the Glucokinase Activity Assay (III).


Embodiment 263: A compound according to embodiment 261, which compound shows no activity in the Ins-1 cells measured as described in the Glucokinase Activity Assay (III).

- Embodiment 264: A compound according to any one of embodiments 1 to 263, which is an agent useful for the treatment of an indication selected from the group consisting of hyperglycemia, IGT, insulin resistance syndrome, syndrome X, type 2 diabetes, type 1 diabetes, dyslipidemia, hypertension, and obesity.

Embodiment 265: A compound according to any one of embodiments 1 to 264 for use as a medicament.

- 20 Embodiment 266: A compound according to any one of embodiments 1 to 264 for treatment of hyperglycemia, for treatment of IGT, for treatment of Syndrome X, for treatment of type 2 diabetes, for treatment of type 1 diabetes, for treatment of dyslipidemia, for treatment of hyperlipidemia, for treatment of hypertension, for treatment of obesity, for lowering of food intake, for appetite regulation, for regulating feeding behaviour, or for enhancing the secretion of enteroincretins, such as GLP-1.
- 25

In a further aspect the invention provides in an Embodiment A1 a compound of the general formula (I)



(I)

- 5 A¹ is selected from the group consisting of arylene, heteroarylene, fused cycloalkylarylene, fused heterocycliarylene, fused cycloalkylheteroarylene, or fused heterocycliheteroarylene; optionally substituted with one or more substituents R²³, R²⁴, R²⁵, R²⁶, and R²⁷, wherein R²³, R²⁴, R²⁵, R²⁶, and R²⁷ independently of each other are selected from the group consisting of
- 10 halogen, -C(O)OR², -C(O)R², -CN, -CF₃, -OCF₃, -NO₂, -OR², -NR²R³, C₁₋₆-alkyl-Z-, C₂₋₆-alkenyl-Z-, C₂₋₆-alkynyl-Z-, aryl-C₁₋₆-alkylene-Z-, heteroaryl-C₁₋₆-alkylene-Z-, heterocycliyl-C₁₋₆-alkylene-Z-, cycloalkyl-C₁₋₆-alkylene-Z-, N(R⁴R⁵)-C₁₋₆-alkylene-Z-, R⁶-W¹-Z-, R⁶-W¹-C₁₋₆-alkylene-Z-, R⁶-W¹-C₁₋₆-arylene-Z-, R⁶-W¹-heteroarylene-Z-, R⁶-W¹-heterocycliylene-Z-, R⁶-W¹-N(R⁴)-Z-, R⁶-N(R⁴)-Z, R⁶-W¹-C₁₋₆-alkylene-Z-,
- 15 heterocycliyl-Z-C₁₋₆-alkylene-, heterocycliyl-C₁₋₆-alkylene-Z-C₁₋₆-alkylene-, C₃₋₁₀-cycloalkyl-Z-C₁₋₆-alkylene-, C₃₋₁₀-cycloalkyl-C₁₋₆-alkylene-Z-C₁₋₆-alkylene-, C₃₋₁₀-cycloalkyl-arylene-Z-, heterocycliyl-arylene-Z-, C₃₋₁₀-cycloalkyl-heteroarylene-Z-, heterocycliyl-heteroarylene-Z-, aryl-C₃₋₁₀-cycloalkylene-Z-, aryl-heterocycliylene-Z-, heteroaryl-C₃₋₁₀-cycloalkylene-Z-, heteroaryl-heterocycliylene-Z-, heterocycliyl-C₃₋₁₀-cycloalkylene-Z-, aryl-heteroarylene-Z-, heteroaryl-arylene-Z-, aryl-arylene-Z-, heteroaryl-heteroarylene-Z-, wherein any mono- or divalent C₁₋₆-alkyl, C₃₋₁₀-cycloalkyl, heterocycliyl, aryl or heteroaryl moiety can optionally be substituted with one or more substituents independently selected from R², and wherein
- 25 R² and R³ independently of each other are hydrogen, halogen, hydroxy, -CN, -CF₃, -OCF₃, -NO₂, -C(O)OH, -NH₂, C₁₋₆-alkyl, C₁₋₆-alkoxy, aryloxy, R⁶-Z-, aryl-C₁₋₆-alkylene-, heteroaryl-C₁₋₆-alkylene-, C₁₋₆-alkyl-arylene-, C₁₋₆-alkyl-heteroarylene-, heteroaryl, or aryl;
- or

R^2 and R^3 , when attached to the same nitrogen atom, together with said nitrogen atom may form a 3 to 8 membered heterocyclic ring optionally containing one or two further heteroatoms selected from nitrogen, oxygen and sulfur, and optionally containing one or two double bonds;

5

Z and W¹ independently of each other are a direct bond, -O-, -N(R^7)-, -N(R^7)C(R^7R^8)-, -S-, -SO₂-, -C(O)N(R^7)-, -N(R^7)C(O)-, -N(R^7)C(O)C(R^7R^8)-, -N(R^7)CON(R^8)-, -N(R^7)SO₂-, -SO₂N(R^7)-, -C(O)-, -C(O)-O-, -N(R^7)SO₂N(R^8)-, or -O-C(O)-, wherein

10

R^7 and R^8 in each individual case independently of each other are hydrogen or C₁₋₆-alkyl; and

R^4 , R^5 , and R^6 independently of each other are selected from the group consisting of hydrogen, cyano, halogen, aryl, heteroaryl, heteroaryl-C₁₋₆-alkylene-, aryl-C₁₋₆-alkylene-, C₃₋₁₀-cycloalkyl, heterocyclyl optionally substituted with one or more C₁₋₆-alkyl, or C₁₋₆-alkyl optionally substituted with halogen, -S(O)₂CH₃ or COOH; or


15

R^4 and R^5 may be taken together to form a ring having the formula -(CH₂)_j-Q-(CH₂)_k- bonded to the nitrogen atom to which R^4 and R^5 are attached, wherein

j and k independently of each other is 1, 2, 3, or 4; and

20

Q is a direct bond, -CH₂-, -O-, -S-, -S(O₂)-, -C(O)-, -C(O)NH-, -NHC(O)-, -NHC(O)NH-, -NHSO₂-, -SO₂NH-, -C(O)-O-, -O-C(O)-, -NHSO₂NH-,



wherein

25

R^9 and R^{10} independently of each other are selected from the group consisting of hydrogen, aryl, C₁₋₆-alkyl, and aryl-alkylene-;

30

L¹ is a bond, -D-C₁₋₆-alkylene-E-, -D-C₂₋₆-alkenylene-E-, -D-C₂₋₆-alkynylene-E-, -D-cycloalkylene-E-, -D-heterocyclylene-E-, -O-, -S-, -S(O)-, -S(O)₂-,

-C(O)-, -N(R¹¹)-, or -C(=N-OR¹²)-, wherein

D and E independently of each other are a direct bond, -O- or -S-;

R¹¹ is selected from hydrogen, C₁₋₆-alkyl, aryl, carbamoyl, aryl-C₁₋₆-alkylene-,

heteroaryl-C₁₋₆-alkylene-, C₁₋₆-alkyl-O-C(O)-, aryl-C₁₋₆-alkylene-O-C(O)-,

5 heteroaryl-C₁₋₆-alkylene-O-C(O)-, C₁₋₆-alkyl-NH-C(O)-, aryl-C₁₋₆-alkylene-NH-C(O)-,

heteroaryl-C₁₋₆-alkylene-NH-C(O)-, C₁₋₆-alkyl-SO₂-, aryl-C₁₋₆-alkylene-SO₂-,

heteroaryl-C₁₋₆-alkylene-SO₂-, aryl-SO₂-, heteroaryl-SO₂-, C₁₋₆-alkyl-NH-SO₂-,

aryl-C₁₋₆-alkylene-NH-SO₂-, heteroaryl-C₁₋₆-alkylene-NH-SO₂-, C₁₋₆-alkyl-C(O)-,

aryl-C₁₋₆-alkylene-C(O)-, heteroaryl-C₁₋₆-alkylene-C(O)-, C₁₋₆-alkyl-Y-, aryl-Y-,

10 heteroaryl-Y-, aryl-C₁₋₆-alkylene-Y-, heteroaryl-C₁₋₆-alkylene-Y-,

N(R¹³)(R¹⁴)-C₁₋₆-alkylene-Y-, and R¹⁵-W²-C₁₋₆-alkylene-Y-, wherein

Y and W² independently of each other are a direct bond, -CH₂-, -SO₂-,

-N(H)CO-, -N(H)SO₂-, or -O-C(O)-;

R¹³ and R¹⁴ independently of each other are selected from hydrogen, aryl,

15 heteroaryl, C₁₋₆-alkyl, C₁₋₆-alkoxy, aryl-C₁₋₆-alkylene-,

heteroaryl-C₁₋₆-alkylene-, aryl-C₁₋₆-alkoxy-, heteroaryl-C₁₋₆-alkoxy-,

C₁₋₆-alkyl-arylene-, C₁₋₆-alkyl-heteroarylene-, C₁₋₆-alkoxy-heteroarylene-, or

C₁₋₆-alkoxy-arylene-;

or

20 R₁₃ and R₁₄ may be taken together to form a ring having the formula

-(CH₂)_o-X-(CH₂)_p- bonded to the nitrogen atom to which R₁₃ and R₁₄ are attached, wherein


o and p are independently of each other are 1, 2, 3, or 4; and

X is a direct bond, -CH₂-, -O-, -S-, -S(O₂)-, -C(O)-, -CON(H)-,

25 -NHC(O)-, -NHCON(H)-, -NHSO₂-, -SO₂N(H)-, -C(O)-O-, -O-C(O)-,

-NHSO₂NH-,

72



wherein

R^{16} and R^{17} are selected from hydrogen, aryl, heteroaryl, C_{1-6} -alkyl, C_{1-6} -alkoxy, aryl- C_{1-6} -alkylene-, heteroaryl- C_{1-6} -alkylene-, C_{1-6} -alkyl-arylene-, C_{1-6} -alkyl-heteroarylene-, C_{1-6} -alkoxy-arylene-, C_{1-6} -alkoxy-heteroarylene-, heteroarylaryl- C_{1-6} -alkoxy-, or aryl- C_{1-6} -alkoxy-; and

R^{15} is selected from the group consisting of aryl, heteroaryl, cycloalkyl,

heterocyclyl, C_{1-6} -alkyl, heteroaryl- C_{1-6} -alkylene-, or aryl- C_{1-6} -alkylene-; and

R^{12} is selected from hydrogen, aryl, heteroaryl, C_{1-6} -alkyl, aryl- C_{1-6} -alkylene-, heteroaryl- C_{1-6} -alkylene-, C_{1-6} -alkyl-arylene-, C_{1-6} -alkyl-heteroarylene-, C_{1-6} -alkoxy-heteroarylene-, or C_{1-6} -alkoxy-arylene-;

G^1 is C_{1-6} -alkyl, C_{3-10} -cycloalkyl, C_{3-10} -cycloalkyl- C_{1-6} -alkylene-, C_{2-6} -alkenyl, or C_{2-6} -alkynyl, all of which may optionally be substituted with one or more substituents independently selected from the group consisting of -CN, -CF₃, -OCF₃, -OR¹⁸, -NR¹⁸R¹⁹, C_{3-10} -cycloalkyl and C_{1-6} -alkyl, wherein

R^{18} and R^{19} independently of each other are hydrogen, C_{1-6} -alkyl, heteroaryl- C_{1-6} -alkylene-, aryl- C_{1-6} -alkylene-, C_{1-6} -alkyl-arylene-, C_{1-6} -alkyl-heteroarylene-, heteroaryl, or aryl;

or

R^{18} and R^{19} , when attached to the same nitrogen atom, together with the said nitrogen atom may form a 3 to 8 membered heterocyclic ring optionally containing one or two further heteroatoms selected from nitrogen, oxygen and sulfur, and optionally containing one or two double bonds;

or

G¹ is aryl, heteroaryl, heterocyclyl, fused cycloalkylheteroaryl, fused heterocyclylaryl, fused arylheterocyclyl, or fused cycloalkylaryl, all of which may optionally be substituted with one or more substituents selected from R⁴⁰, R⁴¹, and R⁴²;

- 5 L² is a direct bond, C₁₋₆-alkylene, C₂₋₆-alkenylene, C₂₋₆-alkynylene, -N(R²⁰)-, -C₁₋₆-alkylene-N(R²⁰)-, -C₂₋₆-alkenylene-N(R²⁰)-, -C₂₋₆-alkynylene-N(R²⁰)-, wherein

R²⁰ is hydrogen, or

R²⁰ is C₁₋₆-alkyl, C₂₋₆-alkenyl, C₂₋₆-alkynyl, cycloalkyl-W³-, heterocyclyl-W³-, aryl-W³-, heteroaryl-W³-, optionally substituted with one or more substituents R³⁰, R³¹, and

- 10 R³², wherein

W³ is C₁₋₆-alkylene or a direct bond;

wherein L¹ and L² are attached to adjacent atoms in A¹;

L³ is -C(O)-, -C(O)-C(O)-, -C(O)CH₂C(O)- or -S(O)₂;

R¹ is hydrogen, or

- 15 R¹ is C₁₋₆-alkyl, C₂₋₆-alkenyl, C₂₋₆-alkynyl, cycloalkyl-W⁴-, heterocyclyl-W⁴-, aryl-W⁴-, or heteroaryl-W⁴-,


optionally substituted with one or more substituents R³³, R³⁴, and R³⁵, wherein

W⁴ is C₁₋₆-alkylene or a direct bond;

G² is heteroaryl, fused heterocyclylheteroaryl, or fused cycloalkylheteroaryl,

- 20 optionally substituted with one or more substituents R⁴³, R⁴⁴, and R⁴⁵, wherein said heteroaryl group possesses a nitrogen atom adjacent to the atom joining said heteroaryl group to -N(R¹)-;

or a group of the formula



- 25 wherein

G³ and G⁵ independently of each other are C₁₋₆-alkyl, C₂₋₆-alkenyl, C₂₋₆-alkynyl, cycloalkyl-R²²-, heterocyclyl-R²²-, aryl-R²²-, heteroaryl-R²²-,

optionally substituted with one or more substituents R⁴⁶, R⁴⁷, and R⁴⁸, wherein

R²² is alkylene or a direct bond; and

- 30 R²¹ is hydrogen, C₁₋₆-alkyl, C₂₋₆-alkenyl, C₂₋₆-alkynyl, cycloalkyl-W⁵-, or heterocyclyl-W⁵-,

optionally substituted with one or more substituents R³⁶, R³⁷, and R³⁸, or

R^{21} is aryl- W^5 -, or heteroaryl- W^5 -, optionally substituted with one or more substituents R^{49} , R^{50} , and R^{51} , wherein

W^5 is C_{1-8} -alkylene or a direct bond;

wherein

- 5 R^{30} , R^{31} , R^{32} , R^{33} , R^{34} , R^{35} , R^{36} , R^{37} , and R^{38} independently of each other are selected from $-CHF_2$, $-CF_3$, $-OCF_3$, $-OCHF_2$, $-OCH_2CF_3$, $-OCF_2CHF_2$, $-S(O)_2CF_3$, $-SCF_3$, $-OR^{52}$, $-NR^{52}R^{53}$, $-SR^{52}$, $-NR^{52}S(O)_2R^{53}$, $-S(O)_2NR^{52}R^{53}$, $-S(O)NR^{52}R^{53}$, $-S(O)R^{52}$, $-S(O)_2R^{52}$, $-C(O)NR^{52}R^{53}$, $-OC(O)NR^{52}R^{53}$, $-NR^{52}C(O)R^{53}$, $-CH_2C(O)NR^{52}R^{53}$, $-OCH_2C(O)NR^{52}R^{53}$, $-CH_2OR^{52}$, $-CH_2NR^{52}R^{53}$, $-OC(O)R^{52}$, $-C(O)R^{52}$ and $-C(O)OR^{52}$; or
- 10 C_{2-8} -alkenyl and C_{2-8} -alkynyl, which may optionally be substituted with one or more substituents selected from $-CN$, $-CF_3$, $-OCF_3$, $-OR^{52}$, $-NR^{52}R^{53}$ and C_{1-8} -alkyl; or C_{3-10} -cycloalkyl, C_{4-8} -cycloalkenyl, heterocyclyl, C_{3-10} -cycloalkyl- C_{1-8} -alkylene-, C_{3-10} -cycloalkyl- C_{1-8} -alkoxy-, C_{3-10} -cycloalkyloxy, C_{3-10} -cycloalkyl- C_{1-8} -alkylthio-, C_{3-10} -cycloalkylthio, C_{3-10} -cycloalkyl- C_{2-8} -alkenylene-, C_{3-10} -cycloalkyl- C_{2-8} -alkynylene-,
- 15 C_{4-8} -cycloalkenyl- C_{1-8} -alkylene-, C_{4-8} -cycloalkenyl- C_{2-8} -alkenylene-, C_{4-8} -cycloalkenyl- C_{2-8} -alkynylene-, heterocyclyl- C_{1-8} -alkylene-, heterocyclyl- C_{2-8} -alkenylene-, heterocyclyl- C_{2-8} -alkynylene-, aryl, aryloxy, aryloxycarbonyl, aroyl, aryl- C_{1-8} -alkoxy-, aryl- C_{1-8} -alkylene-, aryl- C_{2-8} -alkenylene-, aryl- C_{2-8} -alkynylene-, heteroaryl, heteroaryl- C_{1-8} -alkylene-, heteroaryl- C_{2-8} -alkenylene- and heteroaryl- C_{2-8} -alkynylene-, of
- 20 which the aryl and heteroaryl moieties optionally may be substituted with one or more substituents selected from halogen, $-C(O)OR^{52}$, $-CN$, $-CF_3$, $-OCF_3$, $-NO_2$, $-OR^{52}$, $-NR^{52}R^{53}$ and C_{1-8} -alkyl, wherein R^{52} and R^{53} independently of each other are hydrogen, C_{1-8} -alkyl, aryl- C_{1-8} -alkylene-, heteroaryl- C_{1-8} -alkylene- heteroaryl, or aryl;
- 25 or R^{52} and R^{53} , when attached to the same nitrogen atom, together with the said nitrogen atom may form a 3 to 8 membered heterocyclic ring optionally containing one or two further heteroatoms selected from nitrogen, oxygen and sulfur, and optionally containing one or two double bonds;
- 30 R^{40} , R^{41} , R^{42} , R^{43} , R^{44} , R^{45} , R^{46} , R^{47} , R^{48} , R^{49} , R^{50} and R^{51} independently of each other are $-CN$, $-NO_2$, $-S(O)_2CF_3$, $-SCF_3$, $-OR^{54}$, $-NR^{54}R^{55}$, $-SR^{54}$, $-NR^{54}S(O)_2R^{55}$, $-S(O)_2NR^{54}R^{55}$, $-S(O)NR^{54}R^{55}$, $-S(O)R^{54}$, $-S(O)_2R^{54}$, $-C(O)NR^{54}R^{55}$, $-OC(O)NR^{54}R^{55}$, $-NR^{54}C(O)R^{55}$, halogen, $-S-C_{1-8}$ -alkylene- OR^{54} , $-S(O)_2-C_{1-8}$ -alkylene- OR^{54} , $-C_{1-8}$ -alkylene- $S-R^{54}$, $-C_{1-8}$ -alkylene- $S(O)R^{54}$, $-C_{1-8}$ -alkylene- $S(O)_2R^{54}$, $-C_{1-8}$ -alkylene- $N(R^{54})S(O)_2R^{55}$,
- 35 $-N(R^{54})S(O)_2R^{55}$, $-C_{1-8}$ -alkylene- CN ,

- $-C_{1-6}$ -alkylene- $C(O)NR^{54}R^{55}$, $-C_{1-6}$ -alkylene- $N(R^{54})C(O)R^{55}$, $-N(R^{54})C(O)R^{55}$,
 $-C_{1-6}$ -alkylene- $N(R^{54})C(O)NR^{55}R^{56}$, $-C_{1-6}$ -alkylene- $NHC(=NR^{54})NR^{55}R^{56}$,
 $-C_{1-6}$ -alkylene- $N(R^{54})C(O)OR^{55}$, $-N(R^{54})C(O)OR^{55}$, $-C_{1-6}$ -alkylene- $C(O)OR^{54}$,
 $-C_{1-6}$ -alkylene- $N(R^{54})S(O)_2R^{55}$, $-OCH_2C(O)NR^{54}R^{55}$, $-O(CH_2)_{1-3}OR^{54}$, $-C_{1-6}$ -alkylene- $O-R^{54}$,
5 $-C_{1-6}$ -alkylene- $C(O)R^{54}$, $-C_{1-6}$ -alkylene- $NR^{54}R^{55}$, $-C(=NR^{54})-O-R^{55}$, $-C(=N(OR^{54}))C(O)OR^{55}$,
 $-C(=N(OR^{54}))C(O)R^{55}$, $-C_{1-6}$ -alkylene- $C(=N(OR^{54}))C(O)R^{55}$, $-C_{1-6}$ -alkylene- $N-O-R^{54}$,
 $-C_{1-6}$ -alkylene- $N(R^{54})S(O)_2NR^{55}R^{56}$, $-N(R^{54})S(O)_2NR^{55}R^{56}$, $-N(R^{54})S(O)_2NR^{55}R^{56}$, $-OC(O)R^{54}$,
 $-C_{1-6}$ -alkylene- $C(O)N(R^{54})S(O)_2R^{55}$, $-C(O)N(R^{54})S(O)_2R^{55}$, $-C_{1-6}$ -alkylene- $C(R^{54})=N-OR^{55}$,
 $-NHC(=NR^{54})NR^{55}R^{56}$, $-C_{1-6}$ -alkylene- $NHC(=NR^{54})NR^{54}R^{55}$, $-C_{1-6}$ -alkylene- $N=C(N(R^{54}R^{55}))_2$,
10 $-N=C(N(R^{54}R^{55}))_2$, $-C(O)R^{54}$ and $-C(O)OR^{54}$; or
 C_{1-6} -alkyl, C_{2-6} -alkenyl and C_{2-6} -alkynyl, each of which may optionally be substituted with one
or more substituents independently selected from halogen, R^{54} , $-CN$, $-CF_3$, $-OCF_3$, $-OR^{54}$, $-$
 $C(O)OR^{54}$, $-NR^{54}R^{55}$ and C_{1-6} -alkyl; or
 C_{3-10} -cycloalkyl, C_{4-8} -cycloalkenyl, heterocyclyl, C_{3-10} -cycloalkyl- C_{1-6} -alkylene-,
15 C_{3-10} -cycloalkyl- C_{1-6} -alkoxy-, C_{3-10} -cycloalkyloxy, C_{3-10} -cycloalkyl- C_{1-6} -alkylthio-,
 C_{3-10} -cycloalkylthio, C_{3-10} -cycloalkyl- C_{2-6} -alkenylene-, C_{3-10} -cycloalkyl- C_{2-6} -alkynylene-,
 C_{4-8} -cycloalkenyl- C_{1-6} -alkylene-, C_{4-8} -cycloalkenyl- C_{2-6} -alkenylene-,
 C_{4-8} -cycloalkenyl- C_{2-6} -alkynylene-, heterocyclyl- C_{1-6} -alkylene-, heterocyclyl- C_{2-6} -alkenylene-,
heterocyclyl- C_{2-6} -alkynylene- of which the heterocyclyl moieties optionally may be substituted
20 with one or more substituents independently selected from R^{70} ; or
aryl, aryloxy, aryloxycarbonyl, aroyl, aryl- C_{1-6} -alkoxy-, aryl- C_{1-6} -alkylene-,
aryl- C_{2-6} -alkenylene-, aryl- C_{2-6} -alkynylene-, heteroaryl, heteroaryl- C_{1-6} -alkylene-, heteroaryl-S-
 C_{1-6} -alkylene-, heteroaryl- C_{2-6} -alkenylene- and heteroaryl- C_{2-6} -alkynylene-, of which the aryl
and heteroaryl moieties optionally may be substituted with one or more substituents selected
25 from halogen, $-C(O)OR^{54}$, $-CN$, $-CF_3$, $-OCF_3$, $-NO_2$, $-OR^{54}$, $-NR^{54}R^{55}$ or C_{1-6} -alkyl, wherein
 R^{54} , R^{55} and R^{56} independently of each other are hydrogen, C_{1-6} -alkyl, $-Si(C_{1-6}\text{-alkyl})_3$,
 C_{1-6} -alkyl-arylene-, C_{1-6} -alkyl-heteroarylene-, aryl- C_{1-6} -alkylene-,
heteroaryl- C_{1-6} -alkylene-, heterocyclyl, heterocyclyl- C_{1-6} -alkylene-, heteroaryl, or aryl,
each of which is optionally substituted with one or more substituents independently
30 selected from R^{71} ;
or
 R^{54} and R^{55} independently of each other are hydrogen or $-(CHR^{72})_u-(CHR^{73})_v-W^8$,
wherein
35 u is 0, 1 or 2;
 v is 0, 1 or 2;

R^{72} and R^{73} independently of each other are hydrogen, C_{1-6} -alkyl, C_{1-6} -alkyl-arylene-aryl, hydroxy, hydroxyalkyl, $-C(O)O-R^{75}$, amino, or aminoalkyl;

W^6 is hydrogen, $-O-R^{75}$, $-C(O)O-R^{75}$, $-C(O)-R^{75}$, $-CONR^{75}R^{76}$, $-NR^{75}R^{76}$, $-NHCH_2C(O)R^{75}$, $-NHC(O)R^{75}$, $-NHC(O)OR^{75}$, $-S(O)_2R^{75}$, $-NHS(O)_2R^{75}$, alkylamino, or dialkylamino, or

W^6 is a five or six membered ring wherein at least one ring atom is nitrogen and the remaining ring atoms are either carbon or oxygen or optionally substituted with C_{1-6} -alkyl, $-C(O)O-R^{75}$, $-(CH_2)_{1-3}C(O)O-R^{75}$, $=O$;

or

W^6 is phthalimido or heterocyclyl.

or

R^{54} and R^{55} , when attached to the same nitrogen atom, together with the said nitrogen atom may form a 3 to 8 membered heterocyclic ring optionally containing one or two further heteroatoms selected from nitrogen, oxygen and sulfur, and optionally containing one or two double bonds, and optionally substituted with one or more C_{1-6} -alkyl groups;

R^{70} is $=O$, $-C(O)CH_3$, $-S(O)_2CH_3$, $-CF_3$, $-C(O)O-R^{75}$, $-(CH_2)_{1-3}C(O)O-R^{75}$ or C_{1-6} -alkyl;

R^{71} is $=O$, C_{1-6} -alkyl, cycloalkyl, $-C(O)O-R^{75}$, $-(CH_2)_{1-3}C(O)O-R^{75}$, $-(CH_2)_{1-3}NR^{75}R^{76}$, $-OH$ or amino;

R^{75} and R^{76} independently of each other is hydrogen, halogen, $-OH$, $-CF_3$, or C_{1-6} -alkyl optionally substituted with $-NH_2$;

or a pharmaceutically acceptable salt, solvate, or prodrug thereof

Embodiment A2. A compound according to embodiment A1, wherein

A^1 is arylene or heteroarylene, optionally substituted with one or more substituents R^{23} , R^{24} , R^{25} , R^{26} , and R^{27} , wherein

R^{23} , R^{24} , R^{25} , R^{26} , and R^{27} independently of each other are selected from the group consisting of

halogen, $-C(O)OR^2$, $-CN$, $-CF_3$, $-OCF_3$, $-NO_2$, $-OR^2$, $-NR^2R^3$, C_{1-6} -alkyl-Z-, C_{2-6} -alkenyl-Z-, C_{2-6} -alkynyl-Z-, cycloalkyl-Z-, heterocyclyl-Z-, aryl-Z-, heteroaryl-Z-, aryl- C_{1-6} -alkylene-Z-, heteroaryl- C_{1-6} -alkylene-Z-, heterocyclyl- C_{1-6} -alkylene-Z-,

cycloalkyl-C₁₋₆-alkylene-Z-, N(R⁴R⁵)-C₁₋₆-alkylene-Z-, R⁶-W¹-Z-, R⁶-W¹-N(R⁴)-Z-, R⁶-N(R⁴)-Z-, and R⁶-W¹-C₁₋₆-alkylene-Z-, wherein

R², R³, R⁴, R⁵, R⁶, Z, and W¹ are as defined in embodiment A1.

Embodiment A3. A compound according to embodiment A2, wherein

- 5 A¹ is C₆₋₁₀-arylene or C₄₋₁₀-heteroarylene, optionally substituted with one or more substituents R²³, R²⁴, R²⁵, R²⁶, and R²⁷, wherein

R²³, R²⁴, R²⁵, R²⁶, and R²⁷ independently of each other are selected from the group consisting of

- 10 halogen, -C(O)OR², -CN, -CF₃, -OCF₃, -NO₂, -OR², -NR²R³, C₁₋₆-alkyl-Z-, C₂₋₆-alkenyl-Z-, C₂₋₆-alkynyl-Z-, cycloalkyl-Z-, heterocyclyl-Z-, aryl-Z-, heteroaryl-Z-, aryl-C₁₋₆-alkylene-Z-, heteroaryl-C₁₋₆-alkylene-Z-, heterocyclyl-C₁₋₆-alkylene-Z-, cycloalkyl-C₁₋₆-alkylene-Z-, N(R⁴R⁵)-C₁₋₆-alkylene-Z-, R⁶-W¹-Z-, R⁶-W¹-N(R⁴)-Z-, R⁶-N(R⁴)-Z-, and R⁶-W¹-C₁₋₆-alkylene-Z-, wherein

R², R³, R⁴, R⁵, R⁶, Z, and W¹ are as defined in embodiment A1.

15

Embodiment A4. A compound according to embodiment A3 wherein

A¹ is phenylene optionally substituted with one or more substituents R²³, R²⁴, R²⁵, R²⁶, and R²⁷, wherein


R²³, R²⁴, R²⁵, R²⁶, and R²⁷ independently of each other are selected from the group consisting of

- 20 halogen, -C(O)OR², -CN, -CF₃, -OCF₃, -NO₂, -OR², -NR²R³, C₁₋₆-alkyl-Z-, C₂₋₆-alkenyl-Z-, C₂₋₆-alkynyl-Z-, cycloalkyl-Z-, heterocyclyl-Z-, aryl-Z-, heteroaryl-Z-, aryl-C₁₋₆-alkylene-Z-, heteroaryl-C₁₋₆-alkylene-Z-, heterocyclyl-C₁₋₆-alkylene-Z-, cycloalkyl-C₁₋₆-alkylene-Z-, N(R⁴R⁵)-C₁₋₆-alkylene-Z-, R⁶-W¹-Z-, R⁶-W¹-N(R⁴)-Z-, R⁶-N(R⁴)-Z-, and R⁶-W¹-C₁₋₆-alkylene-Z-, wherein

25

R², R³, R⁴, R⁵, R⁶, Z, and W¹ are as defined in embodiment A1.


Embodiment A5. A compound according to embodiment A4 of the formula (Ia)



Formula (Ia)

wherein L^1 , G^1 , L^2 , L^3 , R^1 , G^2 and R^{25} are as defined in embodiment A1.

Embodiment A6. A compound according to embodiment A4 of the formula (Ib)



Formula (Ib)

wherein L^1 , G^1 , L^2 , L^3 , R^1 , G^2 and R^{24} are as defined in embodiment A1.

Embodiment A7. A compound according to any one of the embodiments A1 to A6, wherein R^{23} , R^{24} , R^{25} , R^{26} , and R^{27} independently of each other are selected from the group consisting of

- 10 halogen, $-C(O)OR^2$, $-CN$, $-CF_3$, $-OCF_3$, $-NO_2$, $-OR^2$, $-NR^2R^3$, C_{1-6} -alkyl-Z-, cycloalkyl-Z-, heterocyclyl-Z-, aryl-Z-, or heteroaryl-Z-, $N(R^4R^5)$ - C_{1-6} -alkylene-Z-, R^6 - W^1 -Z-, R^6 - W^1 - $N(R^4)$ -Z-, R^6 - $N(R^4)$ -Z-, and R^6 - W^1 - C_{1-6} -alkylene-Z-, wherein R^2 , R^3 , R^4 , R^5 , R^6 , Z, and W^1 are as defined in embodiment A1.

Embodiment A8. A compound according to embodiment A7 wherein

- 15 R^{23} , R^{24} , R^{25} , R^{26} , and R^{27} independently of each other are selected from the group consisting of halogen, $-CN$, $-CF_3$, $-OR^2$, $-NR^2R^3$, C_{1-6} -alkyl-Z-, cycloalkyl-Z-, heterocyclyl-Z-, aryl-Z-, or heteroaryl-Z-, R^6 - W^1 -Z-, R^6 - W^1 - $N(R^4)$ -Z-, R^6 - $N(R^4)$ -Z-, and R^6 - W^1 - C_{1-6} -alkylene-Z-, wherein

- 20 R^2 , R^3 , R^4 , R^5 , R^6 , Z, and W^1 are as defined in embodiment A1.

Embodiment A9. a compound according to embodiment A8 wherein

R^{23} , R^{24} , R^{25} , R^{26} , and R^{27} independently of each other are selected from the group consisting of

halogen, $-OR^2$, $-NR^2R^3$, C_{1-6} -alkyl-Z-, R^6-W^1-Z -, and $R^6-W^1-C_{1-6}$ -alkylene-Z-, wherein R^2 , R^3 , R^4 , R^5 , R^6 , Z, and W^1 are as defined in embodiment A1.

Embodiment A10. A compound according to embodiment A9 wherein

5 R^{23} , R^{24} , R^{25} , R^{26} , and R^{27} independently of each other are selected from the group consisting of F, Cl, Br, and methyl.

Embodiment A11. A compound according to any one of the embodiments A1 to A10, wherein R^4 , R^5 , and R^6 independently of each other are selected from the group consisting of hydrogen, aryl, C_{1-6} -alkyl, heteroaryl- C_{1-6} -alkylene-, and aryl- C_{1-6} -alkylene-.

10 Embodiment A12. A compound according to embodiment A11, wherein

R^4 , R^5 , and R^6 independently of each other are hydrogen or C_{1-6} -alkyl.

Embodiment A13. A compound according to embodiment A12, wherein


R^4 , R^5 , and R^6 are hydrogen.

Embodiment A14. A compound according to any one of the embodiments A1 to A9, wherein

15 R^4 and R^5 is taken together to form a ring having the formula $-(CH_2)_j-Q-(CH_2)_k$ - bonded to the nitrogen atom to which R^4 and R^5 are attached, wherein

j and k independently of each other is 1, 2, 3, or 4;

Q is a direct bond, $-CH_2-$, $-O-$, $-S-$, $-S(O_2)-$, $-C(O)-$, $-C(O)NH-$, $-NHC(O)-$, $-NHC(O)NH-$, $-NHSO_2-$, $-SO_2NH-$, $-C(O)-O-$, $-O-C(O)-$, $-NHSO_2NH-$,




20

wherein

R^9 and R^{10} independently of each other are selected from the group consisting of hydrogen, aryl, C_{1-6} -alkyl, and aryl- C_{1-6} -alkyl-.

Embodiment A15. A compound according to embodiment A14, wherein

Q is a direct bond, $-\text{CH}_2-$, $-\text{O}-$, $-\text{S}-$, $-\text{S}(\text{O}_2)-$, $-\text{C}(\text{O})-$, $-\text{C}(\text{O})\text{NH}-$, $-\text{NHC}(\text{O})-$, $-\text{NHC}(\text{O})\text{NH}-$, $-\text{NHSO}_2-$, $-\text{SO}_2\text{NH}-$, $-\text{C}(\text{O})-\text{O}-$, $-\text{O}-\text{C}(\text{O})-$, $-\text{NHSO}_2\text{NH}-$,




wherein

- 5 R^9 and R^{10} independently of each other are hydrogen or C_{1-8} -alkyl.

Embodiment A16. A compound according to embodiment A15, wherein

Q is a direct bond, $-\text{CH}_2-$, $-\text{O}-$, $-\text{S}-$, $-\text{S}(\text{O}_2)-$, $-\text{C}(\text{O})-$, $-\text{C}(\text{O})\text{NH}-$, $-\text{NHC}(\text{O})-$, $-\text{NHC}(\text{O})\text{NH}-$, $-\text{NHSO}_2-$, $-\text{SO}_2\text{NH}-$, $-\text{C}(\text{O})-\text{O}-$, $-\text{O}-\text{C}(\text{O})-$, $-\text{NHSO}_2\text{NH}-$,



- 10 Embodiment A17. A compound according to embodiment A16, wherein Q is a direct bond.

Embodiment A18. A compound according to any one of the embodiments A1 to A17, wherein

W^1 is a direct bond, $-\text{O}-$, $-\text{C}(\text{O})-$, $-\text{NH}-$, $-\text{S}-$, $-\text{SO}_2-$, $-\text{C}(\text{O})\text{NH}-$, $-\text{NHC}(\text{O})-$, $-\text{N}(\text{H})\text{CON}(\text{H})-$, $-\text{N}(\text{H})\text{SO}_2-$, $-\text{SO}_2\text{N}(\text{H})-$, $-\text{C}(\text{O})-\text{O}-$, $-\text{N}(\text{H})\text{SO}_2\text{N}(\text{H})-$, or $-\text{O}-\text{C}(\text{O})-$.

Embodiment A19. A compound according to embodiment A18, wherein

- 15 W^1 is a direct bond, $-\text{O}-$, $-\text{C}(\text{O})-$, $-\text{SO}_2-$, $-\text{C}(\text{O})\text{NH}-$, $-\text{NHC}(\text{O})-$, $-\text{N}(\text{H})\text{SO}_2-$, $-\text{C}(\text{O})-\text{O}-$, or $-\text{O}-\text{C}(\text{O})-$.

Embodiment A20. A compound according to embodiment A19, wherein W^1 is a direct bond or $-\text{C}(\text{O})-\text{O}-$.

Embodiment A21. A compound according to embodiment A20, wherein W^1 is a direct bond.

Embodiment A22. A compound according to embodiment A7, wherein

at least one of R^{23} , R^{24} , R^{25} , R^{26} , and R^{27} is halogen, $-C(O)OR^2$, $-CN$, $-CF_3$, $-OCF_3$, $-NO_2$, $-OR^2$, $-NR^2R^3$, C_{1-6} -alkyl-Z-, cycloalkyl-Z-, heterocyclyl-Z-, aryl-Z-, or heteroaryl-Z-, wherein

R^2 , R^3 , and Z are as defined in embodiment A1.

Embodiment A23. A compound according to embodiment A22, wherein

at least one of R^{23} , R^{24} , R^{25} , R^{26} , and R^{27} is halogen, $-C(O)OR^2$, $-CN$, $-CF_3$, $-OCF_3$, $-NO_2$, $-OR^2$, $-NR^2R^3$, C_{1-6} -alkyl-Z-, C_{3-10} -cycloalkyl-Z-, C_{3-10} -heterocyclyl-Z-, C_{3-10} -aryl-Z-, or C_{3-10} -heteroaryl-Z-, wherein

R^2 , R^3 , and Z are as defined in embodiment A1.

Embodiment A24. A compound according to embodiment A23, wherein

at least one of R^{23} , R^{24} , R^{25} , R^{26} , and R^{27} is halogen, $-C(O)OR^2$, $-CN$, $-NO_2$, $-OR^2$, $-NR^2R^3$, C_{1-6} -alkyl-Z-, C_{3-10} -cycloalkyl-Z-, C_{3-10} -heterocyclyl-Z-, C_{3-10} -aryl-Z-, or C_{3-10} -heteroaryl-Z-, wherein

R^2 , R^3 , and Z are as defined in embodiment A1.

Embodiment A25. A compound according to embodiment A24, wherein

at least one of R^{23} , R^{24} , R^{25} , R^{26} , and R^{27} is F, Cl, Br, or methyl.

Embodiment A26. A compound according to embodiment A24, wherein

at least one of R^{23} , R^{24} , R^{25} , R^{26} , and R^{27} is halogen, $-C(O)OR^2$, $-CN$, $-NO_2$, $-OR^2$, or $-NR^2R^3$, wherein

R^2 , R^3 , and Z are as defined in embodiment A1.

Embodiment A27. A compound according to any one of the embodiments A1 to A26, wherein

R^2 and R^3 independently of each other are hydrogen, C_{1-6} -alkyl, aryl- C_{1-6} -alkylene-, heteroaryl- C_{1-6} -alkylene-, C_{1-6} -alkyl-aryl-, C_{1-6} -alkyl-heteroarylene-, heteroaryl, or aryl.

Embodiment A28. A compound according to embodiment A27, wherein

R^2 and R^3 independently of each other, are hydrogen, C_{1-6} -alkyl, aryl- C_{1-6} -alkylene- or aryl.

Embodiment A29. A compound according to embodiment A28, wherein R^2 is hydrogen or C_{1-6} -alkyl.

- 5 Embodiment A30. A compound according to embodiment A29, wherein R^2 is hydrogen.

Embodiment A31. A compound according to any one of the embodiments A1 to A30, wherein R^3 is hydrogen or C_{1-6} -alkyl.

- 10 Embodiment A32. A compound according to embodiment A31, wherein R^3 is hydrogen.


- Embodiment A33. A compound according to any one of the embodiments A1 to A22, wherein R^2 and R^3 , when attached to the same nitrogen atom, together with said nitrogen atom may form a 3 to 8 membered heterocyclic ring optionally containing one or two further heteroatoms selected from nitrogen, oxygen and sulfur, and optionally
15 containing one or two double bonds.

Embodiment A34. A compound according to any one of the embodiments A1 to A33, wherein Z is a direct bond, -O-, -NH-, -NHCH₂-, -S-, -SO₂-, -C(O)NH-, -NHC(O)-, -N(H)CON(H)-, -N(CH₃)CONH-, -N(H)SO₂-, -SO₂N(H)-, -C(O)-O-, -N(H)SO₂N(H)-, or -O-C(O)-.

- 20 Embodiment A35. A compound according to embodiment A34, wherein Z is a direct bond, -O-, -S-, -SO₂-, -C(O)NH-, -NHC(O)-, -N(H)SO₂-, -C(O)-O-, -N(H)SO₂N(H)-, or -O-C(O)-.

Embodiment A36. A compound according to embodiment A35, wherein Z is a direct bond, -NHC(O)-, or -NHS(O)₂-.

- 25 Embodiment A37. A compound according to embodiment A2, wherein A¹ is



, wherein

R^{23} , R^{24} , R^{25} , and R^{26} , independently of each other, are hydrogen or as defined in embodiment A1.

Embodiment A38. A compound according to embodiment A37, wherein

- 5 R^{23} , R^{24} , R^{25} , and R^{26} independently of each other are selected from the group consisting of
 halogen, $-\text{C}(\text{O})\text{OR}^2$, $-\text{CN}$, $-\text{CF}_3$, $-\text{OCF}_3$, $-\text{NO}_2$, $-\text{OR}^2$, $-\text{NR}^2\text{R}^3$, C_{1-6} -alkyl-Z-,
 cycloalkyl-Z-, heterocyclyl-Z-, aryl-Z-, or heteroaryl-Z-, $\text{N}(\text{R}^4\text{R}^5)\text{-C}_{1-6}$ -alkylene-Z-,
 $\text{R}^6\text{-W}^1\text{-Z-}$, $\text{R}^6\text{-W}^1\text{-N}(\text{R}^4)\text{-Z-}$, $\text{R}^6\text{-N}(\text{R}^4)\text{-Z-}$, and $\text{R}^6\text{-W}^1\text{-C}_{1-6}$ -alkylene-Z-, wherein
 10 R^2 , R^3 , R^4 , R^5 , R^6 , Z , and W^1 are as defined in embodiment A1.

Embodiment A39. A compound according to embodiment A37 or A38, wherein

R^4 , R^5 , and R^6 independently of each other are selected from the group consisting of
 hydrogen, aryl, C_{1-6} -alkyl, heteroaryl- C_{1-6} -alkylene-, and aryl- C_{1-6} -alkylene-.

Embodiment A40. A compound according to embodiment A39, wherein


- 15 R^4 , R^5 , and R^6 independently of each other are hydrogen or C_{1-6} -alkyl.

Embodiment A41. A compound according to embodiment A40, wherein

R^4 , R^5 , and R^6 are hydrogen.

Embodiment A42. A compound according to embodiment A37 or A38, wherein

- 20 R^4 and R^5 is taken together to form a ring having the formula $-(\text{CH}_2)_j\text{-Q-(CH}_2)_k\text{-}$
 bonded to the nitrogen atom to which R^4 and R^5 are attached, wherein
 j and k independently of each other is 1, 2, 3, or 4;
 Q is a direct bond, $-\text{CH}_2\text{-}$, $-\text{O-}$, $-\text{S-}$, $-\text{S}(\text{O}_2)\text{-}$, $-\text{C}(\text{O})\text{-}$, $-\text{C}(\text{O})\text{NH-}$, $-\text{NHC}(\text{O})\text{-}$,
 $-\text{NHC}(\text{O})\text{NH-}$, $-\text{NH}\text{SO}_2\text{-}$, $-\text{SO}_2\text{NH-}$, $-\text{C}(\text{O})\text{-O-}$, $-\text{O-C}(\text{O})\text{-}$, $-\text{NH}\text{SO}_2\text{NH-}$,




wherein

R^9 and R^{10} independently of each other are selected from the group consisting of hydrogen, aryl, C_{1-6} -alkyl, and arylalkyl-.

5 Embodiment A43. A compound according to embodiment A42, wherein

Q is a direct bond, $-\text{CH}_2-$, $-\text{O}-$, $-\text{S}-$, $-\text{S}(\text{O}_2)-$, $-\text{C}(\text{O})-$, $-\text{C}(\text{O})\text{NH}-$, $-\text{NHC}(\text{O})-$, $-\text{NHC}(\text{O})\text{NH}-$, $-\text{NHSO}_2-$, $-\text{SO}_2\text{NH}-$, $-\text{C}(\text{O})-\text{O}-$, $-\text{O}-\text{C}(\text{O})-$, $-\text{NHSO}_2\text{NH}-$,




wherein

10 R^9 and R^{10} independently of each other are hydrogen or alkyl.

Embodiment A44. A compound according to embodiment A43, wherein

Q is a direct bond, $-\text{CH}_2-$, $-\text{O}-$, $-\text{S}-$, $-\text{S}(\text{O}_2)-$, $-\text{C}(\text{O})-$, $-\text{C}(\text{O})\text{NH}-$, $-\text{NHC}(\text{O})-$, $-\text{NHC}(\text{O})\text{NH}-$, $-\text{NHSO}_2-$, $-\text{SO}_2\text{NH}-$, $-\text{C}(\text{O})-\text{O}-$, $-\text{O}-\text{C}(\text{O})-$, $-\text{NHSO}_2\text{NH}-$,



15 Embodiment A45. A compound according to embodiment A44, wherein

Q is a direct bond.

Embodiment A46. A compound according to any one of the embodiments A37 to A45, wherein

W^1 is a direct bond, -O-, -C(O)-, -NH-, -S-, -SO₂-, -C(O)NH-, -NHC(O)-, -N(H)CON(H)-, -N(H)SO₂-, -SO₂N(H)-, -C(O)-O-, -N(H)SO₂N(H)-, or -O-C(O)-.

5

Embodiment A47. A compound according to embodiment A46 wherein

W^1 is a direct bond, -O-, -C(O)-, -SO₂-, -C(O)NH-, -NHC(O)-, -N(H)SO₂-, -C(O)-O-, or -O-C(O)-.

Embodiment A48. A compound according to embodiment A47, wherein

10 W^1 is a direct bond, or -C(O)-O-.

Embodiment A49. A compound according to embodiment A48, wherein

W^1 is a direct bond.

Embodiment A50. A compound according to embodiment A37, wherein

15 at least one of R^{23} , R^{24} , R^{25} , and R^{26} is halogen, -C(O)OR², -CN, -CF₃, -OCF₃, -NO₂, -OR², -NR²R³, C₁₋₆-alkyl-Z-, cycloalkyl-Z-, heterocyclyl-Z-, aryl-Z-, or heteroaryl-Z-, wherein

R^2 , R^3 , and Z are as defined in embodiment A37.

Embodiment A51. A compound according to embodiment A50, wherein

20 at least one of R^{23} , R^{24} , R^{25} , and R^{26} is halogen, -C(O)OR², -CN, -CF₃, -OCF₃, -NO₂, -OR², -NR²R³, C₁₋₆-alkyl-Z-, C₃₋₁₀-cycloalkyl-Z-, C₃₋₁₀-heterocyclyl-Z-, C₃₋₁₀-aryl-Z-, or C₃₋₁₀-heteroaryl-Z-, wherein

R^2 , R^3 , and Z are as defined in embodiment A37.

Embodiment A52. A compound according to embodiment A51, wherein

25 at least one of R^{23} , R^{24} , R^{25} , and R^{26} is halogen, -C(O)OR², -CN, -NO₂, -OR², -NR²R³, C₁₋₆-alkyl-Z-, C₃₋₁₀-cycloalkyl-Z-, C₃₋₁₀-heterocyclyl-Z-, C₃₋₁₀-aryl-Z-, or C₃₋₁₀-heteroaryl-Z-, wherein

R^2 , R^3 , and Z are as defined in embodiment A37.

Embodiment A53. A compound according to any one of the embodiments A37 to A52, wherein

Z is a direct bond, -O-, -NH-, -NHCH₂-, -S-, -SO₂-, -C(O)NH-, -NHC(O)-, -N(H)CON(H)-, -N(CH₃)CON(H)-, -N(H)SO₂-, -SO₂N(H)-, -C(O)-O-, -N(H)SO₂N(H)-, or -O-C(O)-.

Embodiment A54. A compound according to embodiment A53, wherein

- 5 Z is a direct bond, -O-, -S-, -SO₂-, -C(O)NH-, -NHC(O)-, -N(H)SO₂-, -C(O)-O-, -N(H)SO₂N(H)-, or -O-C(O)-.

Embodiment A55. A compound according to embodiment A54, wherein

Z is a direct bond, -NHC(O)-, or -NHS(O)₂.

Embodiment A56. A compound according to embodiment A52, wherein

- 10 at least one of R²³, R²⁴, R²⁵, and R²⁶ is halogen, -C(O)OR², -CN, -NO₂, -OR², -NR²R³, or C₁₋₆-alkyl, wherein
R², and R³ are as defined in embodiment A37.

Embodiment A57. A compound according to any one of the embodiments A37 to A56, wherein

- 15 R² and R³ independently of each other are hydrogen, C₁₋₆-alkyl, aryl-C₁₋₆-alkylene-, heteroaryl-C₁₋₆-alkylene-, C₁₋₆-alkyl-arylene-, C₁₋₆-alkyl-heteroarylene-, heteroaryl, or aryl.

Embodiment A58. A compound according to embodiment A57, wherein

- 20 R² and R³ independently of each other, are hydrogen, C₁₋₆-alkyl, aryl-C₁₋₆-alkylene- or aryl.

Embodiment A59. A compound according to embodiment A58, wherein

R² is hydrogen or C₁₋₆-alkyl.

Embodiment A60. A compound according to embodiment A59, wherein

R² is hydrogen.

- 25 Embodiment A61. A compound according to any one of the embodiments A37 to A60, wherein

R³ is hydrogen or C₁₋₆-alkyl.

Embodiment A62. A compound according to embodiment A61, wherein
R³ is hydrogen.

Embodiment A63. A compound according to any one of the embodiments A37 to A50,
wherein

- 5 R² and R³, when attached to the same nitrogen atom, together with said nitrogen atom may form a 3 to 8 membered heterocyclic ring optionally containing one or two further heteroatoms selected from nitrogen, oxygen and sulfur, and optionally containing one or two double bonds.

- Embodiment A64. A compound according to any one of the embodiments A37 to A55,
10 wherein
 at least one of R²³, R²⁴, R²⁵, and R²⁶ are hydrogen.

Embodiment A65. A compound according to embodiment A64, wherein
 at least two of R²³, R²⁴, R²⁵, and R²⁶ are hydrogen.

- Embodiment A66. A compound according to embodiment A65, wherein
15 R²³ and R²⁶ are hydrogen.

Embodiment A67. A compound according to embodiment A65 or embodiment A66, wherein
 at least three of R²³, R²⁴, R²⁵, and R²⁶ are hydrogen.

Embodiment A68. A compound according to any one of the embodiments A37 to A67,
wherein R²⁴ or R²⁵ is halogen.

- 20 69. A compound according to embodiment A68, wherein R²⁴ or R²⁵ is fluoro.

Embodiment A70. A compound according to any one of the embodiments A37 to A67,
wherein R²⁴ or R²⁵ is C₁₋₆-alkyl.

Embodiment A71. A compound according to embodiment A68, wherein R²⁴ or R²⁵ is methyl.

- Embodiment A72. A compound according to any one of the embodiments A37 to A67,
25 wherein
 R²⁴ is hydrogen.

Embodiment A73. A compound according to any one of the embodiments A37 to A72, wherein

R^{25} is hydrogen.

Embodiment A74. A compound according to embodiment A37, wherein

5 R^{23} , R^{24} , R^{25} , and R^{26} are hydrogen.

Embodiment A75. A compound according to any one of the embodiments A1 to A74, wherein

L^1 is a bond, -D-alkylene-E-, -O-, -S-, -S(O)-, -S(O)₂-, -C(O)-, -N(R¹¹)-, or -C(=N-OR¹²), wherein

D, E, R¹¹ and R¹² are as defined in embodiment A1.

10 Embodiment A76. A compound according to embodiment A75, wherein

L^1 is a bond, -D-alkylene-E-, -O-, -C(O)-, -N(R¹¹)-, or -C(=N-OR¹²)-, wherein

D, E, R¹¹ and R¹² are as defined in embodiment A1.

Embodiment A77. A compound according to embodiment A75, wherein

L^1 is -O-.

15 Embodiment A78. A compound according to embodiment A75, wherein

L^1 is -S-.

Embodiment A79. A compound according to embodiment A75, wherein

L^1 is a bond.

80. A compound according to embodiment A75, wherein

20 L^1 is -C(O)-.

Embodiment A81. A compound according to any one of the embodiments A1 to A76, wherein

D is a direct bond or -O-;

E is a direct bond or -O-; and

R¹¹ and R¹² are as defined in embodiment A1.

25 Embodiment A82. A compound according to embodiment A81, wherein

D is a direct bond.

Embodiment A83. A compound according to embodiment A81, wherein
D is -O-.

Embodiment A84. A compound according to any one of the embodiments A81 to A83,
wherein

5 E is a direct bond.

Embodiment A85. A compound according to any one of the embodiments A81 to A83,
wherein

E is -O-.

Embodiment A86. A compound according to any one of the embodiments A1 to A85, wherein

10 R¹¹ is selected from hydrogen, C₁₋₆-alkyl, aryl, carbamoyl, aryl-C₁₋₆-alkylene-,
C₁₋₆-alkyl-NH-C(O)-, aryl-C₁₋₆-alkylene-NH-C(O)-, C₁₋₆-alkyl-SO₂-,
aryl-C₁₋₆-alkylene-SO₂-, aryl-SO₂-, SO₂-, C₁₋₆-alkyl-C(O)-, aryl-C₁₋₆-alkylene-C(O)-,
N(R¹³)(R¹⁴)-C₁₋₆-alkylene-Y-, and R¹⁵-W²-C₁₋₆-alkylene-Y-, wherein

15 Y and W² independently of each other are a direct bond, -CH₂-, -SO₂-,
-N(H)CO-, -N(H)SO₂-, or -O-C(O)-;

R¹³ and R¹⁴ independently of each other are selected from hydrogen, aryl,
C₁₋₆-alkyl, or aryl-C₁₋₆-alkylene-;


or

20 R¹³ and R¹⁴ may be taken together to form a ring having the formula
-(CH₂)_o-X-(CH₂)_p- bonded to the nitrogen atom to which R¹³ and R¹⁴ are
attached, wherein

o and p are independently of each other are 1, 2, 3, or 4; and

25 X is a direct bond, -CH₂-, -O-, -S-, -S(O₂)-, -C(O)-, -CON(H)-,
-NHC(O)-, -NHCON(H)-, -NHSO₂-, -SO₂N(H)-, -C(O)-O-, -O-C(O)-,
-NHSO₂NH-,

90



wherein

R^{16} and R^{17} are selected from hydrogen, aryl, heteroaryl,
 C_{1-6} -alkyl, C_{1-6} -alkoxy, aryl- C_{1-6} -alkylene-,
 heteroaryl- C_{1-6} -alkylene-, C_{1-6} -alkyl-arylene-,
 C_{1-6} -alkyl-heteroarylene-, C_{1-6} -alkoxy-arylene-,
 C_{1-6} -alkoxy-heteroarylene-, heteroarylaryl- C_{1-6} -alkoxy-, or
 aryl- C_{1-6} -alkoxy-; and

R^{15} is selected from the group consisting of aryl, heteroaryl, cycloalkyl, heterocyclyl,
 C_{1-6} -alkyl, or aryl- C_{1-6} -alkylene-.

Embodiment A87. A compound according to embodiment A86, wherein

R^{11} is selected from hydrogen, C_{1-6} -alkyl, aryl, carbamoyl, aryl- C_{1-6} -alkylene-,
 C_{1-6} -alkyl-NH-C(O)-, aryl- C_{1-6} -alkylene-NH-C(O)-, C_{1-6} -alkyl-SO₂-,
 aryl- C_{1-6} -alkylene-SO₂-, aryl-SO₂-, SO₂-, C_{1-6} -alkyl-C(O)-, aryl- C_{1-6} -alkylene-C(O)-,
 $\text{N(R}^{13})(\text{R}^{14})\text{-C}_{1-6}\text{-alkylene-Y-}$, and $\text{R}^{15}\text{-W}^2\text{-C}_{1-6}\text{-alkylene-Y-}$, wherein

Y and W^2 independently of each other are a direct bond, -CH₂-, -SO₂-,
 -N(H)CO-, -N(H)SO₂-, or -O-C(O)-;




R^{13} and R^{14} independently of each other are selected from hydrogen, aryl,
 C_{1-6} -alkyl, or aryl- C_{1-6} -alkylene-;

or

R^{13} and R^{14} may be taken together to form a ring having the formula
 $\text{-(CH}_2\text{)}_o\text{-X-(CH}_2\text{)}_p\text{-}$ bonded to the nitrogen atom to which R^{13} and R^{14} are
 attached, wherein

o and p are independently of each other are 1, 2, 3, or 4; and

X is a direct bond, $-\text{CH}_2-$, $-\text{O}-$, $-\text{S}-$, $-\text{S}(\text{O}_2)-$, $-\text{C}(\text{O})-$, $-\text{CON}(\text{H})-$, $-\text{NHC}(\text{O})-$, $-\text{NHCON}(\text{H})-$, $-\text{NHSO}_2-$, $-\text{SO}_2\text{N}(\text{H})-$, $-\text{C}(\text{O})-\text{O}-$, $-\text{O}-\text{C}(\text{O})-$, $-\text{NHSO}_2\text{NH}-$,



5

wherein

R^{16} and R^{17} are hydrogen; and

R^{15} is selected from the group consisting of aryl, heteroaryl, cycloalkyl, heterocyclyl, C_{1-6} -alkyl, or aryl- C_{1-6} -alkylene-.

Embodiment A88. A compound according to embodiment A87, wherein

10

R^{11} is selected from hydrogen, C_{1-6} -alkyl, aryl, carbamoyl, aryl- C_{1-6} -alkylene-, C_{1-6} -alkyl-NH-C(O)-, aryl- C_{1-6} -alkylene-NH-C(O)-, C_{1-6} -alkyl-SO₂-, aryl- C_{1-6} -alkylene-SO₂-, aryl-SO₂-, SO₂-, C_{1-6} -alkyl-C(O)-, aryl- C_{1-6} -alkylene-C(O)-, $\text{N}(\text{R}^{13})(\text{R}^{14})\text{-C}_{1-6}\text{-alkylene-Y-}$, and $\text{R}^{15}\text{-W}^2\text{-C}_{1-6}\text{-alkylene-Y-}$, wherein

15

Y and W^2 independently of each other are a direct bond, $-\text{CH}_2-$, $-\text{SO}_2-$, $-\text{N}(\text{H})\text{CO}-$, $-\text{N}(\text{H})\text{SO}_2-$, or $-\text{O}-\text{C}(\text{O})-$;

R^{13} and R^{14} independently of each other are selected from hydrogen, aryl, C_{1-6} -alkyl, or aryl- C_{1-6} -alkylene-;

or

20

R^{13} and R^{14} may be taken together to form a ring having the formula $-(\text{CH}_2)_o\text{-X-(CH}_2)_p-$ bonded to the nitrogen atom to which R^{13} and R^{14} are attached, wherein

o and p are independently of each other are 1, 2, 3, or 4;

X is a direct bond; and

R¹⁵ is selected from the group consisting of aryl, heteroaryl, cycloalkyl, heterocyclyl, C₁₋₆-alkyl, or aryl-C₁₋₆-alkylene-.

Embodiment A89. A compound according to embodiment A88, wherein

- 5 R¹¹ is selected from hydrogen, C₁₋₆-alkyl, aryl, carbamoyl, aryl-C₁₋₆-alkylene-, C₁₋₆-alkyl-NH-C(O)-, aryl-C₁₋₆-alkylene-NH-C(O)-, C₁₋₆-alkyl-SO₂-, aryl-C₁₋₆-alkylene-SO₂-, aryl-SO₂-, SO₂-, C₁₋₆-alkyl-C(O)-, aryl-C₁₋₆-alkylene-C(O)-, N(R¹³)(R¹⁴)-C₁₋₆-alkylene-Y-, and R¹⁵-W²-C₁₋₆-alkylene-Y-, wherein
- Y and W² independently of each other are a direct bond, -CH₂-, -SO₂-, -N(H)CO-, -N(H)SO₂-, or -O-C(O)-;
- 10 R¹³ and R¹⁴ independently of each other are selected from hydrogen, aryl, C₁₋₆-alkyl, or aryl-C₁₋₆-alkylene-; and
- R¹⁵ is selected from the group consisting of aryl, heteroaryl, cycloalkyl, heterocyclyl, C₁₋₆-alkyl, or aryl-C₁₋₆-alkylene-.

Embodiment A90. A compound according to embodiment A89, wherein

- 15 R¹¹ is selected from hydrogen, C₁₋₆-alkyl, aryl, carbamoyl, aryl-C₁₋₆-alkylene-, C₁₋₆-alkyl-NH-C(O)-, aryl-C₁₋₆-alkylene-NH-C(O)-, C₁₋₆-alkyl-SO₂-, aryl-C₁₋₆-alkylene-SO₂-, aryl-SO₂-, SO₂-, C₁₋₆-alkyl-C(O)-, aryl-C₁₋₆-alkylene-C(O)-, N(R¹³)(R¹⁴)-C₁₋₆-alkylene-Y-, and R¹⁵-W²-C₁₋₆-alkylene-Y-, wherein
- Y and W² independently of each other are a direct bond, -CH₂-, -SO₂-, -N(H)CO-, -N(H)SO₂-, or -O-C(O)-;
- 20 R¹³ and R¹⁴ are hydrogen; and
- R¹⁵ is selected from the group consisting of aryl, heteroaryl, cycloalkyl, heterocyclyl, C₁₋₆-alkyl, or aryl-C₁₋₆-alkylene-.

Embodiment A91. A compound according to embodiment A90, wherein

- 25 R¹¹ is selected from hydrogen, C₁₋₆-alkyl, aryl, carbamoyl, aryl-C₁₋₆-alkylene-, C₁₋₆-alkyl-NH-C(O)-, aryl-C₁₋₆-alkylene-NH-C(O)-, C₁₋₆-alkyl-SO₂-, aryl-C₁₋₆-alkylene-SO₂-, aryl-SO₂-, SO₂-, C₁₋₆-alkyl-C(O)-, aryl-C₁₋₆-alkylene-C(O)-, N(R¹³)(R¹⁴)-C₁₋₆-alkylene-Y-, and R¹⁵-W²-C₁₋₆-alkylene-Y-, wherein
- Y is a direct bond;
- 30 W² is a direct bond, -CH₂-, -SO₂-, -N(H)CO-, -N(H)SO₂-, or -O-C(O)-;
- R¹³ and R¹⁴ are hydrogen; and

R¹⁵ is selected from the group consisting of aryl, heteroaryl, cycloalkyl, heterocyclyl, C₁₋₆-alkyl, or aryl-C₁₋₆-alkylene-.

Embodiment A92. A compound according to embodiment A90, wherein

- 5 R¹¹ is selected from hydrogen, C₁₋₆-alkyl, aryl, carbamoyl, aryl-C₁₋₆-alkylene-, C₁₋₆-alkyl-NH-C(O)-, aryl-C₁₋₆-alkylene-NH-C(O)-, C₁₋₆-alkyl-SO₂-, aryl-C₁₋₆-alkylene-SO₂-, aryl-SO₂-, SO₂-, C₁₋₆-alkyl-C(O)-, aryl-C₁₋₆-alkylene-C(O)-, N(R¹³)(R¹⁴)-C₁₋₆-alkylene-Y-, and R¹⁵-W²-C₁₋₆-alkylene-Y-, wherein
- Y is a direct bond, -CH₂-, -SO₂-, -N(H)CO-, -N(H)SO₂-, or -O-C(O)-;
- W² is a direct bond;
- 10 R¹³ and R¹⁴ are hydrogen; and
- R¹⁵ is selected from the group consisting of aryl, heteroaryl, cycloalkyl, heterocyclyl, C₁₋₆-alkyl, or aryl-C₁₋₆-alkylene-.

Embodiment A93. A compound according to any one of the embodiments A90 to A92, wherein

- 15 R¹¹ is selected from hydrogen, C₁₋₆-alkyl, aryl, carbamoyl, aryl-C₁₋₆-alkylene-, C₁₋₆-alkyl-NH-C(O)-, aryl-C₁₋₆-alkylene-NH-C(O)-, C₁₋₆-alkyl-SO₂-, aryl-C₁₋₆-alkylene-SO₂-, aryl-SO₂-, SO₂-, C₁₋₆-alkyl-C(O)-, aryl-C₁₋₆-alkylene-C(O)-, NH₂-C₁₋₆-alkylene-, and R¹⁵-C₁₋₆-alkylene-, wherein
- R¹⁵ is selected from the group consisting of aryl, heteroaryl, cycloalkyl,
- 20 heterocyclyl, C₁₋₆-alkyl, or aryl-C₁₋₆-alkylene-.

Embodiment A94. A compound according to embodiment A93, wherein

- R¹¹ is selected from hydrogen, C₁₋₆-alkyl, aryl, carbamoyl, aryl-C₁₋₆-alkylene-, C₁₋₆-alkyl-NH-C(O)-, aryl-C₁₋₆-alkylene-NH-C(O)-, C₁₋₆-alkyl-SO₂-, aryl-C₁₋₆-alkylene-SO₂-, aryl-SO₂-, SO₂-, C₁₋₆-alkyl-C(O)-, aryl-C₁₋₆-alkylene-C(O)-,
- 25 and NH₂-C₁₋₆-alkylene-.

Embodiment A95. A compound according to embodiment A94, wherein

R¹¹ is hydrogen or C₁₋₆-alkyl.

Embodiment A96. A compound according to embodiment A95, wherein

R¹¹ is hydrogen.

Embodiment A97. A compound according to any one of the embodiments A1 to A96, wherein R^{12} is hydrogen or C_{1-6} -alkyl.

Embodiment A98. A compound according to embodiment A97, wherein R^{12} is hydrogen.

5 Embodiment A99. A compound according to any one of the embodiments A1 to A98, wherein G^1 is C_{1-6} -alkyl, C_{2-6} -alkenyl, C_{2-6} -alkynyl, cycloalkyl or heterocyclyl, optionally substituted with one or more substituents selected from the group consisting of -CN, -CF₃, -OCF₃, -OR¹⁸, -NR¹⁸R¹⁹, C_{3-10} -cycloalkyl and C_{1-6} -alkyl, wherein R^{18} and R^{19} are as defined in embodiment A1.

10 Embodiment A100. A compound according to embodiment A99, wherein G^1 is C_{1-6} -alkyl, C_{2-6} -alkenyl, C_{2-6} -alkynyl, cycloalkyl or heterocyclyl, optionally substituted with one or more substituents selected from the group consisting of -CN, -CF₃, -OCF₃, -OR¹⁸, -NR¹⁸R¹⁹ and C_{1-6} -alkyl, wherein R^{18} and R^{19} are as defined in embodiment A1.

15 Embodiment A101. A compound according to embodiment A99, wherein G^1 is C_{1-6} -alkyl, C_{2-6} -alkenyl, C_{2-6} -alkynyl, C_{3-10} -cycloalkyl or C_{3-10} -heterocyclyl, optionally substituted with one or more substituents selected from the group consisting of -CN, -CF₃, -OCF₃, -OR¹⁸, -NR¹⁸R¹⁹, C_{3-10} -cycloalkyl and C_{1-6} -alkyl, wherein
20 R^{18} and R^{19} are as defined in embodiment A1.

Embodiment A102. A compound according to embodiment A100 or embodiment A101, wherein

G^1 is C_{1-6} -alkyl, C_{2-6} -alkenyl, C_{2-6} -alkynyl, C_{3-10} -cycloalkyl or C_{3-10} -heterocyclyl, optionally substituted with one or more substituents selected from the group
25 consisting of -CN, -CF₃, -OCF₃, -OR¹⁸, -NR¹⁸R¹⁹ and C_{1-6} -alkyl, wherein R^{18} and R^{19} are as defined in embodiment A1.

Embodiment A103. A compound according to any one of the embodiments A99 to A102, wherein

G^1 is selected from the group consisting of methyl, ethyl, propyl, butyl, isopropyl,
30 isobutyl, sec-butyl, tert-butyl, 3-pentyl, 2-pentyl, 3-methyl-butyl, 2-propenyl,

cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, cycloheptyl, oxetanyl, tetrahydrofuranyl, tetrahydropyranyl, azetidyl, pyrrolidyl, piperidyl, hexahydroazepinyl, thiolanyl, tetrahydrothiopyranyl, thiepanyl, 1,4-oxathianyl, 1,3-dioxolanyl, 1,2-dithiolanyl, 1,3-dithiolanyl, hexahydro-pyridaziny, imidazolidyl, 1,3-dioxanyl, morpholinyl, 1,3-dithianyl, 1,4-dioxanyl, 1,4-dithianyl, or thiomorpholinyl.

Embodiment A104. A compound according to embodiment A103 wherein

G¹ is selected from the group consisting of methyl, ethyl, propyl, butyl, isopropyl, isobutyl, *sec*-butyl, *tert*-butyl, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, cycloheptyl, tetrahydrofuranyl, tetrahydropyranyl, pyrrolidyl, piperidyl, hexahydroazepinyl, thiolanyl, tetrahydrothiopyranyl, or thiepanyl.

Embodiment A105. A compound according to embodiment A104 wherein

G¹ is selected from the group consisting of methyl, ethyl, propyl, butyl, isopropyl, isobutyl, cyclopentyl, cyclohexyl, tetrahydrofuranyl, tetrahydropyranyl, piperidyl, or hexahydroazepinyl.

Embodiment A106. A compound according to embodiment A105 wherein

G¹ is selected from the group consisting of isobutyl, cyclopentyl, and piperidyl.

Embodiment A107. A compound according to embodiment A105 wherein

G¹ is isobutyl.

Embodiment A108. A compound according to embodiment A105 wherein

G¹ is cyclopentyl.

Embodiment A109. A compound according to embodiment A105 wherein

G¹ is piperidyl.

Embodiment A110. A compound according to any one of the embodiments A99 to A102, wherein

R¹⁸ and R¹⁹, independently of each other, are hydrogen, C₁₋₆-alkyl, heteroaryl-C₁₋₆-alkylene-, aryl-C₁₋₆-alkylene-, C₁₋₆-alkyl-arylene-, C₁₋₆-alkyl-heteroarylene-, heteroaryl, or aryl.

Embodiment A111. A compound according to embodiment A110, wherein

R¹⁸ and R¹⁹, independently of each other, are hydrogen, C₁₋₆-alkyl, C₃₋₁₀-heteroaryl-C₁₋₆-alkylene-, C₃₋₁₀-aryl-C₁₋₆-alkylene-, C₁₋₆-alkyl-C₃₋₁₀-arylene-, C₁₋₆-alkyl-C₃₋₁₀-heteroarylene-, C₃₋₁₀-heteroaryl, or C₃₋₁₀-aryl.

Embodiment A112. A compound according to embodiment A111, wherein

R^{18} and R^{19} , independently of each other, are hydrogen or C_{1-8} -alkyl.

Embodiment A113. A compound according to embodiment A112, wherein

R^{18} is hydrogen.

Embodiment A114. A compound according to embodiment A112 or A113, wherein

5 R^{19} is hydrogen.

Embodiment A115. A compound according to embodiment A100 or embodiment A102, wherein

10 R^{18} and R^{19} , when attached to the same nitrogen atom, together with the said nitrogen atom forms a 3 to 8 membered heterocyclic ring optionally containing one or two further heteroatoms selected from nitrogen, oxygen and sulfur, and optionally containing one or two double bonds.

Embodiment A116. A compound according to any one of the embodiments A1 to A98, wherein

15 G^1 is alkyl or cycloalkyl, optionally substituted with one or more substituents selected from the group consisting of -CN, -CF₃, -OCF₃, -OR¹⁸, -NR¹⁸R¹⁹ and C_{1-8} -alkyl, or G^1 is aryl optionally substituted with one or more substituents R^{40} , R^{41} , and R^{42} , wherein R^{18} , R^{19} , R^{40} , R^{41} , and R^{42} are as defined in embodiment A1.

Embodiment A117. A compound according to embodiment A116, wherein

20 G^1 is C_{1-8} -alkyl or C_{3-10} -cycloalkyl, optionally substituted with one or more substituents selected from the group consisting of -CN, -CF₃, -OCF₃, -OR¹⁸, -NR¹⁸R¹⁹ and C_{1-8} -alkyl, or G^1 is C_{3-10} -aryl optionally substituted with one or more substituents R^{40} , R^{41} , and R^{42} , wherein R^{18} , R^{19} , R^{40} , R^{41} , and R^{42} are as defined in embodiment A1.

25 Embodiment A118. A compound according to embodiment A117, wherein

G^1 is C_{1-8} -alkyl or C_{3-10} -cycloalkyl, optionally substituted with one or more substituents selected from the group consisting of -CN, -CF₃, -OCF₃, -OR¹⁸, -NR¹⁸R¹⁹ and C_{1-8} -alkyl, or G^1 is phenyl optionally substituted with one or more substituents R^{40} , R^{41} , and R^{42} , wherein R^{18} , R^{19} , R^{40} , R^{41} , and R^{42} are as defined in embodiment A1.

30

Embodiment A119. A compound according to embodiment A116, A117, or A118, wherein R^{18} and R^{19} , independently of each other, are hydrogen, C_{1-6} -alkyl, heteroaryl- C_{1-6} -alkylene-, aryl- C_{1-6} -alkylene-, C_{1-6} -alkyl-arylene-, C_{1-6} -alkyl-heteroarylene-, heteroaryl, or aryl.

- 5 Embodiment A120. A compound according to embodiment A119, wherein R^{18} and R^{19} , independently of each other, are hydrogen, C_{1-6} -alkyl, C_{3-10} -heteroaryl- C_{1-6} -alkylene-, C_{3-10} -aryl- C_{1-6} -alkylene-, C_{1-6} -alkyl- C_{3-10} -arylene-, C_{1-6} -alkyl- C_{3-10} -heteroarylene-, C_{3-10} -heteroaryl, or C_{3-10} -aryl.

- Embodiment A121. A compound according to embodiment A120, wherein
10 R^{18} and R^{19} , independently of each other, are hydrogen or C_{1-6} -alkyl.

Embodiment A122. A compound according to embodiment A121, wherein R^{18} is hydrogen.

Embodiment A123. A compound according to embodiment A121 or A122, wherein R^{19} is hydrogen.

- 15 Embodiment A124. A compound according to embodiment A116 or embodiment A117, wherein R^{18} and R^{19} , when attached to the same nitrogen atom, together with the said nitrogen atom forms a 3 to 8 membered heterocyclic ring optionally containing one or two further heteroatoms selected from nitrogen, oxygen and sulfur, and optionally
20 containing one or two double bonds.

Embodiment A125. A compound according to any one of the embodiments A116 to A124, wherein

- R^{40} , R^{41} , and R^{42} independently of each other are
halogen, -CN, -NO₂, C_{1-6} -alkyl, -CHF₂, -CF₃, -OR⁵⁴, -NR⁵⁴R⁵⁵, -SR⁵⁴, -NR⁵⁴S(O)₂R⁵⁵,
25 -S(O)₂NR⁵⁴R⁵⁵, -S(O)NR⁵⁴R⁵⁵, -S(O)R⁵⁴, -S(O)₂R⁵⁴, -C(O)NR⁵⁴R⁵⁵, -OC(O)NR⁵⁴R⁵⁵,
-NR⁵⁴C(O)R⁵⁵, -CH₂C(O)NR⁵⁴R⁵⁵, -CH₂C(O)OR⁵⁴, -OCH₂C(O)NR⁵⁴R⁵⁵, -CH₂OR⁵⁴,
-CH₂NR⁵⁴R⁵⁵, and -C(O)OR⁵⁴; or

C₂₋₈-alkenyl and C₂₋₈-alkynyl, which may optionally be substituted with one or more substituents selected from -CN, -CF₃, -OCF₃, -OR⁵⁴, -NR⁵⁴R⁵⁵ and C₁₋₆-alkyl, wherein

- 5 R⁵⁴ and R⁵⁵ independently of each other are hydrogen, C₁₋₆-alkyl, C₁₋₆-alkyl-arylene-, C₁₋₆-alkyl-heteroarylene-, aryl-C₁₋₆-alkylene-, heteroaryl-C₁₋₆-alkylene-, heteroaryl, or aryl;
- or
- 10 R⁵⁴ and R⁵⁵, when attached to the same nitrogen atom, together with the said nitrogen atom may form a 3 to 8 membered heterocyclic ring optionally containing one or two further heteroatoms selected from nitrogen, oxygen and sulfur, and optionally containing one or two double bonds.

Embodiment A126. A compound according to embodiment A125, wherein

- 15 R⁴⁰, R⁴¹, and R⁴² independently of each other are halogen, -CN, C₁₋₆-alkyl, -CF₃, -OR⁵⁴, -NR⁵⁴R⁵⁵, -SR⁵⁴, -NR⁵⁴S(O)₂R⁵⁵, -S(O)R⁵⁴, -S(O)₂R⁵⁴, -CH₂C(O)OR⁵⁴, -CH₂OR⁵⁴, -CH₂NR⁵⁴R⁵⁵, and -C(O)OR⁵⁴;
- wherein

- 20 R⁵⁴ and R⁵⁵ independently of each other are hydrogen, C₁₋₆-alkyl, C₁₋₆-alkyl-arylene-, C₁₋₆-alkyl-heteroarylene-, aryl-C₁₋₆-alkylene-, heteroaryl-C₁₋₆-alkylene-, heteroaryl, or aryl;
- or
- R⁵⁴ and R⁵⁵, when attached to the same nitrogen atom, together with the said nitrogen atom may form a 3 to 8 membered heterocyclic ring optionally containing one or two further heteroatoms selected from nitrogen, oxygen and sulfur, and optionally containing one or two double bonds.

25 Embodiment A127. A compound according to embodiment A126 wherein

- R⁴⁰, R⁴¹, and R⁴² independently of each other are halogen, or -OR⁵⁴
- wherein
- 30 R⁵⁴ is hydrogen, C₁₋₆-alkyl, C₁₋₆-alkyl-arylene-, C₁₋₆-alkyl-heteroarylene-, aryl-C₁₋₆-alkylene-, heteroaryl-C₁₋₆-alkylene-, heteroaryl, or aryl.

Embodiment A128. A compound according to embodiment A126, wherein

R⁵⁴ and R⁵⁵ independently of each other are hydrogen, or C₁₋₆-alkyl, C₁₋₆-alkyl-C₃₋₁₀-arylene-, C₁₋₆-alkyl-C₃₋₁₀-heteroarylene-, C₃₋₁₀-aryl-C₁₋₆-alkylene-, C₃₋₁₀-heteroaryl-C₁₋₆-alkylene-, C₃₋₁₀-heteroaryl, or C₃₋₁₀-aryl.

Embodiment A129. A compound according to embodiment A128, wherein
R⁵⁴ and R⁵⁵ independently of each other are hydrogen or C₁₋₆-alkyl.

Embodiment A130. A compound according to embodiment A129, wherein
R⁵⁴ is hydrogen.

- 5 Embodiment A131. A compound according to embodiment A129 or embodiment A130,
wherein

R⁵⁵ is hydrogen.

Embodiment A132. A compound according to embodiment A127 wherein
R⁵⁴ is methyl.

- 10 Embodiment A133. A compound according to any one of the embodiments A1 to A98,
wherein

G¹ is aryl or heteroaryl, optionally substituted with one or more substituents R⁴⁰, R⁴¹,
and R⁴², wherein

R⁴⁰, R⁴¹, and R⁴² are as defined in embodiment A1.

- 15 Embodiment A134. A compound according to embodiment A133, wherein

G¹ is C₃₋₁₀-aryl or C₃₋₁₀-heteroaryl, optionally substituted with one or more
substituents R⁴⁰, R⁴¹, and R⁴², wherein

R⁴⁰, R⁴¹, and R⁴² are as defined in embodiment A1.

Embodiment A135. A compound according to embodiment A133, wherein

- 20 G¹ is aryl, optionally substituted with one or more substituents R⁴⁰, R⁴¹, and R⁴²,
wherein

R⁴⁰, R⁴¹, and R⁴² are as defined in embodiment A1.

Embodiment A136. A compound according to embodiment A135, wherein

- 25 G¹ is C₃₋₁₀-aryl, optionally substituted with one or more substituents R⁴⁰, R⁴¹, and
R⁴², wherein

R⁴⁰, R⁴¹, and R⁴² are as defined in embodiment A1.

Embodiment A137. A compound according to embodiment A136, wherein

G¹ is phenyl, optionally substituted with one or more substituents R⁴⁰, R⁴¹, and R⁴²,
wherein

R^{40} , R^{41} , and R^{42} are as defined in embodiment A1.

Embodiment A138. A compound according to any one of the embodiments A133 to A137, wherein

- 5 R^{40} , R^{41} , and R^{42} independently of each other are
halogen, -CN, -NO₂, C₁₋₆-alkyl, -CHF₂, -CF₃, -OR⁵⁴, -NR⁵⁴R⁵⁵, -SR⁵⁴, -NR⁵⁴S(O)₂R⁵⁵,
-S(O)₂NR⁵⁴R⁵⁵, -S(O)NR⁵⁴R⁵⁵, -S(O)R⁵⁴, -S(O)₂R⁵⁴, -C(O)NR⁵⁴R⁵⁵, -OC(O)NR⁵⁴R⁵⁵,
-NR⁵⁴C(O)R⁵⁵, -CH₂C(O)NR⁵⁴R⁵⁵, -CH₂C(O)OR⁵⁴, -OCH₂C(O)NR⁵⁴R⁵⁵, -CH₂OR⁵⁴,
-CH₂NR⁵⁴R⁵⁵, and -C(O)OR⁵⁴; or
C₂₋₆-alkenyl and C₂₋₆-alkynyl, which may optionally be substituted with one or more
10 substituents selected from -CN, -CF₃, -OCF₃, -OR⁵⁴, -NR⁵⁴R⁵⁵ and C₁₋₆-alkyl,
wherein
 R^{54} and R^{55} independently of each other are hydrogen, C₁₋₆-alkyl,
C₁₋₆-alkyl-arylene-, C₁₋₆-alkyl-heteroarylene-, aryl-C₁₋₆-alkylene-,
heteroaryl-C₁₋₆-alkylene-, heteroaryl, or aryl;
15 or
 R^{54} and R^{55} , when attached to the same nitrogen atom, together with the
said nitrogen atom may form a 3 to 8 membered heterocyclic ring optionally
containing one or two further heteroatoms selected from nitrogen, oxygen
and sulfur, and optionally containing one or two double bonds.

20

Embodiment A139. A compound according to embodiment A138, wherein

- R^{40} , R^{41} , and R^{42} independently of each other are
halogen, -CN, C₁₋₆-alkyl, -CF₃, -OR⁵⁴, -NR⁵⁴R⁵⁵, -SR⁵⁴, -NR⁵⁴S(O)₂R⁵⁵, -S(O)R⁵⁴,
-S(O)₂R⁵⁴, -CH₂C(O)OR⁵⁴, -CH₂OR⁵⁴, -CH₂NR⁵⁴R⁵⁵, and -C(O)OR⁵⁴;
25 wherein
 R^{54} and R^{55} independently of each other are hydrogen, C₁₋₆-alkyl,
C₁₋₆-alkyl-arylene-, C₁₋₆-alkyl-heteroarylene-, aryl-C₁₋₆-alkylene-,
heteroaryl-C₁₋₆-alkylene-, heteroaryl, or aryl;
or
30 R^{54} and R^{55} , when attached to the same nitrogen atom, together with the
said nitrogen atom may form a 3 to 8 membered heterocyclic ring optionally
containing one or two further heteroatoms selected from nitrogen, oxygen
and sulfur, and optionally containing one or two double bonds.

Embodiment A140. A compound according to embodiment A139 wherein

R^{40} , R^{41} , and R^{42} independently of each other are halogen, or $-OR^{54}$

wherein

5 R^{54} is hydrogen, C_{1-6} -alkyl, C_{1-6} -alkyl-arylene-, C_{1-6} -alkyl-heteroarylene-, aryl- C_{1-6} -alkylene-, heteroaryl- C_{1-6} -alkylene-, heteroaryl, or aryl.

Embodiment A141. A compound according to any one of the embodiments A138 to A139, wherein

10 R^{54} and R^{55} independently of each other are hydrogen, or C_{1-6} -alkyl, C_{1-6} -alkyl- C_{3-10} -arylene-, C_{1-6} -alkyl- C_{3-10} -heteroarylene-, C_{3-10} -aryl- C_{1-6} -alkylene-, C_{3-10} -heteroaryl- C_{1-6} -alkylene-, C_{3-10} -heteroaryl, or C_{3-10} -aryl.

Embodiment A142. A compound according to embodiment A141, wherein

R^{54} and R^{55} independently of each other are hydrogen or C_{1-6} -alkyl.

Embodiment A143. A compound according to embodiment A142, wherein

R^{54} is hydrogen.


15 Embodiment A144. A compound according to embodiment A142 or embodiment A143, wherein

R^{55} is hydrogen.

Embodiment A145. A compound according to embodiment A140 wherein

R^{54} is methyl.

20 Embodiment A146. A compound according to embodiment A135, wherein G^1 is



, wherein

25 R^{56} , R^{57} , R^{58} , R^{59} , and R^{60} , independently of each other, are hydrogen or halogen, $-CN$, $-NO_2$, C_{1-6} -alkyl, $-CHF_2$, $-CF_3$, $-OCF_3$, $-OCHF_2$, $-OCH_2CF_3$, $-OCF_2CHF_2$, $-S(O)_2CF_3$, $-SCF_3$, $-OR^{61}$, $-NR^{61}R^{62}$, $-SR^{61}$, $-NR^{61}S(O)_2R^{62}$, $-S(O)_2NR^{61}R^{62}$, $-S(O)NR^{61}R^{62}$, $-S(O)R^{61}$, $-S(O)_2R^{61}$,

- C(O)NR⁶¹R⁶², -OC(O)NR⁶¹R⁶², -NR⁶¹C(O)R⁶², -CH₂C(O)NR⁶¹R⁶², -CH₂C(O)OR⁶¹, -OCH₂C(O)NR⁶¹R⁶², -CH₂OR⁶¹, -CH₂NR⁶¹R⁶², -OC(O)R⁶¹, -C(O)R⁶¹ and -C(O)OR⁶¹; or C₂₋₆-alkenyl and C₂₋₆-alkynyl, which may optionally be substituted with one or more substituents selected from -CN, -CF₃, -OCF₃, -OR⁶¹, -NR⁶¹R⁶² and C₁₋₆-alkyl;
- 5 C₃₋₁₀-cycloalkyl, C₄₋₈-cycloalkenyl, heterocyclyl, C₃₋₁₀-cycloalkyl-C₁₋₆-alkylene-, C₃₋₁₀-cycloalkyl-C₁₋₆-alkoxy-, C₃₋₁₀-cycloalkyloxy, C₃₋₁₀-cycloalkyl-C₁₋₆-alkylthio-, C₃₋₁₀-cycloalkylthio-, C₃₋₁₀-cycloalkyl-C₂₋₆-alkenylene-, C₃₋₁₀-cycloalkyl-C₂₋₆-alkynylene-, C₄₋₈-cycloalkenyl-C₁₋₆-alkylene-, C₄₋₈-cycloalkenyl-C₂₋₆-alkenylene-, C₄₋₈-cycloalkenyl-C₂₋₆-alkynylene-, heterocyclyl-C₁₋₆-alkylene-, heterocyclyl-C₂₋₆-alkenylene-, heterocyclyl-C₂₋₆-alkynylene-; or
- 10 aryl, aryloxy, aryloxycarbonyl, aroyl, aryl-C₁₋₆-alkoxy-, aryl-C₁₋₆-alkylene-, aryl-C₂₋₆-alkenylene-, aryl-C₂₋₆-alkynylene-, heteroaryl, heteroaryl-C₁₋₆-alkylene-, heteroaryl-C₂₋₆-alkenylene- and heteroaryl-C₂₋₆-alkynylene-, of which the aryl and heteroaryl moieties optionally may be substituted with one or more substituents selected from halogen, -C(O)OR⁶¹, -CN, -CF₃, -OCF₃, -NO₂, -OR⁶¹, -NR⁶¹R⁶² or C₁₋₆-alkyl, wherein
- 15 R⁶¹ and R⁶² independently of each other are hydrogen, C₁₋₆-alkyl, C₁₋₆-alkyl-arylene-, C₁₋₆-alkyl-heteroarylene-, aryl-C₁₋₆-alkylene-, heteroaryl-C₁₋₆-alkylene-, heteroaryl, or aryl;
- or
- R⁶¹ and R⁶², when attached to the same nitrogen atom, together with the said
- 20 nitrogen atom may form a 3 to 8 membered heterocyclic ring optionally containing one or two further heteroatoms selected from nitrogen, oxygen and sulfur, and optionally containing one or two double bonds.

Embodiment A147. A compound according to embodiment A146, wherein

- 25 R⁵⁶, R⁵⁷, R⁵⁸, R⁵⁹, and R⁶⁰ independently of each other are halogen, -CN, -NO₂, C₁₋₆-alkyl, -CHF₂, -CF₃, -OR⁶¹, -NR⁶¹R⁶², -SR⁶¹, -NR⁶¹S(O)₂R⁶², -S(O)₂NR⁶¹R⁶², -S(O)NR⁶¹R⁶², -S(O)R⁶¹, -S(O)₂R⁶¹, -C(O)NR⁶¹R⁶², -OC(O)NR⁶¹R⁶², -NR⁶¹C(O)R⁶², -CH₂C(O)NR⁶¹R⁶², -CH₂C(O)OR⁶¹, -OCH₂C(O)NR⁶¹R⁶², -CH₂OR⁶¹, -CH₂NR⁶¹R⁶², and -C(O)OR⁶¹; or
- C₂₋₆-alkenyl and C₂₋₆-alkynyl, which may optionally be substituted with one or more
- 30 substituents selected from -CN, -CF₃, -OCF₃, -OR⁶¹, -NR⁶¹R⁶² and C₁₋₆-alkyl, wherein
- R⁶¹ and R⁶² independently of each other are hydrogen, C₁₋₆-alkyl, C₁₋₆-alkyl-arylene-, C₁₋₆-alkyl-heteroarylene-, aryl-C₁₋₆-alkylene-, heteroaryl-C₁₋₆-alkylene-, heteroaryl, or aryl;

or

R^{61} and R^{62} , when attached to the same nitrogen atom, together with the said nitrogen atom may form a 3 to 8 membered heterocyclic ring optionally containing one or two further heteroatoms selected from nitrogen, oxygen and sulfur, and optionally containing one or two double bonds.

Embodiment A148. A compound according to embodiment A146 or embodiment A147, wherein R^{58} or R^{57} is $-OR^{61}$.

Embodiment A149. A compound according to embodiment A148 wherein R^{61} is methyl.

Embodiment A150. A compound according to embodiment A147, wherein

R^{61} and R^{62} independently of each other are hydrogen, or C_{1-6} -alkyl, C_{1-6} -alkyl- C_{3-10} -arylene-, C_{1-6} -alkyl- C_{3-10} -heteroarylene-, C_{3-10} -aryl- C_{1-6} -alkylene-, C_{3-10} -heteroaryl- C_{1-6} -alkylene-, C_{3-10} -heteroaryl, or C_{3-10} -aryl.

Embodiment A151. A compound according to embodiment A150, wherein

R^{61} and R^{62} independently of each other are hydrogen or C_{1-6} -alkyl.

Embodiment A152. A compound according to embodiment A151, wherein

R^{61} is hydrogen.

Embodiment A153. A compound according to embodiment A151 or embodiment A152, wherein

R^{62} is hydrogen.

Embodiment A154. A compound according to any one of the embodiments A146 to A153, wherein

at least one of R^{56} , R^{57} , R^{58} , R^{59} , and R^{60} are hydrogen.

Embodiment A155. A compound according to embodiment A154, wherein

at least two of R^{23} , R^{24} , R^{25} , and R^{26} are hydrogen.

Embodiment A156. A compound according to embodiment A155, wherein

at least three of R^{23} , R^{24} , R^{25} , and R^{26} are hydrogen.

Embodiment A157. A compound according to any one of the embodiments A1 to A156, wherein

L^2 is a direct bond, C_{1-6} -alkylene, C_{2-6} -alkenylene, C_{2-6} -alkynylene, $-N-R^{20}-$, $-C_{1-6}$ -alkylene- $N(R^{20})-$, $-C_{2-6}$ -alkenylene- $N(R^{20})-$, or $-C_{2-6}$ -alkynylene- $N(R^{20})-$, wherein
 5 R^{20} is as defined in embodiment A1.

Embodiment A158. A compound according to any one of the embodiments A1 to A153, wherein

L^2 is $-N-R^{20}-$, $-alkylene-N(R^{20})-$, $-alkenylene-N(R^{20})-$, or $-alkynylene-N(R^{20})-$, wherein
 R^{20} is as defined in embodiment A1.

10 Embodiment A159. A compound according to embodiment A157 or embodiment A158, wherein

L^2 is $-N-R^{20}-$, $-C_{1-6}$ -alkylene- $N(R^{20})-$, $-C_{2-6}$ -alkenylene- $N(R^{20})-$, or $-C_{2-6}$ -alkynylene- $N(R^{20})-$, wherein
 R^{20} is as defined in embodiment A1.

15 Embodiment A160. A compound according to embodiment A159, wherein

L^2 is $-N-R^{20}-$, wherein
 R^{20} is as defined in embodiment A1.

Embodiment A161. A compound according to embodiment A157, wherein

L^2 is a direct bond.

20 Embodiment A162. A compound according to any one of the embodiments A1 to A160, wherein

R^{20} is hydrogen, or
 R^{20} is C_{1-6} -alkyl, C_{2-6} -alkenyl, C_{2-6} -alkynyl, C_{3-10} -cycloalkyl- W^3- ,
 C_{3-10} -heterocyclyl- W^3- , C_{3-10} -aryl- W^3- , or C_{4-10} -heteroaryl- W^3- , optionally substituted
 25 with one or more substituents R^{30} , R^{31} , and R^{32} , wherein
 W^3 , R^{30} , R^{31} , and R^{32} are as defined in embodiment A1.

Embodiment A163. A compound according to embodiment A162, wherein

W^3 is alkylene.

Embodiment A164. A compound according to embodiment A163, wherein
 W^3 is C_{2-6} -alkylene.

Embodiment A165. A compound according to embodiment A162, wherein
 W^3 is a direct bond.

5 Embodiment A166. A compound according to any one of the embodiments A1 to A160, wherein

R^{20} is hydrogen, C_{1-6} -alkyl, C_{2-6} -alkenyl, or C_{2-6} -alkynyl, optionally substituted with one or more substituents R^{30} , R^{31} , and R^{32} , wherein

R^{30} , R^{31} , and R^{32} are as defined in embodiment A1.

10 Embodiment A167. A compound according to any one of the embodiments A162 to A166, wherein

R^{20} is hydrogen, or

R^{20} is C_{1-6} -alkyl, C_{1-6} -alkenyl, or C_{1-6} -alkynyl, optionally substituted with one or more substituents R^{30} , R^{31} , and R^{32} , wherein

15 R^{30} , R^{31} , and R^{32} are as defined in embodiment A1.

Embodiment A168. A compound according to any one of the embodiments A1 to A167, wherein

20 R^{30} , R^{31} , and R^{32} independently of each other are selected from $-CHF_2$, $-CF_3$, $-OCF_3$, $-OCHF_2$, $-OCH_2CF_3$, $-OCF_2CHF_2$, $-S(O)_2CF_3$, $-SCF_3$, $-OR^{52}$, $-NR^{52}R^{53}$, $-SR^{52}$, $-NR^{52}S(O)_2R^{53}$, $-S(O)_2NR^{52}R^{53}$, $-S(O)NR^{52}R^{53}$, $-S(O)R^{52}$, $-S(O)_2R^{52}$, $-C(O)NR^{52}R^{53}$, $-OC(O)NR^{52}R^{53}$, $-NR^{52}C(O)R^{53}$, $-CH_2C(O)NR^{52}R^{53}$, $-OCH_2C(O)NR^{52}R^{53}$, $-CH_2OR^{52}$, $-CH_2NR^{52}R^{53}$, $-OC(O)R^{52}$, $-C(O)R^{52}$ and $-C(O)OR^{52}$, wherein

R^{52} and R^{53} are as defined in embodiment A1.

Embodiment A169. A compound according to embodiment A168, wherein

25 R^{52} and R^{53} , when attached to the same nitrogen atom, together with the said nitrogen atom may form a 3 to 8 membered heterocyclic ring optionally containing one or two further heteroatoms selected from nitrogen, oxygen and sulfur, and optionally containing one or two double bonds.

Embodiment A170. A compound according to any one of the embodiments A1 to A168, wherein

R^{52} and R^{53} independently of each other are hydrogen, C_{1-6} -alkyl, aryl- C_{1-6} -alkylene-heteroaryl- C_{1-6} -alkylene- heteroaryl, or aryl.

- 5 Embodiment A171. A compound according to embodiment A170, wherein
 R^{52} and R^{53} independently of each other are hydrogen, C_{1-6} -alkyl, aryl- C_{1-6} -alkylene-or aryl.

Embodiment A172. A compound according to embodiment A171, wherein

R^{52} and R^{53} independently of each other are hydrogen or C_{1-6} -alkyl.

- 10 Embodiment A173. A compound according to embodiment A172, wherein
 R^{52} is hydrogen.

Embodiment A174. A compound according to embodiment A172 or embodiment A173, wherein

R^{53} is hydrogen.

- 15 Embodiment A175. A compound according to embodiment A167, wherein
 R^{20} is hydrogen.

Embodiment A176. A compound according to any one of the embodiments A1 to A175, wherein

L^3 is -C(O)-, -C(O)-C(O)- or -C(O)CH₂C(O)-.

- 20 Embodiment A177. A compound according to embodiment A176 wherein
 L^3 is -C(O)-.

Embodiment A178. A compound according to embodiment A176 wherein
 L^3 is -C(O)-C(O)-.

Embodiment A179. A compound according to embodiment A176 wherein

- 25 L^3 is -C(O)CH₂C(O)-.

Embodiment A180. A compound according to any one of the embodiments A1 to A179, wherein

R¹ is hydrogen, or

5 R¹ is C₁₋₆-alkyl, C₂₋₆-alkenyl, C₂₋₆-alkynyl, C₃₋₁₀-cycloalkyl-W⁴-, C₃₋₁₀-heterocyclyl-W⁴-, C₃₋₁₀-aryl-W⁴-, or C₄₋₁₀-heteroaryl-W⁴-, optionally substituted with one or more substituents R³³, R³⁴, and R³⁵, wherein W⁴, R³³, R³⁴, and R³⁵ are as defined in embodiment A1.

Embodiment A181. A compound according to embodiment A180, wherein W⁴ is alkylene.

10 Embodiment A182. A compound according to embodiment A181, wherein W⁴ is C₂₋₆-alkylene.

Embodiment A183. A compound according to embodiment A180, wherein W⁴ is a direct bond.

15 Embodiment A184. A compound according to any one of the embodiments A1 to A176, wherein R¹ is hydrogen, C₁₋₆-alkyl, C₂₋₆-alkenyl, or C₂₋₆-alkynyl, optionally substituted with one or more substituents R³³, R³⁴, and R³⁵, wherein R³³, R³⁴, and R³⁵ are as defined in embodiment A1.

20 Embodiment A185. A compound according to any one of the embodiments A180 to A184, wherein R¹ is hydrogen, C₁₋₆-alkyl, C₂₋₆-alkenyl, or C₂₋₆-alkynyl, optionally substituted with one or more substituents R³³, R³⁴, and R³⁵, wherein R³³, R³⁴, and R³⁵ are as defined in embodiment A1.

25 Embodiment A186. A compound according to embodiment A185, wherein R¹ is hydrogen or C₁₋₆-alkyl optionally substituted with one or more substituents R³³, R³⁴, and R³⁵, wherein R³³, R³⁴, and R³⁵ are as defined in embodiment A1.

Embodiment A187. A compound according to any one of the embodiments A1 to A186, wherein

- 5 R^{33} , R^{34} , and R^{35} independently of each other are selected from $-\text{CHF}_2$, $-\text{CF}_3$, $-\text{OCF}_3$, $-\text{OCHF}_2$, $-\text{OCH}_2\text{CF}_3$, $-\text{OCF}_2\text{CHF}_2$, $-\text{S}(\text{O})_2\text{CF}_3$, $-\text{SCF}_3$, $-\text{OR}^{52}$, $-\text{NR}^{52}\text{R}^{53}$, $-\text{SR}_{52}$, $-\text{NR}^{52}\text{S}(\text{O})_2\text{R}^{53}$, $-\text{S}(\text{O})_2\text{NR}^{52}\text{R}^{53}$, $-\text{S}(\text{O})\text{NR}^{52}\text{R}^{53}$, $-\text{S}(\text{O})\text{R}^{52}$, $-\text{S}(\text{O})_2\text{R}^{52}$, $-\text{C}(\text{O})\text{NR}^{52}\text{R}^{53}$, $-\text{OC}(\text{O})\text{NR}^{52}\text{R}^{53}$, $-\text{NR}^{52}\text{C}(\text{O})\text{R}^{53}$, $-\text{CH}_2\text{C}(\text{O})\text{NR}^{52}\text{R}^{53}$, $-\text{OCH}_2\text{C}(\text{O})\text{NR}^{52}\text{R}^{53}$, $-\text{CH}_2\text{OR}^{52}$, $-\text{CH}_2\text{NR}^{52}\text{R}^{53}$, $-\text{OC}(\text{O})\text{R}^{52}$, $-\text{C}(\text{O})\text{R}^{52}$ and $-\text{C}(\text{O})\text{OR}^{52}$, wherein R^{52} and R^{53} are as defined in embodiment A1.

- 10 Embodiment A188. A compound according to embodiment A186, wherein R^{52} and R^{53} , when attached to the same nitrogen atom, together with the said nitrogen atom may form a 3 to 8 membered heterocyclic ring optionally containing one or two further heteroatoms selected from nitrogen, oxygen and sulfur, and optionally containing one or two double bonds.

- 15 Embodiment A189. A compound according to embodiment A186, wherein R^{52} and R^{53} independently of each other are hydrogen, C_{1-6} -alkyl, aryl- C_{1-6} -alkylene-heteroaryl- C_{1-6} -alkylene- heteroaryl, or aryl.

Embodiment A190. A compound according to embodiment A189, wherein R^{52} and R^{53} independently of each other are hydrogen, C_{1-6} -alkyl, aryl- C_{1-6} -alkylene-or aryl.

- 20 Embodiment A191. A compound according to embodiment A190, wherein R^{52} and R^{53} independently of each other are hydrogen or C_{1-6} -alkyl.

Embodiment A192. A compound according to embodiment A191, wherein R^{52} is hydrogen.

- 25 Embodiment A193. A compound according to embodiment A191 or embodiment A192, wherein R^{53} is hydrogen.

Embodiment A194. A compound according to embodiment A186, wherein R^1 is hydrogen.

Embodiment A195. A compound according to any one of the embodiments A1 to A194, wherein

5 G² is heteroaryl, fused heterocycl/heteroaryl, or fused cycloalkylheteroaryl, optionally substituted with one or more substituents R⁴³, R⁴⁴, and R⁴⁵, wherein said heteroaryl group possesses a nitrogen atom adjacent to the atom joining G² with -N(R¹)-, and wherein R⁴³, R⁴⁴, and R⁴⁵ are as defined in embodiment A1.

Embodiment A196. A compound according to embodiment A195, wherein

10 G² is fused cycloalkylheteroaryl optionally substituted with one or more substituents R⁴³, R⁴⁴, and R⁴⁵, wherein said heteroaryl group possesses a nitrogen atom adjacent to the atom joining G² with -N(R¹)-, and wherein R⁴³, R⁴⁴, and R⁴⁵ are as defined in embodiment A1.

Embodiment A197. A compound according to embodiment A196 wherein

15 G² is fused cycloalkylthiazolyl optionally substituted with one or more substituents R⁴³, R⁴⁴, and R⁴⁵, and wherein R⁴³, R⁴⁴, and R⁴⁵ are as defined in embodiment A1.

Embodiment A198. A compound according to embodiment A197, wherein

20 G² is fused cycloalkylthiazolyl optionally substituted with -COOH

Embodiment A199. A compound according to embodiment A195, wherein

25 G² is heteroaryl optionally substituted with one or more substituents R⁴³, R⁴⁴, and R⁴⁵, wherein said heteroaryl group possesses a nitrogen atom adjacent to the atom joining G² with -N(R¹)-, and wherein R⁴³, R⁴⁴, and R⁴⁵ are as defined in embodiment A1.

Embodiment A200. A compound according to embodiment A199, wherein

G² is furanyl, thienyl, thiophenyl, pyrrolyl, imidazolyl, pyrazolyl, triazolyl, tetrazolyl, thiazolyl, oxazolyl, isoxazolyl, oxadiazolyl, thiadiazolyl, isothiazolyl, pyridinyl, pyridazinyl, pyrazinyl, pyrimidinyl, quinolinyl, isoquinolinyl, benzofuranyl,

benzothiophenyl, indolyl, or indazolyl, optionally substituted with one or more substituents R^{43} , R^{44} , and R^{45} , wherein


R^{43} , R^{44} , and R^{45} are as defined in embodiment A1.

Embodiment A201. A compound according to embodiment A200 wherein G^2 is thiazolyl,

5 optionally substituted with one or more substituents R^{43} , R^{44} , and R^{45} , wherein

R^{43} , R^{44} , and R^{45} are as defined in embodiment A1.


Embodiment A202. A compound according to embodiment A201 wherein G^2 is



wherein

10 R^{43} and R^{44} are as defined in embodiment A1.


Embodiment A203. A compound according to embodiment A202 wherein G^2 is



wherein

R^{43} is as defined in embodiment A1.


15 Embodiment A204. A compound according to embodiment A202 wherein G^2 is



wherein

R^{43} and R^{44} are as defined in embodiment A1.

Embodiment A205. A compound according to embodiment A200 wherein G^2 is




20

wherein

R^{43} is as defined in embodiment A1.

Embodiment A206. A compound according to embodiment A202 wherein G^2 is



wherein

R^{44} is as defined in embodiment A1.

Embodiment A207. A compound according to any one of the embodiments A1 to A206,

5 wherein

R^{43} , R^{44} , and R^{45} independently of each other are selected from
 $-\text{CN}$, $-\text{NO}_2$, $-\text{SCF}_3$, $-\text{OR}^{54}$, $-\text{NR}^{54}\text{R}^{55}$, $-\text{SR}^{54}$, $-\text{S}(\text{O})_2\text{NR}^{54}\text{R}^{55}$,
 $-\text{S}(\text{O})\text{NR}^{54}\text{R}^{55}$, $-\text{S}(\text{O})\text{R}^{54}$, $-\text{S}(\text{O})_2\text{R}^{54}$, $-\text{C}(\text{O})\text{NR}^{54}\text{R}^{55}$, $-\text{OC}(\text{O})\text{NR}^{54}\text{R}^{55}$, $-\text{NR}^{54}\text{C}(\text{O})\text{R}^{55}$,
10 $-\text{C}_{1-6}\text{-alkylene-S}(\text{O})\text{R}^{54}$, $-\text{C}_{1-6}\text{-alkylene-S}(\text{O})_2\text{R}^{54}$, $-\text{C}_{1-6}\text{-alkylene-N}(\text{R}^{54})\text{S}(\text{O})_2\text{R}^{55}$,
 $-\text{N}(\text{R}^{54})\text{S}(\text{O})_2\text{R}^{55}$, $-\text{C}_{1-6}\text{-alkylene-C}(\text{O})\text{NR}^{54}\text{R}^{55}$, $-\text{C}_{1-6}\text{-alkylene-N}(\text{R}^{54})\text{C}(\text{O})\text{R}^{55}$,
 $-\text{N}(\text{R}^{54})\text{C}(\text{O})\text{R}^{55}$, $-\text{C}_{1-6}\text{-alkylene-N}(\text{R}^{54})\text{C}(\text{O})\text{NR}^{55}\text{R}^{56}$,
 $\text{C}_{1-6}\text{-alkylene-NHC}(=\text{NR}^{54})\text{NR}^{55}\text{R}^{56}$, $-\text{C}_{1-6}\text{-alkylene-N}(\text{R}^{54})\text{C}(\text{O})\text{OR}^{55}$,
 $-\text{N}(\text{R}^{54})\text{C}(\text{O})\text{OR}^{55}$, $-\text{C}_{1-6}\text{-alkylene-C}(\text{O})\text{OR}^{54}$, $-\text{OCH}_2\text{C}(\text{O})\text{NR}^{54}\text{R}^{55}$, $-\text{O}(\text{CH}_2)_{1-3}\text{OR}^{54}$,
15 $-\text{C}_{1-6}\text{-alkylene-O-R}^{54}$, $-\text{C}_{1-6}\text{-alkylene-C}(\text{O})\text{R}^{54}$, $-\text{C}_{1-6}\text{-alkylene-NR}^{54}\text{R}^{55}$,
 $-\text{C}_{1-6}\text{-alkylene=N-O-R}^{54}$, $-\text{C}_{1-6}\text{-alkylene-N}(\text{R}^{54})\text{S}(\text{O})_2\text{NR}^{55}\text{R}^{56}$, $-\text{N}(\text{R}^{54})\text{S}(\text{O})_2\text{NR}^{55}\text{R}^{56}$,
 $-\text{N}(\text{R}^{54})\text{S}(\text{O})_2\text{NR}^{56}\text{R}^{56}$, $-\text{OC}(\text{O})\text{R}^{54}$, $-\text{C}(\text{O})\text{N}(\text{R}^{54})\text{S}(\text{O})_2\text{R}^{55}$,
 $-\text{C}_{1-6}\text{-alkylene-C}(\text{R}^{54})=\text{N-OR}^{55}$, $-\text{NHC}(=\text{NR}^{54})\text{NR}^{55}\text{R}^{56}$,
 $-\text{C}_{1-6}\text{-alkylene-N}=\text{C}(\text{N}(\text{R}^{54}\text{R}^{55}))_2$, $-\text{N}=\text{C}(\text{N}(\text{R}^{54}\text{R}^{55}))_2$, $-\text{C}(\text{O})\text{R}^{54}$ and $-\text{C}(\text{O})\text{OR}^{54}$; or
20 $\text{C}_{1-6}\text{-alkyl}$ or $\text{C}_{2-6}\text{-alkenyl}$, each of which may optionally be substituted with one or
more substituents independently selected from halogen, R^{54} , $-\text{CN}$, $-\text{CF}_3$, $-\text{OCF}_3$, $-\text{OR}^{54}$,
 $-\text{C}(\text{O})\text{OR}^{54}$, $-\text{NR}^{54}\text{R}^{55}$ and $\text{C}_{1-6}\text{-alkyl}$; or
 $\text{C}_{3-10}\text{-cycloalkyl}$, heterocyclyl, $\text{C}_{3-10}\text{-cycloalkyl-C}_{1-6}\text{-alkylene-}$,
 $\text{C}_{3-10}\text{-cycloalkyl-C}_{1-6}\text{-alkoxy-}$, $\text{C}_{3-10}\text{-cycloalkyloxy}$, heterocyclyl- $\text{C}_{1-6}\text{-alkylene-}$, of which
25 the heterocyclyl moieties optionally may be substituted with one or more
substituents independently selected from R^{70} ; or
aryl, aryloxy, aryl- $\text{C}_{1-6}\text{-alkoxy-}$, aryl- $\text{C}_{1-6}\text{-alkylene-}$, heteroaryl,
heteroaryl- $\text{C}_{1-6}\text{-alkylene-}$, of which the aryl and heteroaryl moieties optionally may be
substituted with one or more substituents selected from halogen, $-\text{C}(\text{O})\text{OR}^{54}$, $-\text{CN}$,
30 $-\text{CF}_3$, $-\text{OCF}_3$, $-\text{NO}_2$, $-\text{OR}^{54}$, $-\text{NR}^{54}\text{R}^{55}$ or $\text{C}_{1-6}\text{-alkyl}$, wherein
 R^{54} , R^{55} and R^{56} are as defined in embodiment A1.

Embodiment A208. A compound according to embodiment A207, wherein

- R^{43} , R^{44} , and R^{45} independently of each other are selected from
 $-OR^{54}$, $-NR^{54}R^{55}$, $-SR^{54}$, $-S(O)R^{54}$, $-S(O)_2R^{54}$, $-C(O)NR^{54}R^{55}$, halogen,
 $-S-C_{1-6}\text{-alkylene-}OR^{54}$, $-C_{1-6}\text{-alkylene-S-}R^{54}$, $-C_{1-6}\text{-alkylene-S(O)}R^{54}$,
5 $-C_{1-6}\text{-alkylene-S(O)}_2R^{54}$, $-C_{1-6}\text{-alkylene-N(R}^{54})S(O)_2R^{55}$, $-C_{1-6}\text{-alkylene-C(O)NR}^{54}R^{55}$,
 $-C_{1-6}\text{-alkylene-N(R}^{54})C(O)R^{55}$, $-C_{1-6}\text{-alkylene-N(R}^{54})C(O)NR^{55}R^{56}$,
 $C_{1-6}\text{-alkylene-NHC(=NR}^{54})NR^{55}R^{56}$, $-C_{1-6}\text{-alkylene-N(R}^{54})C(O)OR^{55}$,
 $-C_{1-6}\text{-alkylene-C(O)OR}^{54}$, $-C_{1-6}\text{-alkylene-O-}R^{54}$, $-C_{1-6}\text{-alkylene-C(O)}R^{54}$,
 $-C_{1-6}\text{-alkylene-NR}^{54}R^{55}$, $-C_{1-6}\text{-alkylene=N-O-}R^{54}$, $-C_{1-6}\text{-alkylene-N=C(N(R}^{54}R^{55}))_2$,
10 $-N=C(N(R^{54}R^{55}))_2$, $-C(O)R^{54}$ and $-C(O)OR^{54}$; or
 C_{1-6} -alkyl optionally substituted with one or more substituents independently
selected from halogen, R^{54} , $-CN$, $-CF_3$, $-OCF_3$, $-OR^{54}$, $-C(O)OR^{54}$, $-NR^{54}R^{55}$ and
 C_{1-6} -alkyl; or
Heterocyclyl or heterocyclyl- C_{1-6} -alkylene-, of which the heterocyclyl moieties
15 optionally may be substituted with one or more substituents independently selected
from R^{70} ; or
aryl, aryl- C_{1-6} -alkylene-, heteroaryl, heteroaryl- C_{1-6} -alkylene-, of which the aryl and
heteroaryl moieties optionally may be substituted with one or more substituents
selected from halogen, $-C(O)OR^{54}$, $-CN$, $-CF_3$, $-OCF_3$, $-NO_2$, $-OR^{54}$, $-NR^{54}R^{55}$ or
20 C_{1-6} -alkyl, wherein
 R^{54} , R^{55} , R^{56} , and R^{70} are as defined in embodiment A1.

Embodiment A209. A compound according to embodiment A208, wherein

- R^{43} , R^{44} , and R^{45} independently of each other are selected from
 $-SR^{54}$, $-S(O)R^{54}$, $-S(O)_2R^{54}$, halogen, $-C_{1-6}\text{-alkylene-S-}R^{54}$, $-C_{1-6}\text{-alkylene-S(O)}R^{54}$,
25 $-C_{1-6}\text{-alkylene-S(O)}_2R^{54}$, $-C_{1-6}\text{-alkylene-N(R}^{54})C(O)OR^{55}$, $-C_{1-6}\text{-alkylene-C(O)OR}^{54}$,
 $-C_{1-6}\text{-alkylene-O-}R^{54}$, $-C_{1-6}\text{-alkylene-NR}^{54}R^{55}$, $-C_{1-6}\text{-alkylene-C(O)}R^{54}$, $-C(O)R^{54}$ and
 $-C(O)OR^{54}$; or
 C_{1-6} -alkyl optionally substituted with one or more substituents independently
selected from halogen, R^{54} , $-CN$, $-CF_3$, $-OCF_3$, $-OR^{54}$, $-C(O)OR^{54}$, $-NR^{54}R^{55}$ and
30 C_{1-6} -alkyl; or
Heterocyclyl or heterocyclyl- C_{1-6} -alkylene-, of which the heterocyclyl moieties
optionally may be substituted with one or more substituents independently selected
from R^{70} ; or

heteroaryl or heteroaryl-C₁₋₆-alkylene-, of which the aryl and heteroaryl moieties optionally may be substituted with one or more substituents selected from halogen, -C(O)OR⁵⁴, -CN, -CF₃, -OCF₃, -NO₂, -OR⁵⁴, -NR⁵⁴R⁵⁵ or C₁₋₆-alkyl, wherein R⁵⁴, R⁵⁵, R⁵⁶, and R⁷⁰ are as defined in embodiment A1.

- 5 Embodiment A210. A compound according to embodiment A209, wherein R⁴³, R⁴⁴, and R⁴⁵ independently of each other are selected from -SR⁵⁴, -S(O)₂R⁵⁴, halogen, -C₁₋₆-alkylene-C(O)OR⁵⁴, and -C(O)OR⁵⁴; or Heterocyclyl or heterocyclyl-C₁₋₆-alkylene-, of which the heterocyclyl moieties optionally may be substituted with one or more substituents independently selected
10 from R⁷⁰; or heteroaryl or heteroaryl-C₁₋₆-alkylene-, of which the heteroaryl moieties optionally may be substituted with one or more substituents selected from halogen, -C(O)OR⁵⁴, -CN, -CF₃, -OCF₃, -NO₂, -OR⁵⁴, -NR⁵⁴R⁵⁵ or C₁₋₆-alkyl, wherein R⁵⁴, R⁵⁵, R⁵⁶, and R⁷⁰ are as defined in embodiment A1.
- 15 Embodiment A211. A compound according to embodiment A210, wherein R⁴³, R⁴⁴, and R⁴⁵ independently of each other are selected from -SR⁵⁴, -S(O)₂R⁵⁴, halogen, -C₁₋₆-alkylene-C(O)OR⁵⁴, and -C(O)OR⁵⁴; or heterocyclyl-C₁₋₆-alkylene-, wherein heterocyclyl is selected from imidazolyl, piperidyl, piperazinyl, and morpholinyl, and
20 of which the heterocyclyl moieties optionally may be substituted with one or more substituents independently selected from R⁷⁰; or heteroaryl or heteroaryl-C₁₋₆-alkylene-, wherein heteroaryl is selected from thiazolyl, triazolyl, or tetrazolyl, and of which the heteroaryl moieties optionally may be substituted with one or more substituents selected from halogen, -C(O)OR⁵⁴, -CN,
25 -CF₃, -OCF₃, -NO₂, -OR⁵⁴, -NR⁵⁴R⁵⁵ or C₁₋₆-alkyl, wherein R⁵⁴, R⁵⁵, R⁵⁶, and R⁷⁰ are as defined in embodiment A1.
- Embodiment A212. A compound according to embodiment A209, wherein R⁴³, R⁴⁴, and R⁴⁵ independently of each other are selected from -C₁₋₆-alkylene-S-R⁵⁴, -C₁₋₆-alkylene-O-R⁵⁴, -C₁₋₆-alkylene-S(O)₂R⁵⁴ or
30 -C₁₋₆-alkylene-NR⁵⁴R⁵⁵, wherein R⁵⁴ and R⁵⁵ are as defined in embodiment A1.
- Embodiment A213. A compound according to embodiment A209, wherein R⁴³ is -C₁₋₆-alkylene-S-R⁵⁴, wherein

R⁵⁴ is as defined in embodiment A1.

Embodiment A214. R⁴³ is -C₁₋₈-alkylene-O-R⁵⁴, wherein

R⁵⁴ is as defined in embodiment A1.

Embodiment A215. R⁴³ is -C₁₋₈-alkylene-NR⁵⁴R⁵⁵, wherein

5 R⁵⁴ and R⁵⁵ are as defined in embodiment A1.

Embodiment A216. R⁴³ is -C₁₋₈-alkylene-S(O)₂R⁵⁴, wherein

R⁵⁴ is as defined in embodiment A1.

Embodiment A217. A compound according to any one of the embodiments A207 to A210 wherein R⁴³ is heteroaryl or heteroaryl-C₁₋₈-alkylene- optionally substituted with one or more
10 substituents selected from halogen, -C(O)OR⁵⁴, -CN, -CF₃, -OCF₃, -NO₂, -OR⁵⁴, -NR⁵⁴R⁵⁵ or C₁₋₈-alkyl, wherein R⁵⁴ and R⁵⁵ are as defined in embodiment A1.

Embodiment A218. A compound according to embodiment A217 wherein R⁴³ is heteroaryl optionally substituted with one or more substituents selected from halogen, -C(O)OR⁵⁴, -CN, -CF₃, -OCF₃, -NO₂, -OR⁵⁴, -NR⁵⁴R⁵⁵ or C₁₋₈-alkyl, wherein R⁵⁴ and R⁵⁵ are as defined in
15 embodiment A1.

Embodiment A219. A compound according to embodiment A217 wherein R⁴³ is heteroaryl-C₁₋₈-alkylene- optionally substituted with one or more substituents selected from halogen, -C(O)OR⁵⁴, -CN, -CF₃, -OCF₃, -NO₂, -OR⁵⁴, -NR⁵⁴R⁵⁵ or C₁₋₈-alkyl, wherein R⁵⁴ and R⁵⁵ are as defined in embodiment A1.

20 Embodiment A220. A compound according to any one of the embodiments A207 to A211 wherein R⁷⁰ is =O, methyl or C(O)OR⁷⁵.

Embodiment A221. A compound according to any one of the embodiments A207 to A211 wherein R⁷⁰ is -C(O)OH

Embodiment A222. A compound according to any one of the embodiments A207 to A211
25 wherein R⁷⁰ is -(CH₂)₁₋₃C(O)OH.

Embodiment A223. A compound according to any one of the embodiments A207 to A211 wherein R⁷⁰ is -S(O)₂CH₃.

Embodiment A224. A compound according to embodiment A211 wherein R⁴³ is -C₁₋₈-alkylene-C(O)OR⁵⁴.

30 Embodiment A225. A compound according to embodiment A224 wherein R⁴³ is -CH₂-C(O)OR⁵⁴.

Embodiment A226. A compound according to embodiment A211 wherein R⁴⁴ is -SR⁵⁴.

Embodiment A227. A compound according to embodiment A211 wherein R⁴⁴ is -S(O)₂R⁵⁴.

Embodiment A228. A compound according to embodiment A211 wherein R⁴⁴ is halogen.

Embodiment A229. A compound according to any one of the embodiments A1 to A228 wherein

R^{54} , R^{55} and R^{56} independently of each other are hydrogen, C_{1-6} -alkyl, aryl- C_{1-6} -alkylene-, heteroaryl- C_{1-6} -alkylene-, heterocyclyl, heterocyclyl- C_{1-6} -alkylene-, heteroaryl, or aryl, each of which is optionally substituted with one or more substituents independently selected from R^{71} ;

or

R^{54} and R^{55} independently of each other are hydrogen or $-(CHR^{72})_u-(CHR^{73})_v-W^6$,

or

R^{54} and R^{55} , when attached to the same nitrogen atom, together with the said nitrogen atom may form a 3 to 8 membered heterocyclic ring optionally containing one or two further heteroatoms selected from nitrogen, oxygen and sulfur, and optionally containing one or two double bonds, and optionally substituted with one or more C_{1-6} -alkyl groups,

wherein R^{71} , R^{72} , R^{73} , u , v , and W^6 are as defined in embodiment A1.

Embodiment A230. A compound according to embodiment A229 wherein

R^{54} , R^{55} and R^{56} independently of each other are hydrogen, C_{1-6} -alkyl, heterocyclyl, heterocyclyl- C_{1-6} -alkylene-, heteroaryl, or aryl, each of which is optionally substituted with one or more substituents independently selected from R^{71} ;

or

R^{54} and R^{55} independently of each other are hydrogen or $-(CHR^{72})_u-(CHR^{73})_v-W^6$,

wherein R^{71} , R^{72} , R^{73} , u , v , and W^6 are as defined in embodiment A1.

Embodiment A231. A compound according to any one of the embodiments A1 to A230 wherein

R^{54} , R^{55} and R^{56} independently of each other are hydrogen, methyl, ethyl, propyl, butyl, isopropyl, isobutyl, *sec*-butyl, *tert*-butyl, 3-pentyl, 2-pentyl, or 3-methyl-butyl.

Embodiment A232. A compound according to embodiment A231 wherein

R^{54} , R^{55} and R^{56} independently of each other are hydrogen, methyl, ethyl, propyl, butyl, isopropyl, isobutyl, *sec*-butyl, or *tert*-butyl.

Embodiment A233. A compound according to embodiment A232 wherein

R^{54} , R^{55} and R^{56} independently of each other are hydrogen, methyl, ethyl, propyl, butyl, isopropyl, isobutyl, *sec*-butyl, or *tert*-butyl.

Embodiment A234. A compound according to embodiment A233 wherein

R⁵⁴, R⁵⁵ and R⁵⁶ independently of each other are hydrogen, methyl or ethyl.

Embodiment A235. A compound according to embodiment A234 wherein

R⁵⁴ is hydrogen.

Embodiment A236. A compound according to any one of the embodiments A1 to A230

5 wherein

R⁵⁴, R⁵⁵ and R⁵⁶ independently of each other are oxetanyl, tetrahydrofuranyl, tetrahydropyranyl, oxepanyl, azetidyl, pyrrolidyl, piperidyl, hexahydroazepinyl, thietanyl, thiolanyl, tetrahydrothiopyranyl, thiepanyl, 1,4-oxathianyl, 1,3-dioxolanyl, 1,2-dithiolanyl, 1,3-dithiolanyl, hexahydro-pyridazinyl, imidazolidyl, 1,3-dioxanyl, 10 morpholiny, 1,3-dithianyl, 1,4-dioxanyl, 1,4-dithianyl, or thiomorpholiny, each of which is optionally substituted with one or more substituents independently selected from R⁷¹, wherein R⁷¹ is as defined in embodiment A1.

Embodiment A237. A compound according to embodiment A236 wherein

R⁵⁴, R⁵⁵ and R⁵⁶ independently of each other are tetrahydropyranyl, oxepanyl, 15 piperidyl, hexahydroazepinyl, tetrahydrothiopyranyl, thiepanyl, 1,4-oxathianyl, 1,3-dithiolanyl, hexahydro-pyridazinyl, 1,3-dioxanyl, morpholiny, 1,3-dithianyl, 1,4-dioxanyl, 1,4-dithianyl, or thiomorpholiny, each of which is optionally substituted with one or more substituents independently selected from R⁷¹, wherein R⁷¹ is as defined in embodiment A1.

20 Embodiment A238. A compound according to embodiment A237 wherein

R⁵⁴, R⁵⁵ and R⁵⁶ independently of each other are tetrahydropyranyl, piperidyl, tetrahydrothiopyranyl, 1,4-oxathianyl, hexahydro-pyridazinyl, morpholiny, 1,4-dioxanyl, 1,4-dithianyl, or thiomorpholiny, each of which is optionally substituted with one or more substituents independently selected from R⁷¹, wherein R⁷¹ is as defined in embodiment A1.

25 Embodiment A239. A compound according to embodiment A238 wherein

R⁵⁴, R⁵⁵ and R⁵⁶ independently of each other are tetrahydropyranyl, piperidyl, tetrahydrothiopyranyl, or morpholiny, each of which is optionally substituted with one or more substituents independently selected from R⁷¹, wherein R⁷¹ is as defined in embodiment A1.

30 Embodiment A240. A compound according to embodiment A239 wherein

R⁵⁴, R⁵⁵ and R⁵⁶ independently of each other are piperidyl or morpholiny, each of which is optionally substituted with one or more substituents independently selected from R⁷¹, wherein R⁷¹ is as defined in embodiment A1.

Embodiment A241. A compound according to embodiment A240 wherein

35 R⁵⁴, R⁵⁵ and R⁵⁶ independently of each other are piperidyl or morpholiny.

242.

Embodiment A243. A compound according to any one of the embodiments A1 to A230 wherein

5 R^{54} , R^{55} and R^{56} independently of each other are furanyl, thienyl, thiophenyl, pyrrolyl, imidazolyl, pyrazolyl, triazolyl, tetrazolyl, thiazolyl, oxazolyl, isoxazolyl, oxadiazolyl, thiadiazolyl, isothiazolyl, pyridinyl, pyridazinyl, pyrazinyl, pyrimidinyl, quinolinyl, isoquinolinyl, benzofuranyl, benzothiophenyl, indolyl, purinyl, or indazolyl, each of which is optionally substituted with one or more substituents independently selected from R^{71} , wherein R^{71} is as defined in embodiment A1.

10 Embodiment A244. A compound according to embodiment A243 wherein

R^{54} , R^{55} and R^{56} independently of each other are furanyl, thienyl, thiophenyl, pyrrolyl, imidazolyl, pyrazolyl, triazolyl, tetrazolyl, thiazolyl, oxazolyl, isoxazolyl, oxadiazolyl, thiadiazolyl, isothiazolyl, pyridinyl, pyridazinyl, pyrazinyl, pyrimidinyl, benzofuranyl, indolyl, or purinyl, each of which is optionally substituted with one or more
15 substituents independently selected from R^{71} , wherein R^{71} is as defined in embodiment A1.

Embodiment A245. A compound according to embodiment A244 wherein

R^{54} , R^{55} and R^{56} independently of each other are imidazolyl, triazolyl, tetrazolyl, thiazolyl, pyridinyl, pyrimidinyl, benzofuranyl, indolyl, or purinyl, each of which is
20 optionally substituted with one or more substituents independently selected from R^{71} , wherein R^{71} is as defined in embodiment A1.

Embodiment A246. A compound according to embodiment A245 wherein

R^{54} , R^{55} and R^{56} independently of each other are imidazolyl, triazolyl, tetrazolyl, thiazolyl, pyridinyl, pyrimidinyl, or purinyl, each of which is optionally substituted with
25 one or more substituents independently selected from R^{71} , wherein R^{71} is as defined in embodiment A1.

Embodiment A247. A compound according to embodiment A246 wherein

R^{54} , R^{55} and R^{56} independently of each other are imidazolyl, triazolyl, tetrazolyl, thiazolyl, pyridinyl, pyrimidinyl, each of which is optionally substituted with one or
30 more substituents independently selected from R^{71} , wherein R^{71} is as defined in embodiment A1.

Embodiment A248. A compound according to embodiment A247 wherein R^{54} is imidazolyl optionally substituted with one or more substituents independently selected from R^{71} , wherein R^{71} is as defined in embodiment A1.

Embodiment A249. A compound according to embodiment A247 wherein R^{54} is triazolyl optionally substituted with one or more substituents independently selected from R^{71} , wherein R^{71} is as defined in embodiment A1.

5 Embodiment A250. A compound according to embodiment A247 wherein R^{54} is tetrazolyl optionally substituted with one or more substituents independently selected from R^{71} , wherein R^{71} is as defined in embodiment A1.

Embodiment A251. A compound according to embodiment A247 wherein R^{54} is thiazolyl optionally substituted with one or more substituents independently selected from R^{71} , wherein R^{71} is as defined in embodiment A1.

10 Embodiment A252. A compound according to embodiment A247 wherein R^{54} is pyridinyl optionally substituted with one or more substituents independently selected from R^{71} , wherein R^{71} is as defined in embodiment A1.

Embodiment A253. A compound according to embodiment A247 wherein R^{54} is pyrimidinyl optionally substituted with one or more substituents independently selected from R^{71} , wherein
15 R^{71} is as defined in embodiment A1.

Embodiment A254. A compound according to embodiment A246 wherein R^{54} is purinyl optionally substituted with one or more substituents independently selected from R^{71} , wherein R^{71} is as defined in embodiment A1.

Embodiment A255. A compound according to any one of the embodiments A1 to A254
20 wherein R^{71} is methyl or =O.

Embodiment A256. A compound according to any one of the embodiments A1 to A254 wherein R^{71} is $-C(O)OH$

Embodiment A257. A compound according to any one of the embodiments A1 to A254 wherein R^{71} is $-(CH_2)_{1-3}C(O)OH$.

25 Embodiment A258. A compound according to any one of the embodiments A1 to A254 wherein R^{70} is $-(CH_2)_{1-3}NR^{75}R^{76}$.

Embodiment A259. A compound according to any one of the embodiments A1 to A230 wherein u is 0 or 1.

Embodiment A260. A compound according to embodiment A259 wherein u is 0.

30 Embodiment A261. A compound according to embodiment A259 wherein u is 1.

Embodiment A262. A compound according to any one of the embodiments A1 to A230 wherein v is 0 or 1.

Embodiment A263. A compound according to embodiment A262 wherein v is 0.

Embodiment A264. A compound according to embodiment A262 wherein v is 1.

Embodiment A265. A compound according to any one of the embodiments A1 to A230 wherein u is 0 and v is 1.

Embodiment A266. A compound according to any one of the embodiments A1 to A230 wherein u and v are both 0.

- 5 Embodiment A267. A compound according to any one of the embodiments A1 to A230 or A259 to A266 wherein R^{72} and R^{73} are independently selected from hydrogen, hydroxy or $-C(O)OR^{75}$.

Embodiment A268. A compound according to embodiment A267 wherein R^{72} and R^{73} are independently selected from hydrogen or $-C(O)OR^{75}$, wherein R^{75} is as defined in
10 embodiment A1.

Embodiment A269. A compound according to embodiment A267 wherein R^{72} and R^{73} are hydrogen.

Embodiment A270. A compound according to any one of the embodiments A1 to A230 or A259 to A269 wherein

- 15 W^6 is $-O-R^{75}$, $-C(O)O-R^{75}$, $-C(O)-R^{75}$, $-NR^{75}R^{76}$, $-NHCH_2C(O)R^{75}$, $-NHC(O)R^{75}$, $-S(O)_2R^{75}$, $-NHS(O)_2R^{75}$, or
 W^6 is heterocyclyl, wherein R^{75} and R^{76} are as defined in embodiment A1.

Embodiment A271. A compound according to embodiment A270 wherein

- 20 W^6 is $-O-R^{75}$, $-C(O)O-R^{75}$, $-C(O)-R^{75}$, $-NR^{75}R^{76}$, $-NHCH_2C(O)R^{75}$, $-NHC(O)R^{75}$, $-S(O)_2R^{75}$, $-NHS(O)_2R^{75}$, or
 W^6 is oxetanyl, tetrahydrofuranyl, tetrahydropyranyl, oxepanyl, azetidyl, pyrrolidyl, piperidyl, hexahydroazepinyl, thietanyl, thiolanyl, tetrahydrothiopyranyl, thiepanyl, 1,4-oxathianyl, 1,3-dioxolanyl, 1,2-dithiolanyl, 1,3-dithiolanyl, hexahydro-pyridazinyl, imidazolidyl, 1,3-dioxanyl,
25 morpholinyl, 1,3-dithianyl, 1,4-dioxanyl, 1,4-dithianyl, or thiomorpholinyl, wherein R^{75} and R^{76} are as defined in embodiment A1.

Embodiment A272. A compound according to embodiment A270 wherein

- 30 W^6 is $-O-R^{75}$, $-C(O)O-R^{75}$, $-C(O)-R^{75}$, $-NR^{75}R^{76}$, $-NHCH_2C(O)R^{75}$, $-NHC(O)R^{75}$, $-S(O)_2R^{75}$, $-NHS(O)_2R^{75}$, or
 W^6 is tetrahydropyranyl, oxepanyl, piperidyl, hexahydroazepinyl, tetrahydrothiopyranyl, thiepanyl, 1,4-oxathianyl, morpholinyl, 1,4-dioxanyl, 1,4-dithianyl, or thiomorpholinyl, wherein R^{75} and R^{76} are as defined in embodiment A1.

Embodiment A273. A compound according to embodiment A272 wherein

W^6 is $-O-R^{75}$, $-C(O)O-R^{75}$, $-C(O)-R^{75}$, $-NR^{75}R^{76}$, $-NHCH_2C(O)R^{75}$,
 $-NHC(O)R^{75}$, $-S(O)_2R^{75}$, $-NHS(O)_2R^{75}$, or

W^6 is tetrahydropyranyl, piperidyl, tetrahydrothiopyranyl, or morpholinyl,
wherein R^{75} and R^{76} are as defined in embodiment A1.

5 Embodiment A274. A compound according to embodiment A273 wherein

W^6 is $-O-R^{75}$, $-C(O)O-R^{75}$, $-NR^{75}R^{76}$, $-NHC(O)R^{75}$, $-S(O)_2R^{75}$, or

W^6 is tetrahydropyranyl, piperidyl, tetrahydrothiopyranyl, or morpholinyl,
wherein R^{75} and R^{76} are as defined in embodiment A1.

Embodiment A275. A compound according to embodiment A274 wherein

10 W^6 is $-O-R^{75}$, or $-C(O)O-R^{75}$, wherein R^{75} is as defined in embodiment A1.

Embodiment A276. A compound according to embodiment A275 wherein W^6 is $-C(O)O-R^{75}$,
wherein R^{75} is as defined in embodiment A1..

Embodiment A277. A compound according to any one of the embodiments A1 to A230 or
A259 to A276 wherein R^{75} and R^{76} are independently selected from hydrogen, $-OH$, or

15 C_{1-6} -alkyl optionally substituted with $-NH_2$.

Embodiment A278. A compound according to embodiment A277 wherein R^{75} and R^{76} are
independently selected from hydrogen, $-OH$, or methyl.

Embodiment A279. A compound according to embodiment A278 wherein R^{75} and R^{76} are
independently selected from hydrogen or $-OH$.

20 Embodiment A280. A compound according to embodiment A279 wherein R^{75} is hydrogen.

Embodiment A281. A compound according to any one of the embodiments A1 to A280,
which compound is an activator of glucokinase, when tested in the Glucokinase Activation
Assay (I) disclosed herein at a glucose concentration of 2 mM.

Embodiment A282. A compound according to any one of the embodiments A1 to A281,
25 which compound is an activator of glucokinase, when tested in the Glucokinase Activation
Assay (I) disclosed herein at a glucose concentration of from 10 to 15 mM.

Embodiment A283. A compound according to any one of the embodiments A1 to A282,
which compound, at a concentration of 30 μM , is capable of providing an at least 1.5, such
as at least 1.7, for instance at least 2.0 fold activation of glucokinase in the Glucokinase
30 Activation Assay (I) disclosed herein at a glucose concentration of 2 mM.

Embodiment A284. A compound according to any one of the embodiments A1 to A283, which compound, at a concentration of 30 μ M, is capable of providing an at least 1.5, such as at least 1.7, for instance at least 2.0 fold activation of glucokinase in the Glucokinase Activation Assay (I) disclosed herein at a glucose concentration of from 10 to 15 mM.

- 5 Embodiment A285. A compound according to any one of the embodiments A1 to A284, which at a concentration of 5 μ M is capable of providing an at least 1.5, such as at least 1.7, for instance at least 2.0 fold activation of glucokinase in the Glucokinase Activation Assay (I) disclosed herein at a glucose concentration of 2 mM.

- 10 Embodiment A286. A compound according to any one of the embodiments A1 to A285, which at a concentration of 5 μ M is capable of providing an at least 1.5, such as at least 1.7, for instance at least 2.0 fold activation of glucokinase in the Glucokinase Activation Assay (I) disclosed herein at a glucose concentration of from 10 to 15 mM.

- 15 Embodiment A287. A glucose kinase activator compound defined as a compound which at a compound concentration of at or below 30 μ M at a glucose concentration of 2 mM in the Glucokinase Activation Assay (I) disclosed herein gives 1.5 - fold higher glucokinase activity than measured at a glucose concentration of 2 mM in the Glucokinase Activation Assay (I) without compound, where the increase in glucokinase activity provided by the compound increases with increasing concentrations of glucose.

- 20 Embodiment A288. A compound according to any one of the embodiments A1 to A286, which compound provides an increase in glucokinase activity, where the increase in glucokinase activity provided by the compound increases with increasing concentrations of glucose.

- 25 Embodiment A289. A compound according to embodiment A287 or embodiment A288, which provides an increase in glucokinase activity in Glucokinase Activation Assay (I) disclosed herein at a glucose concentration of 15 mM, which increase is significantly higher than the increase in glucokinase activity provided by the compound in Glucokinase Activation Assay (I) disclosed herein at a glucose concentration of 5 mM.

Embodiment A290. A compound according to any one of the embodiments A287 to A289, which at a compound concentration of 10 μ M provides an increase in glucokinase activity in

Glucokinase Activation Assay (I) disclosed herein at a glucose concentration of 15 mM, which increase is significantly higher than the increase in glucokinase activity provided by the compound at a compound concentration of 10 μ M in Glucokinase Activation Assay (I) disclosed herein at a glucose concentration of 5 mM.

- 5 Embodiment A291. A compound according to any one of the embodiments A287 to A290, which at a compound concentration of 10 μ M provides an increase in glucokinase activity in Glucokinase Activation Assay (I) disclosed herein at a glucose concentration of 15 mM, which increase is at least 1.1 fold higher, such as at least 1.2 fold higher, for instance at least 1.3 fold higher, such as at least 1.4 fold higher, for instance 1.5 fold higher, such as at least
10 1.6 fold higher, for instance at least 1.7 fold higher, such as at least 1.8 fold higher, for instance at least 1.9 fold higher, such as at least 2.0 fold higher than the increase in glucokinase activity provided by the compound at a compound concentration of 10 μ M in Glucokinase Activation Assay (I) disclosed herein at a glucose concentration of 5 mM.

- Embodiment A292. A glucose kinase activator compound defined as a compound which at a
15 compound concentration of at or below 30 μ M at a glucose concentration of 2 mM in the Glucokinase Activation Assay (I) disclosed herein gives 1.5 - fold higher glucokinase activity than measured at a glucose concentration of 2 mM in the Glucokinase Activation Assay (I) without compound, which glucose kinase activator compound increases glucose utilization in the liver without inducing any increase in insulin secretion in response to glucose.

- 20 Embodiment A293. A glucose kinase activator compound defined as a compound which at a compound concentration of at or below 30 μ M at a glucose concentration of 2 mM in the Glucokinase Activation Assay (I) disclosed herein gives 1.5 - fold higher glucokinase activity than measured at a glucose concentration of 2 mM in the Glucokinase Activation Assay (I) without compound, which glucokinase activator compound shows a significantly higher
25 activity in isolated hepatocytes compared to the activity of the glucokinase compound in Ins-1 cells.

Embodiment A294. A compound according to any one of the embodiments A1 to A291, which compound increases glucose utilization in the liver without inducing any increase in insulin secretion in response to glucose.

Embodiment A295. A compound according to any one of the embodiments A1 to A291, which compound shows a significantly higher activity in isolated hepatocytes compared to the activity of the compound in Ins-1 cells.

5 Embodiment A296. A compound according to any one of the embodiments A292 to A295, which compound shows a significantly higher activity in isolated hepatocytes measured as described in the Glucokinase Activity Assay (II) compared to the activity of the compound in Ins-1 cells measured as described in the Glucokinase Activity Assay (III).

Embodiment A297. A compound according to embodiment A296, which compound shows an activity in isolated hepatocytes measured as described in the Glucokinase Activity Assay (II)
10 which activity is at least 1.1 fold higher, such as at least 1.2 fold higher, for instance at least 1.3 fold higher, such as at least 1.4 fold higher, for instance 1.5 fold higher, such as at least 1.6 fold higher, for instance at least 1.7 fold higher, such as at least 1.8 fold higher, for instance at least 1.9 fold higher, such as at least 2.0 fold higher, for instance at least a 3.0 fold higher, such as at least a 4.0 fold higher, for instance at least 5.0 fold higher, such as at
15 least 10 fold higher than the activity of the compound in Ins-1 cells measured as described in the Glucokinase Activity Assay (III).

Embodiment A298. A compound according to embodiment A296, which compound shows no activity in the Ins-1 cells measured as described in the Glucokinase Activity Assay (III).

Embodiment A299. A method of preventing hypoglycaemia comprising administration of a
20 compound according to any one of the embodiments A1 to A298.

Embodiment A300. The use of a compound according to any one of the embodiments A1 to Embodiment A298 for the preparation of a medicament for the prevention of hypoglycaemia.

Embodiment A301. A compound according to any one of embodiments A1 to A298, which is an agent useful for the treatment of an indication selected from the group consisting of
25 hyperglycemia, IGT, insulin resistance syndrome, syndrome X, type 2 diabetes, type 1 diabetes, dyslipidemia, hypertension, and obesity.

Embodiment A302. A compound according to any one of embodiments A1 to A301 for use as a medicament.

Embodiment A303. A compound according to any one of embodiments A1 to A301 for treatment of hyperglycemia, for treatment of IGT, for treatment of Syndrome X, for treatment of type 2 diabetes, for treatment of type 1 diabetes, for treatment of dyslipidemia, for treatment of hyperlipidemia, for treatment of hypertension, for treatment of obesity, for lowering of food intake, for appetite regulation, for regulating feeding behaviour, or for enhancing the secretion of enteroincretins, such as GLP-1.

Embodiment A304. A pharmaceutical composition comprising, as an active ingredient, at least one compound according to any one of embodiments A1 to 303 together with one or more pharmaceutically acceptable carriers or excipients.

10 Embodiment A305. A pharmaceutical composition according to embodiment A304 in unit dosage form, comprising from about 0.05 mg to about 1000 mg, preferably from about 0.1 mg to about 500 mg and especially preferred from about 0.5 mg to about 200 mg of the compound according to any one of embodiments A1 to 303.

15 Embodiment A306. Use of a compound according to any one of the embodiments A1 to 303 for increasing the activity of glucokinase.

Embodiment A307. Use of a compound according to any one of embodiments A1 to A303 for the preparation of a medicament for the treatment of metabolic disorders, for blood glucose lowering, for the treatment of hyperglycemia, for the treatment of IGT, for the treatment of Syndrome X, for the treatment of impaired fasting glucose (IFG), for the treatment of type 2 diabetes, for the treatment of type 1 diabetes, for delaying the progression of impaired glucose tolerance (IGT) to type 2 diabetes, for delaying the progression of non-insulin requiring type 2 diabetes to insulin requiring type 2 diabetes, for the treatment of dyslipidemia, for the treatment of hyperlipidemia, for the treatment of hypertension, for lowering of food intake, for appetite regulation, for the treatment of obesity, for regulating feeding behaviour, or for enhancing the secretion of enteroincretins.

Embodiment A308. Use of a compound according to any one of embodiments A1 to A303 for the preparation of a medicament for the adjuvant treatment of type 1 diabetes for preventing the onset of diabetic complications.

Embodiment A309. Use of a compound according to any one of embodiments A1 to A303 for the preparation of a medicament for increasing the number and/or the size of beta cells in a mammalian subject, for treatment of beta cell degeneration, in particular apoptosis of beta cells, or for treatment of functional dyspepsia, in particular irritable bowel syndrome.

- 5 Embodiment A310. Use according to any one of the embodiments A307 to A309 in a regimen which comprises treatment with a further antidiabetic agent.

Embodiment A311. Use according to any one of the embodiments A307 to A310 in a regimen which comprises treatment with a further antihyperlipidemic agent.

- 10 Embodiment A312. Use according to any one of embodiments A307 to A311 in a regimen which comprises treatment with a further antiobesity agent.

Embodiment A313. Use according to any one of embodiments A307 to A312 in a regimen which comprises treatment with a further antihypertensive agent.

- 15 Embodiment A314. Use of a compound according to any one of the embodiments A1 to A303 or a pharmaceutical composition according to embodiment A304 or embodiment A305 for the treatment of metabolic disorders, for blood glucose lowering, for the treatment of hyperglycemia, for treatment of IGT, for treatment of Syndrome X, for the treatment of impaired fasting glucose (IFG), for treatment of type 2 diabetes, for treatment of type 1 diabetes, for delaying the progression of impaired glucose tolerance (IGT) to type 2 diabetes, for delaying the progression of non-insulin requiring type 2 diabetes to insulin requiring type 2
20 diabetes, for treatment of dyslipidemia, for treatment of hyperlipidemia, for treatment of hypertension, for the treatment or prophylaxis of obesity, for lowering of food intake, for appetite regulation, for regulating feeding behaviour, or for enhancing the secretion of enteroinsulins.

- 25 Embodiment A315. Use of a compound according to any one of the embodiments A1 to A303 or a pharmaceutical composition according to embodiment A304 or embodiment A305 for the adjuvant treatment of type 1 diabetes for preventing the onset of diabetic complications.

Embodiment A316. Use of a compound according to any one of the embodiments A1 to A303 or a pharmaceutical composition according to embodiment A304 or embodiment A305

for increasing the number and/or the size of beta cells in a mammalian subject, for treatment of beta cell degeneration, in particular apoptosis of beta cells, or for treatment of functional dyspepsia, in particular irritable bowel syndrome.

Further embodiments are clear from the appended claims.

5 Included within the scope of the present invention are the individual enantiomers of the compounds represented by formula (I) above as well as any wholly or partially racemic mixtures thereof. The present invention also covers the individual enantiomers of the compounds represented by formula (I) above as mixtures with diastereoisomers thereof in which one or more stereocenters are inverted.

10 The present invention provides glucose sensitive glucokinase activators, that is glucokinase activators, for instance of the general formula (I), which provides a higher increase in glucokinase activity at lower concentrations of glucose. This should be taken to mean that when the glucose concentration is low, then the glucose sensitive glucokinase activator provides an increase in the glucokinase activity, which increase is higher than the
15 increase in glucokinase activity provided by the compound when glucose concentration is high. The compound may for instance provide a 4.0 fold increase in glucokinase activity at a glucose concentration of 5 mM and a 2.0 fold increase in glucokinase activity at a glucose concentration of 15 mM, thus providing an increase in glucokinase activity at a glucose concentration of 5 mM, which increase is 2.0 fold higher than the increase in glucokinase
20 activity provided by the compound at a glucose concentration of 15 mM. For the purpose of describing the present invention, the glucose sensitivity may be assayed by use of Glucokinase Activity Assay (I) where the activity of the glucokinase activator is measured at different concentrations of glucose.

The glucose sensitivity of a glucokinase activator may for instance be measured at a
25 glucose concentration of 5 mM and at a glucose concentration of 15 mM using the same concentration of glucokinase activator, such as a concentration of 10 μ M. The two measurements may then be compared and if the fold activity at a glucose concentration of 5 mM (the lower glucose concentration) - in the above example 4.0 fold - is significantly higher than the fold activity at a glucose concentration of 15 mM (the higher glucose concentration -
30 in the above example 2.0 fold - then the glucokinase activator is deemed to be a glucose sensitive glucokinase activator. In the above example, the increase in glucokinase activity at a glucose concentration of 5 mM is 2.0 fold higher than the increase in glucokinase activity at a glucose concentration of 15 mM. The increase in glucokinase activity provided by the glucokinase activator at 5 mM glucose may for instance be at least 1.1 fold higher, such as at
35 least 1.2 fold higher, for instance at least 1.3 fold higher, such as at least 1.4 fold higher, for

instance 1.5 fold higher, such as at least 1.6 fold higher, for instance at least 1.7 fold higher, such as at least 1.8 fold higher, for instance at least 1.9 fold higher, such as at least 2.0 fold higher than the activity of the glucokinase activator at 15 mM glucose.

The present invention provides liver specific glucokinase activators, that is, glucokinase activators, for instance of the general formula (I), which increase glucose utilization in the liver (i.e. increase glycogen deposition) without inducing any increase in insulin secretion in response to glucose. For the purpose of describing this invention, the potential liver selectivity of a glucokinase activator may be assayed by comparison of the results obtained in response to the glucokinase activator in isolated hepatocytes and the results obtained in response to the glucokinase activator in Ins-1 cells. Glucokinase activators, which show a significantly higher activity in isolated hepatocytes measured as described in the Glucokinase Activity Assay (II) compared to the activity in Ins-1 cells measured as described in the Glucokinase Activity Assay (III), are deemed to be liver specific glucokinase activators. The activity of the glucokinase activator in Glucokinase Activity Assay (II) (hepatocytes) may for instance be at least 1.1 fold higher, such as at least 1.2 fold higher, for instance at least 1.3 fold higher, such as at least 1.4 fold higher, for instance 1.5 fold higher, such as at least 1.6 fold higher, for instance at least 1.7 fold higher, such as at least 1.8 fold higher, for instance at least 1.9 fold higher, such as at least 2.0 fold higher, for instance at least a 3.0 fold higher, such as at least a 4.0 fold higher, for instance at least 5.0 fold higher, such as at least 10 fold higher than the activity of the glucokinase activator in Glucokinase Activity Assay (III) (Ins-1 cells). Alternatively, the glucokinase activator may show no activity in the Ins-1 cells measured as described in the Glucokinase Activity Assay (III), while showing a significant activity in hepatocytes measured as described in the Glucokinase Activity Assay (II).

Such liver-specific glucokinase activators may be particularly useful in patients that are at risk of experiencing hypoglycaemia. Since liver glucokinase is highly sensitive to the serum concentration of glucose, the blood glucose-decreasing effect of the GK in the liver will only occur when the serum concentration of glucose is relatively high. When the serum concentration of glucose is relatively low, the effect of the GK in the liver decreases, and thus does not further lower the glucose concentration in the blood. This mechanism remains even when the liver GK is affected by a GK activator. The effect of GK on the pancreatic beta cells is not similarly glucose-sensitive. Therefore a GK activator which affects both liver and beta cells may have a glucose-lowering effect even at low serum glucose concentration, resulting in a risk of hypoglycaemia. A GK activator which affects only, or which primarily effects, the liver GK will thus provide a treatment with a lower risk of hypoglycaemia. Thus the invention

provides a method of preventing hypoglycaemia comprising administration of a liver-specific glucokinase activator, as well as the use of a liver-specific glucokinase activator for the preparation of a medicament for the prevention of hypoglycaemia.

Examples of liver specific glucokinase activators are

- 5 2-{3-[2-(2,3-dimethoxyphenoxy)-5-fluorophenyl]ureido}thiazole-4-carboxylic acid ethyl ester, (2-{3-[2-(2,3-dimethoxyphenoxy)-5-fluorophenyl]ureido}thiazol-4-yl)acetic acid ethyl ester, (2-{3-[5-fluoro-2-(2-fluoro-6-methoxyphenoxy)phenyl]ureido}thiazol-4-yl)acetic acid, 2-{3-[2-(2,3-dimethoxyphenoxy)-5-fluorophenyl]ureido}thiazole-4-carboxylic acid, 2-(2-{3-[5-fluoro-2-(2-fluoro-6-methoxyphenoxy)phenyl]ureido}thiazol-4-yl)-N-(2-morpholin-4-yl)ethyl)acetamide,
- 10 [2-(2-{3-[5-fluoro-2-(2-fluoro-6-methoxyphenoxy)phenyl]ureido}thiazol-4-yl)acetylamino]acetic acid, {2-[3-(2-cyclopentanecarbonyl-4-methyl-phenyl)-ureido]-thiazol-4-yl}acetic acid, and {2-[3-(4-methyl-2-[2-methylpropoxy]phenyl)-ureido]-thiazol-4-yl}-acetic acid.

- 15 In one embodiment, a compound according to the present invention is for use as a medicament in the treatment of hyperglycemia.

In one embodiment, a compound according to the present invention is for use as a medicament in the treatment of IGT.

- In one embodiment, a compound according to the present invention is for use as a medicament in the treatment of Syndrome X.
- 20

In one embodiment, a compound according to the present invention is for use as a medicament in the treatment of type 2 diabetes.

In one embodiment, a compound according to the present invention is for use as a medicament in the treatment of type 1 diabetes.

- 25 In one embodiment, a compound according to the present invention is for use as a medicament in the treatment of hyperlipidemia.

In one embodiment, a compound according to the present invention is for use as a medicament in the treatment of dyslipidemia.

- In one embodiment, a compound according to the present invention is for use as a medicament in the treatment of hypertension.
- 30

In one embodiment, a compound according to the present invention is for use as a medicament in the treatment of obesity.

In one embodiment, a compound according to the present invention is for use as a medicament for lowering of food intake.

In one embodiment, a compound according to the present invention is for use as a medicament for appetite regulation.

In one embodiment, a compound according to the present invention is for use as a medicament for regulating feeding behaviour.

5 In one embodiment, a compound according to the present invention is for use as a medicament for enhancing the secretion of enteroinsulin, such as GLP-1.

The present compounds are activators of glucokinase and are as such useful for the activation of glucokinase.

10 Accordingly, the present invention provides a method for activating glucokinase in a patient in need thereof, which method comprises administering to a subject in need thereof a compound according to the present invention, preferably in a pharmacologically effective amount, more preferably in a therapeutically effective amount. The present invention also provides a method for lowering blood glucose in a patient in need thereof, which method comprises administering to a subject in need thereof a compound according to the present
15 invention, preferably in a pharmacologically effective amount, more preferably in a therapeutically effective amount. The present invention also provides a method for prevention and/or treatment of glucokinase deficiency-mediated human diseases, the method comprising administration to a human in need thereof a therapeutically effective amount of a compound according to the present invention. As used herein, the phrase "a
20 subject in need thereof" includes mammalian subjects, preferably humans, who either suffer from one or more of the aforesaid diseases or disease states or are at risk for such. Accordingly, in the context of the therapeutic method of the present invention, this method also is comprised of a method for treating a mammalian subject prophylactically, or prior to the onset of diagnosis such disease(s) or disease state(s).

25 In one embodiment, the present invention provides a method for the treatment of hyperglycemia, the method comprising administering to a subject in need thereof an effective amount of a compound or a pharmaceutical composition according to the present invention.

In one embodiment, the present invention provides a method for the treatment of IGT, the method comprising administering to a subject in need thereof an effective amount of
30 a compound or a pharmaceutical composition according to the present invention.

In one embodiment, the present invention provides a method for the treatment of Syndrome X, the method comprising administering to a subject in need thereof an effective amount of a compound or a pharmaceutical composition according to the present invention.

In one embodiment, the present invention provides a method for the treatment of
35 type 2 diabetes, the method comprising administering to a subject in need thereof an

effective amount of a compound or a pharmaceutical composition according to the present invention.

In one embodiment, the present invention provides a method for the treatment of type 1 diabetes, the method comprising administering to a subject in need thereof an effective amount of a compound or a pharmaceutical composition according to the present invention.

In one embodiment, the present invention provides a method for the treatment of dyslipidemia or hyperlipidemia, the method comprising administering to a subject in need thereof an effective amount of a compound or a pharmaceutical composition according to the present invention.

In one embodiment, the present invention provides a method for the treatment of obesity, the method comprising administering to a subject in need thereof an effective amount of a compound or a pharmaceutical composition according to the present invention.

In one embodiment, the effective amount of the compound is in the range of from about 0.05 mg to about 2000 mg, preferably from about 0.1 mg to about 1000 mg and especially preferred from about 0.5 mg to about 500 mg per day.

In one embodiment, the method according to the present invention is part of a regimen, which comprises treatment with a further antidiabetic agent, for example an antidiabetic agent such as insulin or an insulin analogue, a sulphonylurea, a biguanide, a meglitinide, an insulin sensitizer, a thiazolidinedione insulin sensitizer, an α -glucosidase inhibitor, a glycogen phosphorylase inhibitor, or an agent acting on the ATP-dependent potassium channel of the pancreatic β -cells.

In one embodiment, the method according to the present invention is part of a regimen, which comprises treatment with a further antihyperlipidemic agent, for example an antihyperlipidemic agent such as cholestyramine, colestipol, clofibrate, gemfibrozil, lovastatin, pravastatin, simvastatin, probucol or dextrothyroxine.

In one embodiment, the method according to the present invention is part of a regimen, which comprises treatment with a further antihypertensive agent.

In one embodiment, the method according to the present invention is part of a regimen, which comprises treatment with a further antiobesity or appetite regulating agent.

In one embodiment, the method according to the present invention is part of a regimen, which comprises treatment with a further antihypertensive agent.

Other embodiments of such methods will be clear from the following description.

Compounds according to the present invention are useful for the treatment of disorders, diseases and conditions, wherein the activation of glucokinase is beneficial.

Accordingly, the present compounds are useful for the treatment of hyperglycemia, IGT (impaired glucose tolerance), insulin resistance syndrome, syndrome X, type 1 diabetes, type 2 diabetes, dyslipidemia, dyslipoproteinemia (abnormal lipoproteins in the blood) including diabetic dyslipidemia, hyperlipidemia, hyperlipoproteinemia (excess of lipoproteins in the blood) including type I, II-a (hypercholesterolemia), II-b, III, IV (hypertriglyceridemia) and V (hypertriglyceridemia) hyperlipoproteinemias, and obesity. Furthermore, they may be useful for the treatment of albuminuria, cardiovascular diseases such as cardiac hypertrophy, hypertension and arteriosclerosis including atherosclerosis; gastrointestinal disorders; acute pancreatitis; and appetite regulation or energy expenditure disorders.

Accordingly, in a further aspect the invention relates to a compound according to the present invention for use as a medicament.

The invention also relates to pharmaceutical compositions comprising, as an active ingredient, at least one compound according to the present invention together with one or more pharmaceutically acceptable carriers or excipients.

The pharmaceutical composition is preferably in unit dosage form, comprising from about 0.05 mg to about 1000 mg, preferably from about 0.1 mg to about 500 mg and especially preferred from about 0.5 mg to about 200 mg of the compound according to the present invention.

In one embodiment, the pharmaceutical composition according to the present invention comprises a further antidiabetic agent, for example an antidiabetic agent such as insulin, an insulin derivative or an insulin analogue, a sulphonylurea, a biguanide, a meglitinide, an insulin sensitizer, a thiazolidinedione insulin sensitizer, an α -glucosidase inhibitor, a glycogen phosphorylase inhibitor, or an agent acting on the ATP-dependent potassium channel of the pancreatic β -cells.

In one embodiment, the pharmaceutical composition according to the present invention comprises a further antihyperlipidemic agent, for example an antihyperlipidemic agent such as cholestyramine, colestipol, clofibrate, gemfibrozil, lovastatin, pravastatin, simvastatin, probucol or dextrothyroxine.

In one embodiment, the pharmaceutical composition according to the present invention comprises a further antiobesity or appetite regulating agent.

In one embodiment, the pharmaceutical composition according to the present invention comprises a further antihypertensive agent.

In one embodiment of the present invention, the pharmaceutical composition according to the present invention comprises a compound according to the present invention in combination with one or more of the agents mentioned above eg in combination with

metformin and a sulphonylurea such as glyburide; a sulphonylurea and acarbose; nateglinide and metformin; acarbose and metformin; a sulphonylurea, metformin and troglitazone; insulin and a sulphonylurea; insulin and metformin; insulin, metformin and a sulphonylurea; insulin and troglitazone; insulin and lovastatin; etc.

5 In one embodiment of the present invention, the present compounds are used for the preparation of a medicament for the treatment of hyperglycemia. As used herein hyperglycemia is to be taken as generally understood in the art, with reference for example to the Report of the Expert Committee of the Diagnosis and Classification of Diabetes Mellitus, published in *Diabetes Care* 20, 1183-1197, (1997), but is usually taken to mean an
10 elevated plasma glucose level exceeding about 110 mg/dl. The present compounds are effective in lowering the blood glucose both in the fasting and postprandial stage.

 In one embodiment of the present invention, the present compounds are used for the preparation of a pharmaceutical composition for the treatment of IGT.

 In one embodiment of the present invention, the present compounds are used for
15 the preparation of a pharmaceutical composition for the treatment of Syndrome X.

 In one embodiment of the present invention, the present compounds are used for the preparation of a pharmaceutical composition for the treatment of type 2 diabetes. Such treatment includes a treatment for the purpose of the delaying of the progression from IGT to type 2 diabetes as well as delaying the progression from non-insulin requiring type 2
20 diabetes to insulin requiring type 2 diabetes.

 In one embodiment of the present invention the present compounds are used for the preparation of a pharmaceutical composition for the treatment of type 1 diabetes. Such therapy is normally accompanied by insulin administration.

 In one embodiment of the present invention the present compounds are used for the
25 preparation of a pharmaceutical composition for the treatment of dyslipidemia and hyperlipidemia.

 In one embodiment of the present invention the present compounds are used for the preparation of a pharmaceutical composition for the treatment of obesity.

 In another aspect of the present invention treatment of a patient with the present
30 compounds are combined with diet and/or exercise.

 The present invention provides methods of activating glucokinase activity in a mammal, which methods comprise administering, to a mammal in need of activation of glucokinase activity, a therapeutically defined amount of a compound according to the present invention defined above, as a single or polymorphic crystalline form or forms, an
35 amorphous form, a single enantiomer, a racemic mixture, a single stereoisomer, a mixture of

stereoisomers, a single diastereoisomer, a mixture of diastereoisomers, a solvate, a pharmaceutically acceptable salt, a solvate, a prodrug, a biohydrolyzable ester, or a biohydrolyzable amide thereof.

The present invention provides a method of activating glucokinase, comprising the
5 step of administering to a mammal in need thereof a pharmacologically effective amount of a compound according to the present invention. The invention further provides a pharmaceutical composition comprising a pharmaceutically acceptable carrier and a pharmacologically effective amount of a compound according to the present invention sufficient to activate glucokinase. A glucokinase - activating amount can be an amount that
10 reduces or inhibits a PTPase activity in the subject.

Additionally provided by the present invention is a pharmaceutical composition comprising a pharmaceutically acceptable carrier and a pharmacologically effective amount of a compound according to the present invention sufficient to treat type I diabetes.

Also provided by the present invention is a pharmaceutical composition comprising
15 a pharmaceutically acceptable carrier and a pharmacologically effective amount of a compound according to the present invention sufficient to treat type II diabetes.

The compounds of the present invention can be administered to any mammal in need of activation of glucokinase activity. Such mammals can include, for example, horses, cows, sheep, pigs, mice, dogs, cats, primates such as chimpanzees, gorillas, rhesus
20 monkeys, and, most preferably humans.

In a further aspect of the present invention the present compounds are administered in combination with one or more further active substances in any suitable ratios. Such further active agents may be selected from antidiabetic agents, antihyperlipidemic agents, antiobesity agents, antihypertensive agents and agents for the treatment of complications
25 resulting from or associated with diabetes.

Suitable antidiabetic agents include insulin, GLP-1 (glucagon like peptide-1) derivatives such as those disclosed in WO 98/08871 (Novo Nordisk A/S), which is incorporated herein by reference, as well as orally active hypoglycemic agents.

Suitable orally active hypoglycemic agents preferably include imidazolines,
30 sulfonylureas, biguanides, meglitinides, oxadiazolidinediones, thiazolidinediones, insulin sensitizers, α -glucosidase inhibitors, agents acting on the ATP-dependent potassium channel of the pancreatic β -cells eg potassium channel openers such as those disclosed in WO 97/26265, WO 99/03861 and WO 00/37474 (Novo Nordisk A/S) which are incorporated herein by reference, potassium channel openers, such as ornitiglinide, potassium channel
35 blockers such as nateglinide or BTS-67582, glucagon antagonists such as those disclosed in

WO 99/01423 and WO 00/39088 (Novo Nordisk A/S and Agouron Pharmaceuticals, Inc.), all of which are incorporated herein by reference, GLP-1 agonists such as those disclosed in WO 00/42026 (Novo Nordisk A/S and Agouron Pharmaceuticals, Inc.), which are incorporated herein by reference, DPP-IV (dipeptidyl peptidase-IV) inhibitors, PTPase
5 (protein tyrosine phosphatase) inhibitors, inhibitors of hepatic enzymes involved in stimulation of gluconeogenesis and/or glycogenolysis, glucose uptake modulators, GSK-3 (glycogen synthase kinase-3) inhibitors, compounds modifying the lipid metabolism such as antihyperlipidemic agents and antilipidemic agents, compounds lowering food intake, and PPAR (peroxisome proliferator-activated receptor) and RXR (retinoid X receptor) agonists
10 such as ALRT-268, LG-1268 or LG-1069.

In one embodiment of the present invention, the present compounds are administered in combination with insulin, insulin derivatives or insulin analogues.

In one embodiment of the present invention, the present compounds are administered in combination with a sulphonylurea eg tolbutamide, chlorpropamide,
15 tolazamide, glibenclamide, glipizide, glimepiride, glicazide or glyburide.

In one embodiment of the present invention, the present compounds are administered in combination with a biguanide eg metformin.

In one embodiment of the present invention, the present compounds are administered in combination with a meglitinide eg repaglinide or senaglinide/nateglinide.

20 In one embodiment of the present invention, the present compounds are administered in combination with a thiazolidinedione insulin sensitizer eg troglitazone, ciglitazone, pioglitazone, rosiglitazone, isaglitazone, darglitazone, englitazone, CS-011/CI-1037 or T 174 or the compounds disclosed in WO 97/41097 (DRF-2344), WO 97/41119, WO 97/41120, WO 00/41121 and WO 98/45292 (Dr. Reddy's Research Foundation), which are
25 incorporated herein by reference.

In one embodiment of the present invention the present compounds may be administered in combination with an insulin sensitizer eg such as GI 262570, YM-440, MCC-555, JTT-501, AR-H039242, KRP-297, GW-409544, CRE-16336, AR-H049020, LY510929, MBX-102, CLX-0940, GW-501516 or the compounds disclosed in WO 99/19313
30 (NN622/DRF-2725), WO 00/50414, WO 00/63191, WO 00/63192, WO 00/63193 (Dr. Reddy's Research Foundation) and WO 00/23425, WO 00/23415, WO 00/23451, WO 00/23445, WO 00/23417, WO 00/23416, WO 00/63153, WO 00/63196, WO 00/63209, WO 00/63190 and WO 00/63189 (Novo Nordisk A/S), which are incorporated herein by reference.

In one embodiment of the present invention the present compounds are administered in combination with an α -glucosidase inhibitor eg voglibose, emiglitate, miglitol or acarbose.

In one embodiment of the present invention the present compounds are
5 administered in combination with a glycogen phosphorylase inhibitor eg the compounds described in WO 97/09040 (Novo Nordisk A/S).

In one embodiment of the present invention the present compounds are administered in combination with an agent acting on the ATP-dependent potassium channel of the pancreatic β -cells eg tolbutamide, glibenclamide, glipizide, glicazide, BTS-67582 or
10 repaglinide.

In one embodiment of the present invention the present compounds are administered in combination with nateglinide.

In one embodiment of the present invention the present compounds are administered in combination with an antihyperlipidemic agent or a antilipidemic agent eg
15 cholestyramine, colestipol, clofibrate, gemfibrozil, lovastatin, pravastatin, simvastatin, probucol or dextrothyroxine.

In another aspect of the present invention, the present compounds are administered in combination with more than one of the above-mentioned compounds eg in combination with metformin and a sulphonylurea such as glyburide; a sulphonylurea and acarbose;
20 nateglinide and metformin; acarbose and metformin; a sulphonylurea, metformin and troglitazone; insulin and a sulphonylurea; insulin and metformin; insulin, metformin and a sulphonylurea; insulin and troglitazone; insulin and lovastatin; etc.

Furthermore, the compounds according to the invention may be administered in combination with one or more antiobesity agents or appetite regulating agents.

25 Such agents may be selected from the group consisting of CART (cocaine amphetamine regulated transcript) agonists, NPY (neuropeptide Y) antagonists, MC3 (melanocortin 3) agonists, MC4 (melanocortin 4) agonists, orexin antagonists, TNF (tumor necrosis factor) agonists, CRF (corticotropin releasing factor) agonists, CRF BP (corticotropin releasing factor binding protein) antagonists, urocortin agonists, β 3 adrenergic
30 agonists such as CL-316243, AJ-9677, GW-0604, LY362884, LY377267 or AZ-40140, MSH (melanocyte-stimulating hormone) agonists, MCH (melanocyte-concentrating hormone) antagonists, CCK (cholecystokinin) agonists, serotonin reuptake inhibitors (fluoxetine, seroxat or citalopram), serotonin and norepinephrine reuptake inhibitors, 5HT (serotonin) agonists, bombesin agonists, galanin antagonists, growth hormone, growth factors such as
35 prolactin or placental lactogen, growth hormone releasing compounds, TRH (thyreotropin

releasing hormone) agonists, UCP 2 or 3 (uncoupling protein 2 or 3) modulators, leptin agonists, DA (dopamine) agonists (bromocriptin, dorexin), lipase/amylase inhibitors, PPAR modulators, RXR modulators, TR β agonists, adrenergic CNS stimulating agents, AGRP (agouti related protein) inhibitors, H3 histamine antagonists such as those disclosed in WO 00/42023, WO 00/63208 and WO 00/64884, which are incorporated herein by reference, 5 exendin-4, GLP-1 agonists and ciliary neurotrophic factor. Further antiobesity agents are bupropion (antidepressant), topiramate (anticonvulsant), ecopipam (dopamine D1/D5 antagonist) and naltrexone (opioid antagonist).

In one embodiment of the present invention the antiobesity agent is leptin.

10 In one embodiment of the present invention the antiobesity agent is a serotonin and norepinephrine reuptake inhibitor eg sibutramine.

In one embodiment of the present invention the antiobesity agent is a lipase inhibitor eg orlistat.

In one embodiment of the present invention the antiobesity agent is an adrenergic 15 CNS stimulating agent eg dexamphetamine, amphetamine, phentermine, mazindol phendimetrazine, diethylpropion, fenfluramine or dexfenfluramine.

Furthermore, the present compounds may be administered in combination with one or more antihypertensive agents. Examples of antihypertensive agents are β -blockers such as alprenolol, atenolol, timolol, pindolol, propranolol and metoprolol, ACE (angiotensin 20 converting enzyme) inhibitors such as benazepril, captopril, enalapril, fosinopril, lisinopril, quinapril and ramipril, calcium channel blockers such as nifedipine, felodipine, nicardipine, isradipine, nimodipine, diltiazem and verapamil, and α -blockers such as doxazosin, urapidil, prazosin and terazosin. Further reference can be made to Remington: The Science and Practice of Pharmacy, 19th Edition, Gennaro, Ed., Mack Publishing Co., Easton, PA, 1995.

25 In a further aspect the invention provides a pharmaceutical preparation comprising an activator of glucokinase and an insulin derivative.

In one embodiment of the invention the insulin derivative is selected from the group consisting of B29-N^E-myristoyl-des(B30) human insulin, B29-N^E-palmitoyl-des(B30) human insulin, B29-N^E-myristoyl human insulin, B29-N^E-palmitoyl human insulin, B28-N^E-myristoyl 30 Lys^{B28} Pro^{B29} human insulin, B28-N^E-palmitoyl Lys^{B28} Pro^{B29} human insulin, B30-N^E-myristoyl-Thr^{B29} Lys^{B30} human insulin, B30-N^E-palmitoyl-Thr^{B29} Lys^{B30} human insulin, B29-N^E-(N-palmitoyl- γ -glutamyl)-des(B30) human insulin, B29-N^E-(N-lithocholyl- γ -glutamyl)-des(B30) human insulin, B29-N^E-(ω -carboxyheptadecanoyl)-des(B30) human insulin and B29-N^E-(ω -carboxyheptadecanoyl) human insulin.

In another embodiment of the invention the insulin derivative is B29-N^ε-myristoyl-des(B30) human insulin.

It should be understood that any suitable combination of the compounds according to the invention with diet and/or exercise, one or more of the above-mentioned compounds
5 and optionally one or more other active substances are considered to be within the scope of the present invention.

PHARMACEUTICAL COMPOSITIONS

The compounds of the present invention may be administered alone or in combination with pharmaceutically acceptable carriers or excipients, in either single or
10 multiple doses. The pharmaceutical compositions according to the invention may be formulated with pharmaceutically acceptable carriers or diluents as well as any other known adjuvants and excipients in accordance with conventional techniques such as those disclosed in Remington: The Science and Practice of Pharmacy, 19th Edition, Gennaro, Ed., Mack Publishing Co., Easton, PA, 1995.

15 The pharmaceutical compositions may be specifically formulated for administration by any suitable route such as the oral, rectal, nasal, pulmonary, topical (including buccal and sublingual), transdermal, intracisternal, intraperitoneal, vaginal and parenteral (including subcutaneous, intramuscular, intrathecal, intravenous and intradermal) route, the oral route being preferred. It will be appreciated that the preferred route will depend on the general
20 condition and age of the subject to be treated, the nature of the condition to be treated and the active ingredient chosen.

Pharmaceutical compositions for oral administration include solid dosage forms such as hard or soft capsules, tablets, troches, dragees, pills, lozenges, powders and granules. Where appropriate, they can be prepared with coatings such as enteric coatings or
25 they can be formulated so as to provide controlled release of the active ingredient such as sustained or prolonged release according to methods well known in the art.

Liquid dosage forms for oral administration include solutions, emulsions, aqueous or oily suspensions, syrups and elixirs.

Pharmaceutical compositions for parenteral administration include sterile aqueous
30 and non-aqueous injectable solutions, dispersions, suspensions or emulsions as well as sterile powders to be reconstituted in sterile injectable solutions or dispersions prior to use. Depot injectable formulations are also contemplated as being within the scope of the present invention.

Other suitable administration forms include suppositories, sprays, ointments, cremes, gels, inhalants, dermal patches, implants etc.

A typical oral dosage is in the range of from about 0.001 to about 100 mg/kg body weight per day, preferably from about 0.01 to about 50 mg/kg body weight per day, and more preferred from about 0.05 to about 10 mg/kg body weight per day administered in one or more dosages such as 1 to 3 dosages. The exact dosage will depend upon the frequency and mode of administration, the sex, age, weight and general condition of the subject treated, the nature and severity of the condition treated and any concomitant diseases to be treated and other factors evident to those skilled in the art.

The formulations may conveniently be presented in unit dosage form by methods known to those skilled in the art. A typical unit dosage form for oral administration one or more times per day such as 1 to 3 times per day may contain from 0.05 to about 1000 mg, preferably from about 0.1 to about 500 mg, and more preferred from about 0.5 mg to about 200 mg.

For parenteral routes such as intravenous, intrathecal, intramuscular and similar administration, typically doses are in the order of about half the dose employed for oral administration.

The compounds of this invention are generally utilized as the free substance or as a pharmaceutically acceptable salt thereof. Examples are an acid addition salt of a compound having the utility of a free base and a base addition salt of a compound having the utility of a free acid. The term "pharmaceutically acceptable salts" refers to non-toxic salts of the compounds of this invention which are generally prepared by reacting the free base with a suitable organic or inorganic acid or by reacting the acid with a suitable organic or inorganic base. When a compound according to the present invention contains a free base such salts are prepared in a conventional manner by treating a solution or suspension of the compound with a chemical equivalent of a pharmaceutically acceptable acid. When a compound according to the present invention contains a free acid such salts are prepared in a conventional manner by treating a solution or suspension of the compound with a chemical equivalent of a pharmaceutically acceptable base. Physiologically acceptable salts of a compound with a hydroxy group include the anion of said compound in combination with a suitable cation such as sodium or ammonium ion. Other salts which are not pharmaceutically acceptable may be useful in the preparation of compounds of the present invention and these form a further aspect of the present invention.

For parenteral administration, solutions of the novel compounds of the formula (I) in sterile aqueous solution, aqueous propylene glycol or sesame or peanut oil may be

employed. Such aqueous solutions should be suitably buffered if necessary and the liquid diluent first rendered isotonic with sufficient saline or glucose. The aqueous solutions are particularly suitable for intravenous, intramuscular, subcutaneous and intraperitoneal administration. The sterile aqueous media employed are all readily available by standard techniques known to those skilled in the art.

Suitable pharmaceutical carriers include inert solid diluents or fillers, sterile aqueous solution and various organic solvents. Examples of solid carriers are lactose, terra alba, sucrose, cyclodextrin, talc, gelatine, agar, pectin, acacia, magnesium stearate, stearic acid and lower alkyl ethers of cellulose. Examples of liquid carriers are syrup, peanut oil, olive oil, phospholipids, fatty acids, fatty acid amines, polyoxyethylene and water. Similarly, the carrier or diluent may include any sustained release material known in the art, such as glyceryl monostearate or glyceryl distearate, alone or mixed with a wax. The pharmaceutical compositions formed by combining the novel compounds of the present invention and the pharmaceutically acceptable carriers are then readily administered in a variety of dosage forms suitable for the disclosed routes of administration. The formulations may conveniently be presented in unit dosage form by methods known in the art of pharmacy.

Formulations of the present invention suitable for oral administration may be presented as discrete units such as capsules or tablets, each containing a predetermined amount of the active ingredient, and which may include a suitable excipient. Furthermore, the orally available formulations may be in the form of a powder or granules, a solution or suspension in an aqueous or non-aqueous liquid, or an oil-in-water or water-in-oil liquid emulsion.

Compositions intended for oral use may be prepared according to any known method, and such compositions may contain one or more agents selected from the group consisting of sweetening agents, flavoring agents, coloring agents, and preserving agents in order to provide pharmaceutically elegant and palatable preparations. Tablets may contain the active ingredient in admixture with non-toxic pharmaceutically-acceptable excipients which are suitable for the manufacture of tablets. These excipients may be for example, inert diluents, such as calcium carbonate, sodium carbonate, lactose, calcium phosphate or sodium phosphate; granulating and disintegrating agents, for example corn starch or alginic acid; binding agents, for example, starch, gelatin or acacia; and lubricating agents, for example magnesium stearate, stearic acid or talc. The tablets may be uncoated or they may be coated by known techniques to delay disintegration and absorption in the gastrointestinal tract and thereby provide a sustained action over a longer period. For example, a time delay material such as glyceryl monostearate or glyceryl distearate may be employed. They may

also be coated by the techniques described in U.S. Patent Nos. 4,356,108; 4,166,452; and 4,265,874, incorporated herein by reference, to form osmotic therapeutic tablets for controlled release.

Formulations for oral use may also be presented as hard gelatin capsules where the active ingredient is mixed with an inert solid diluent, for example, calcium carbonate, calcium phosphate or kaolin, or a soft gelatin capsules wherein the active ingredient is mixed with water or an oil medium, for example peanut oil, liquid paraffin, or olive oil.

Aqueous suspensions may contain the active compounds in admixture with excipients suitable for the manufacture of aqueous suspensions. Such excipients are suspending agents, for example sodium carboxymethylcellulose, methylcellulose, hydroxypropylmethylcellulose, sodium alginate, polyvinylpyrrolidone, gum tragacanth and gum acacia; dispersing or wetting agents may be a naturally-occurring phosphatide such as lecithin, or condensation products of an alkylene oxide with fatty acids, for example polyoxyethylene stearate, or condensation products of ethylene oxide with long chain aliphatic alcohols, for example, heptadecaethyl-eneoxycetanol, or condensation products of ethylene oxide with partial esters derived from fatty acids and a hexitol such as polyoxyethylene sorbitol monooleate, or condensation products of ethylene oxide with partial esters derived from fatty acids and hexitol anhydrides, for example polyethylene sorbitan monooleate. The aqueous suspensions may also contain one or more coloring agents, one or more flavoring agents, and one or more sweetening agents, such as sucrose or saccharin.

Oily suspensions may be formulated by suspending the active ingredient in a vegetable oil, for example arachis oil, olive oil, sesame oil or coconut oil, or in a mineral oil such as a liquid paraffin. The oily suspensions may contain a thickening agent, for example beeswax, hard paraffin or cetyl alcohol. Sweetening agents such as those set forth above, and flavoring agents may be added to provide a palatable oral preparation. These compositions may be preserved by the addition of an anti-oxidant such as ascorbic acid.

Dispersible powders and granules suitable for preparation of an aqueous suspension by the addition of water provide the active compound in admixture with a dispersing or wetting agent, suspending agent and one or more preservatives. Suitable dispersing or wetting agents and suspending agents are exemplified by those already mentioned above. Additional excipients, for example, sweetening, flavoring, and coloring agents may also be present.

The pharmaceutical compositions of the present invention may also be in the form of oil-in-water emulsions. The oily phase may be a vegetable oil, for example, olive oil or arachis oil, or a mineral oil, for example a liquid paraffin, or a mixture thereof. Suitable

emulsifying agents may be naturally-occurring gums, for example gum acacia or gum tragacanth, naturally-occurring phosphatides, for example soy bean, lecithin, and esters or partial esters derived from fatty acids and hexitol anhydrides, for example sorbitan monooleate, and condensation products of said partial esters with ethylene oxide, for example polyoxyethylene sorbitan monooleate. The emulsions may also contain sweetening and flavoring agents.

Syrups and elixirs may be formulated with sweetening agents, for example glycerol, propylene glycol, sorbitol or sucrose. Such formulations may also contain a demulcent, a preservative and flavoring and coloring agents. The pharmaceutical compositions may be in the form of a sterile injectible aqueous or oleaginous suspension. This suspension may be formulated according to the known methods using suitable dispersing or wetting agents and suspending agents described above. The sterile injectable preparation may also be a sterile injectable solution or suspension in a non-toxic parenterally-acceptable diluent or solvent, for example as a solution in 1,3-butanediol. Among the acceptable vehicles and solvents that may be employed are water, Ringer's solution, and isotonic sodium chloride solution. In addition, sterile, fixed oils are conveniently employed as solvent or suspending medium. For this purpose, any bland fixed oil may be employed using synthetic mono- or diglycerides. In addition, fatty acids such as oleic acid find use in the preparation of injectables.

The compositions may also be in the form of suppositories for rectal administration of the compounds of the present invention. These compositions can be prepared by mixing the drug with a suitable non-irritating excipient which is solid at ordinary temperatures but liquid at the rectal temperature and will thus melt in the rectum to release the drug. Such materials include cocoa butter and polyethylene glycols, for example.

For topical use, creams, ointments, jellies, solutions or suspensions, etc., containing the compounds of the present invention are contemplated. For the purpose of this application, topical applications shall include mouth washes and gargles.

The compounds of the present invention may also be administered in the form of liposome delivery systems, such as small unilamellar vesicles, large unilamellar vesicles, and multilamellar vesicles. Liposomes may be formed from a variety of phospholipids, such as cholesterol, stearylamine, or phosphatidylcholines.

In addition, some of the compounds of the present invention may form solvates with water or common organic solvents. Such solvates are also encompassed within the scope of the present invention.

Thus, in a further embodiment, there is provided a pharmaceutical composition comprising a compound according to the present invention, or a pharmaceutically acceptable

salt, solvate, or prodrug thereof, and one or more pharmaceutically acceptable carriers, excipients, or diluents.

If a solid carrier is used for oral administration, the preparation may be tableted, placed in a hard gelatine capsule in powder or pellet form or it can be in the form of a troche or lozenge. The amount of solid carrier will vary widely but will usually be from about 25 mg to about 1 g. If a liquid carrier is used, the preparation may be in the form of a syrup, emulsion, soft gelatine capsule or sterile injectable liquid such as an aqueous or non-aqueous liquid suspension or solution.

A typical tablet that may be prepared by conventional tableting techniques may contain:

Core:

| | |
|--|---------|
| Active compound (as free compound or salt thereof) | 5.0 mg |
| Lactosum Ph. Eur. | 67.8 mg |
| Cellulose, microcryst. (Avicel) | 31.4 mg |
| Amberlite®IRP88* | 1.0 mg |
| Magnesii stearas Ph. Eur. | q.s. |

Coating:

| | | |
|-------------------------------|---------|--------|
| Hydroxypropyl methylcellulose | approx. | 9 mg |
| Mywacett 9-40 T** | approx. | 0.9 mg |

* Polacrillin potassium NF, tablet disintegrant, Rohm and Haas.

** Acylated monoglyceride used as plasticizer for film coating.

If desired, the pharmaceutical composition of the present invention may comprise a compound according to the present invention in combination with further active substances such as those described in the foregoing.

The present invention also provides a method for the synthesis of compounds useful as intermediates in the preparation of compounds of formula (I) along with methods for the preparation of compounds of formula (I). The compounds can be prepared readily according to the following reaction Schemes (in which all variables are as defined before, unless so specified) using readily available starting materials, reagents and conventional synthesis procedures. In these reactions, it is also possible to make use of variants which are themselves known to those of ordinary skill in this art, but are not mentioned in greater detail.

ABBREVIATIONS

Abbreviations used in the Schemes and Examples are as follows:

| | | |
|----|----------------|--|
| | d | = days |
| | g | = grams |
| 5 | h | = hours |
| | Hz | = hertz |
| | kD | = kiloDalton |
| | L | = liters |
| | M | = molar |
| 10 | mbar | = millibar |
| | mg | = milligrams |
| | min | = minutes |
| | ml | = milliliters |
| | mM | = millimolar |
| 15 | mmol | = millimoles |
| | mol | = moles |
| | N | = normal |
| | ppm | = parts per million |
| | psi | = pounds per square inch |
| 20 | APCI | = atmospheric pressure chemical ionization |
| | ESI | = electrospray ionization |
| | i.v. | = intravenous |
| | m/z | = mass to charge ratio |
| | mp | = melting point |
| 25 | MS | = mass spectrometry |
| | NMR | = nuclear magnetic resonance spectroscopy |
| | p.o. | = per oral |
| | R _f | = relative TLC mobility |
| | rt | = room temperature |
| 30 | s.c. | = subcutaneous |
| | TLC | = thin layer chromatography |
| | t _r | = retention time |
| | BOP | = (1-benzotriazolyloxy)tris(dimethylamino)phosphonium hexafluorophosphate |
| 35 | DCM | = dichloromethane |


| | | |
|----|-------|--|
| | DIEA | = diisopropylethylamine |
| | DMF | = N, N-dimethylformamide |
| | DMPU | = 1,3-dimethypropylene urea |
| | DMSO | = dimethylsulfoxide |
| 5 | EDC | = 1-ethyl-3-(3-dimethylaminopropyl)-carbodiimide hydrochloride |
| | ether | = diethyl ether |
| | EtOAc | = ethyl acetate |
| | HMPA | = hexamethylphosphoric triamide |
| | HOBt | = 1-hydroxybenzotriazole |
| 10 | LAH | = lithium aluminum hydride |
| | LDA | = lithium diisopropylamide |
| | MeOH | = methanol |
| | NMM | = N-methylmorpholine, 4-methylmorpholine |
| | TEA | = triethylamine |
| 15 | TFA | = trifluoroacetic acid |
| | THF | = tetrahydrofuran |
| | THP | = tetrahydropyranyl |
| | TTF | = Fluoro-N,N,N'-tetramethylformamidinium hexafluorophosphate |

REACTION SCHEMES

20 Unless otherwise specified, the variables in the Schemes are as defined for formula (I).

Scheme 1 describes the preparation of compounds of formula (74).

Scheme 1



A^2 is heteroaryl, fused heterocycl/heteroaryl, or fused cycloalkyl/heteroaryl.


R^{100} and R^{101} , independently of each other, are substituents such as, but not limited to, H, alkyl, alkenyl, alkynyl, -alkylene-aryl, alkylene-cycloalkyl, and the like.

K is halogen or 1-imidazolyl.

The amine (71) may be treated with carbonyldiimidazole, 4-nitrophenyl chloroformate, phosgene or a derivative of phosgene such as diphosgene or triphosgene, in a solvent such as DCM or DCE. DMAP may be used as a catalyst in this reaction. The reaction may be conducted at a temperature of from 0°C to 100°C. The reaction mixture may be then be treated with the compound (73) and the whole may be incubated at a temperature of from 25°C to 100°C to afford the urea (75). It is also understood that (73) may be treated with the reagent (72) under similar conditions, followed by treatment with the amine (71), to afford (74).

Scheme 2 describes the preparation of a compound of formula (79).

Scheme 2



L¹¹ has the meaning of L¹ in formula (I), with the proviso that when L¹¹ is -D-alkylene-E-, -D-alkenylene-E-, -D-alkynylene-E-, -D-cycloalkylene-E-, or -D-heterocyclylene-E-, then D is selected from -O- or -S-, and that L¹¹ is not -S(O)-, -S(O)₂-, -C(O)-, or -C(=N-OR¹²)-.

Lg¹ is a leaving group such as F, Cl, Br, or I.


R¹⁰¹ is a substituent such as but not limited to H, alkyl, alkenyl, alkynyl, -alkylene-aryl, alkylene-cycloalkyl, and the like.

A nitro-substituted aryl or heteroaryl ring compound such as (75) may be treated with (76) in the presence of a base such as NaH or potassium tert-butoxide, in a solvent such as THF, DMF, or NMP at a temperature of from 0°C to 100°C, to afford (77). The resulting adduct (77) may be treated with tin(II) chloride in ethanol or other alcoholic solvent, at a temperature of from 25°C to 100°C, in the presence of aqueous HCl, to afford the amine (78). The amine (78) may, is desired, be treated with an alkyl halide R¹⁰¹-Lg², wherein Lg² is a leaving group such as Br, I, or p-toluenesulfonate, and a base such as DBU or sodium hydride, to afford (79). Alternatively, (78) may be treated with a reagent R¹⁰²-C(O)-R¹⁰³, wherein R¹⁰² and R¹⁰³ independently of each other are substituents such as, but not limited to, H, alkyl, alkenyl, alkynyl, -alkylene-aryl, alkylene-cycloalkyl, and the like, in the presence of a reducing agent such as sodium cyanoborohydride or sodium triacetoxyborohydride, to afford (79) wherein R¹⁰¹ should be understood as R¹⁰²-C(H)(R¹⁰³)-.

Alternatively, (78) may be treated with a reagent $R^{102}C(O)-OH$ in the presence of a dehydrating agent such as EDC, to afford an intermediate amide, which may be reduced with a reagent such as DIBAL or LAH, in a solvent such as THF, at a temperature of from $0^{\circ}C$ to $80^{\circ}C$, to afford (71) wherein R^{102} should be understood as $-CH_2-R^{102}$. Alternatively, (79) wherein R^{101} is $-CH_3$ may be prepared by treatment of (78) with a reagent $R^{102}-O-CO-Cl$, or $R^{102}O-CO-O-CO-O-R^{102}$, in the presence of a base such as TEA or aqueous alkali, to afford an intermediate which may be reduced as above employing DIBAL or LAH giving (79). Compound (79) may be employed in the same manner as is compound (73) according to the chemistry in Scheme (1).

Scheme 3 describes the synthesis of a compound of formula (71).

Scheme 3



A^2 is (un)substituted heteroaryl, (un)substituted fused heterocyclylheteroaryl, or (un)substituted fused cycloalkylheteroaryl.

R^{100} is a substituent such as, but not limited to, H, (un)substituted alkyl, (un)substituted alkenyl, (un)substituted alkynyl, (un)substituted -alkylene-aryl, (un)substituted -alkylene-cycloalkyl, and the like.


The amine (80) may, if desired, be treated with an alkyl halide $R^{100}-Lg^2$, wherein Lg^2 is a leaving group such as Br, I, or p-toluenesulfonate, and a base such as DBU or sodium hydride, to afford (71). Alternatively, (80) may be treated with a reagent $R^{102}-C(O)-R^{103}$ wherein R^{102} and R^{103} , independently of each other, are substituents such as, but not limited to, H, alkyl, alkenyl, alkynyl, -alkylene-aryl, -alkylene-cycloalkyl, and the like, in the presence of a reducing agent such as sodium cyanoborohydride or sodium triacetoxyborohydride, to afford (71) wherein R^{100} should be understood as $R^{102}-C(H)(R^{103})-$.

Alternatively, (78) may be treated with a reagent such as $R^{102}C(O)-Cl$ and a base such as TEA, or $R^{102}C(O)-OH$ in the presence of a dehydrating agent such as EDC, to afford an intermediate amide, which may be reduced with a reagent such as DIBAL or LAH, in a

solvent such as THF, at a temperature of from 0°C to 80°C, to afford (71) wherein R^{100} should be understood as $-\text{CH}_2-\text{R}^{102}$. Alternatively, (71) wherein R^{100} is $-\text{CH}_3$ may be prepared by treatment of (78) with a reagent $\text{R}^{102}-\text{O}-\text{C}(\text{O})-\text{Cl}$, or $\text{R}^{102}\text{O}-\text{C}(\text{O})-\text{O}-\text{C}(\text{O})-\text{O}-\text{R}^{102}$, in the presence of a base such as TEA or aqueous alkali, to afford an intermediate which may be reduced as above employing DIBAL or LAH giving (71).

Scheme 4 describes the synthesis of a compound of formula (81).

Scheme 4




A^2 is (un)substituted heteroaryl, (un)substituted fused heterocyclylheteroaryl, or (un)substituted fused cycloalkylheteroaryl.

R^{100} and R^{101} are, independently of each other, substituents such as but not limited to H, (un)substituted alkyl, (un)substituted alkenyl, (un)substituted alkynyl, (un)substituted -alkylene-aryl, (un)substituted -alkylene-cycloalkyl, and the like.

The amine (73) may be treated with the reagent chlorocarbonyl isocyanate in the presence of a base such as DIEA, in a solvent such as THF, DCE, or dioxane, at a temperature of from -60°C to 25°C. The intermediate thus formed may be treated at a temperature of from 0°C to 80°C with (71) to afford (81).

Scheme 5 describes the preparation of an intermediate of formula (87).

Scheme 5




L¹¹ is, in this instance, a group such as (un)substituted alkylene or a direct bond.

R¹⁰⁴ is a substituent such as but not limited to (un)substituted alkyl, (un)substituted aryl, (un)substituted alkenyl, (un)substituted alkynyl, (un)substituted -alkylene-aryl, (un)substituted -alkylene-cycloalkyl, and the like.

The anthranilic acid (82) may be treated with an acid chloride R¹⁰⁴-CO-Cl in the presence of a base such as TEA or aqueous alkali to afford an amide intermediate, which may be treated with a dehydrating agent such as POCl₃ or SOCl₂ in a solvent such as DCE, at a temperature of from 0°C to 80°C, to afford (83). A reagent (84) derived from an active metallating agent such as lithium or magnesium metal and G¹-L¹¹-Br or G¹-L¹¹-I may be prepared. For example, where G¹ is aryl and L¹¹ is a direct bond, G¹-L¹¹-Br may be treated with n-butyllithium in a solvent such as ether, at a temperature of from -78°C to 0°C, to afford the reagent (84) where M¹ is Li. (84) may be treated with (83) in a solvent such as THF, at a temperature of from -78°C to 50°C, to afford (85). The amide (85) may be treated with aqueous alkali in a solvent such as ethanol, at a temperature of from 25°C to 100°C, to afford (86).

Scheme 6



L¹² is, in this instance, a group such as but not limited to (un)substituted alkylene, (un)substituted cycloalkylene, or a direct bond.


- 5 An amine compound (87) may be treated with an acid chloride, or other acid halide, in the presence of boron trichloride at a temperature of from -40 °C to 25 °C, followed by treatment with gallium (III) chloride and chlorobenzene and heating at a temperature of from 50 °C to 150 °C, to afford (88).

10

Scheme 7 describes synthesis of intermediates of formulae (91), (92), (93), (94), and (95).

151

Scheme 7



L^{13} is a group such as oxygen, or may be a group broadly defined as for L^{12} and L^{11} .

R^{106} , R^{107} and R^{108} are groups such as but not limited to (un)substituted alkyl,

5 (un)substituted -alkylene-aryl, or H.

The nitrotoluene (89) can be brominated with a reagent such as N-bromosuccinimide in carbon tetrachloride to get the bromide intermediate (90). The methyl group in (89) may also be a more elaborate alkyl group with hydrogen(s) on the carbon adjacent to A^1 . The bromide (90) may be treated with sodium methanesulfinate to afford intermediate (91) and with secondary or primary amines to obtain the intermediate (92). Alternately, the R^{106} and R^{107} groups in the compound $R^{106}R^{107}NH$ may be taken together for constitute an heteroaryl or heterocyclic group, and treatment of (90) with such a compound in

10

the presence of a base such as potassium tert-butoxide affords (92) where the R^{108} and R^{107} groups are taken together for constitute a heteroaryl or heterocyclic group. Alternately, (90) may be treated with sodium thiolacetate, followed by hydrolysis with aqueous alkali, to afford the thiol (91). From (91), various compounds may be prepared. For example, treatment of


5 (91) with an alkylating agent such as an alkyl bromide in the presence of base such as sodium hydride affords (93) where L^{14} is S and R^{105} is alkyl. Oxidation of this species with a reagent such as m-chloroperbenzoic acid may afford the compound where L_{14} is $-SO_2-$. Compound (92) where R^{108} is H may be treated with a compound $R^{108}SO_2Cl$ in the presence of a base such as pyridine to form (95). Alternately, (92) where R^{108} is H may be treated with

10 a carboxylic acid $R^{108}COOH$ in the presence of a peptide coupling agent such as dicyclohexylcarbodiimide to form (94).

Scheme 8 describes synthesis of compounds of formula (97).

15

Scheme 8




L^{18} is oxygen. G^1 and L^{18} , in this instance, preferably contain no ketone, aldehyde, or primary or secondary amine groups.

20 R^{109} , R^{110} , and R^{111} are groups such as but not limited to (un)substituted alkyl, H, or (un)substituted alkylene-aryl. R^{110} and R^{111} may optionally be taken together to constitute a heterocyclic ring.

25 Ureas of formula (91) can be reductively aminated using amines $R^{110}NHR^{111}$ and a reagent such as sodium triacetoxyborohydride in a solvent such as 1,2-dichloroethane in the presence or absence of acetic acid to obtain compounds of formula (92).

Scheme (9) describes synthesis of compounds of formulae (100) and (101).



5

L¹⁷ is carbonyl or sulfonyl group.


Nitrophenylureas (98) can be reduced to aniline derivatives of formula (99).

Treatment of intermediate (99) with acid chlorides or sulfonyl chlorides can yield compounds of formula (100). Alkylation of intermediate (99) using aldehydes or ketones in the presense of sodium triacetoxyborohydride affords (101). Alternately, (99) may be treated with a dialkyl halide and a base such as DIEA to afford (101) where R¹¹³ and R¹¹⁴ and the nitrogen to which they are attached constitute a ring.

10

5

Scheme 10



10 L^{19} in this instance is a group such as (un)substituted alkylene. R^{115} and R^{116} are independently, groups such as (un)substituted alkyl, (un)substituted alkylene-aryl, or H. Alternately, R^{115} and R^{116} may be taken together to constitute a heterocyclic ring.

The acid (102) may be coupled with an amine $R^{115}NHR^{116}$ in the presence of a
 15 coupling agent such as dicyclohexylcarbodiimide in a solvent such as THF or dichloromethane to afford (103).


20

Scheme 11

25

155

Scheme 11



A^3 is a group such as (un)substituted heteroarylene, (un)substituted fused heterocyclylheteroarylene, or (un)substituted fused cycloalkylheteroarylene.


5 R^{117} is a group such as (un)substituted alkyl, (un)substituted alkylene-aryl, (un)substituted aryl, or (un)substituted heteroaryl.

The compound (104) may be treated with a thiol reagent in the presence of a base such as DIEA at temperatures of from 50 °C to 150 °C to afford the thioether (105). (105) may be oxidized with an oxidizing reagent such as m-chloroperoxybenzoic acid in a solvent such as dichloromethane to afford the sulfone (106). Where only one equivalent of the oxidant is employed, the sulfoxide may be obtained. Where 2 or more equivalents of oxidant are employed, the sulfone is obtained.

15 Scheme 12 describes the synthesis of compounds of formula (109).

156

Scheme 12



A^3 is a group such as (un)substituted heteroarylene, (un)substituted fused heterocycl/heteroarylene, or (un)substituted fused cycloalkylheteroarylene.


5 L^{20} in this instance is a group such as (un)substituted alkylene. R^{117} is a group such as (un)substituted alkyl, (un)substituted alkylene-aryl, (un)substituted aryl, or (un)substituted heteroaryl. Lg_2 is a leaving group such as chloride, methanesulfonate, or p-toluenesulfonate.

The compound (107) where Lg_2 is methanesulfonate may be synthesized from the precursor where Lg_2 is hydroxyl by treatment with methanesulfonyl chloride in the presence of pyridine. (107) then may be treated with a thiol reagent in the presence of a base such as DIEA, potassium tert-butoxide, or sodium hydride, to afford the displacement product (108). The thioether product (108) may be oxidized to the sulfoxide or sulfone (109) as described in Scheme 11.

15

Scheme 13 describes the synthesis of compounds of formula (111).

Scheme 13




L^{21} in this instance is a group such as alkylene. R^{115} and R^{116} may have the meaning denoted previously, or may be, independently, groups such as (un)substituted alkyl, (un)substituted alkylene-aryl, (un)substituted aryl, H, or (un)substituted heteroaryl. A^3 is a group such as (un)substituted heteroarylene, (un)substituted fused heterocyclylheteroarylene, or (un)substituted fused cycloalkylheteroarylene.

The acid (110) may be coupled with an amine $R^{115}NHR^{116}$ in the presence of a coupling agent such as dicyclohexylcarbodiimide to afford (111).

Scheme 14 describes a synthesis of intermediates of formula (113).

Scheme 14



R^{118} and R^{119} may be, independently, groups such as (un)substituted alkyl, (un)substituted alkylene-aryl, (un)substituted aryl, H, or (un)substituted heteroaryl.


Thiourea may be condensed with the bromo carbonyl compound (112) in the presence or absence of a mild base such as potassium carbonate or triethylamine, in a solvent such as ethanol, at a temperature of from 25 °C to 120 °C, to afford (113). The bromo compound (112) may be accessed by a variety of methods known in art. For example, bromination of the ketone with pyrrrolidinium hydrotribromide in THF or N-

bromosuccinimide in THF in the presence of a mild base such as potassium carbonate affords (112).

Scheme 15 describes synthesis of intermediates of formula (117)

5

Scheme (15)



10 Scheme 15 shows the synthetic route to diamides of the type (117), where A^1 , L^1 , R^{20} , R^1 , G^1 , G^2 are as defined in Formula (I). Amine (114) can be coupled with an activated oxalic or malonic esters using EDC/HOBT to give amide (115). Deprotection of the t-butyl ester of B is done with lithium hydroxide to give the carboxylic acid (116), which can be coupled using standard amide coupling reagents (eg. PyBOP) to give diamides of the type

15 (115).

The intermediates in the above Schemes may be substituted with amino, hydroxyl, or carboxyl groups which may require protection and deprotection during the course of preparation of Example compounds.

20

"Amino protection" refers to substituents of the amino group commonly employed to block or protect the amino functionality while reacting other functional groups on the

compound. Examples of such amino-protecting groups include the formyl group, the trityl group, the phthalimido group, the trichloroacetyl group, the chloroacetyl, bromoacetyl and iodoacetyl groups, urethane-type blocking groups such as benzyloxycarbonyl, 4-phenylbenzyloxycarbonyl, 2-methylbenzyloxycarbonyl, 4-methoxybenzyloxycarbonyl, 4-fluorobenzyloxycarbonyl, 4-chlorobenzyloxycarbonyl, 3-chlorobenzyloxycarbonyl, 2-chlorobenzyloxycarbonyl, 2,4-dichlorobenzyloxycarbonyl, 4-bromobenzyloxycarbonyl, 3-bromobenzyloxycarbonyl, 4-nitrobenzyloxycarbonyl, 4-cyanobenzyloxy-carbonyl, 2-(4-xenyl)iso-propoxycarbonyl, 1,1-diphenyleth-1-yloxycarbonyl, 1,1-diphenylprop-1-yloxycarbonyl, 2-phenylprop-2-yloxycarbonyl, 2-(p-toluy)prop-2-yloxycarbonyl, cyclopentanyloxycarbonyl, 1-methylcyclopentanyloxycarbonyl, cyclohexanyloxycarbonyl, 1-methylcyclohexanyloxycarbonyl, 2-methylcyclohexanyloxycarbonyl, 2-(4-toluylsulfonyl)ethoxycarbonyl, 2(methylsulfonyl)ethoxycarbonyl, 2-(triphenylphosphino)ethoxycarbonyl, 9-fluorenylmethoxycarbonyl ("Fmoc"), t-butoxycarbonyl ("BOC"), 2-(trimethylsilyl)ethoxycarbonyl, allyloxycarbonyl, 1-(trimethylsilylmethyl)prop-1-enyloxycarbonyl, 5-benzisoxalylmethoxycarbonyl, 4-acetoxybenzyloxycarbonyl, 2,2,2-trichloroethoxycarbonyl, 2-ethynyl-2-propoxycarbonyl, cyclopropylmethoxycarbonyl, 4-(decyloxy)benzyloxycarbonyl, isobornyloxycarbonyl, 1-piperidyloxycarbonyl and the like; the benzoylmethylsulfonyl group, the 2-(nitro)phenylsulfonyl group, the diphenylphosphine oxide group and like amino-protecting groups. The species of amino-protecting group employed is not critical so long as the derivatized amino group is stable to the condition of subsequent reaction(s) on other positions of the compound of Formula (I) and can be removed at the desired point without disrupting the remainder of the molecule. Preferred amino-protecting groups are the allyloxycarbonyl, the t-butoxycarbonyl, 9-fluorenylmethoxycarbonyl, and the trityl groups. Similar amino-protecting groups used in the cephalosporin, penicillin and peptide art are also embraced by the above terms. Further examples of groups referred to by the above terms are described by J. W. Barton, "Protective Groups In Organic Chemistry", J. G. W. McOmie, Ed., Plenum Press, New York, N.Y., 1973, and T. W. Greene, "Protective Groups in Organic Synthesis", John Wiley and Sons, New York, N.Y., 1981. The related term "protected amino" defines an amino group substituted with an amino-protecting group discussed above.

"Hydroxyl protection" refers to substituents of the alcohol group commonly employed to block or protect the alcohol functionality while reacting other functional groups on the compound. Examples of such alcohol -protecting groups include the 2-tetrahydropyranyl

group, 2-ethoxyethyl group, the trityl group, the trichloroacetyl group, urethane-type blocking groups such as benzyloxycarbonyl, and the trialkylsilyl group, examples of such being trimethylsilyl, tert-butyldimethylsilyl, phenyldimethylsilyl, triisopropylsilyl and thexyldimethylsilyl. The choice of alcohol-protecting group employed is not critical so long as the derivatized alcohol group is stable to the condition of subsequent reaction(s) on other positions of the compound of the formulae and can be removed at the desired point without disrupting the remainder of the molecule. Further examples of groups referred to by the above terms are described by J. W. Barton, "Protective Groups In Organic Chemistry", J. G. W. McOmie, Ed., Plenum Press, New York, N.Y., 1973, and T. W. Greene, "Protective Groups in Organic Synthesis", John Wiley and Sons, New York, N.Y., 1981. The related term "protected hydroxyl" or "protected alcohol" defines a hydroxyl group substituted with a hydroxyl - protecting group as discussed above.

"Carboxyl protection" refers to substituents of the carboxyl group commonly employed to block or protect the -OH functionality while reacting other functional groups on the compound. Examples of such alcohol -protecting groups include the 2-tetrahydropyranyl group, 2-ethoxyethyl group, the trityl group, the allyl group, the trimethylsilylethoxymethyl group, the 2,2,2-trichloroethyl group, the benzyl group, and the trialkylsilyl group, examples of such being trimethylsilyl, tert-butyldimethylsilyl, phenyldimethylsilyl, triisopropylsilyl and thexyldimethylsilyl. The choice of carboxyl protecting group employed is not critical so long as the derivatized alcohol group is stable to the condition of subsequent reaction(s) on other positions of the compound of the formulae and can be removed at the desired point without disrupting the remainder of the molecule. Further examples of groups referred to by the above terms are described by J. W. Barton, "Protective Groups In Organic Chemistry", J. G. W. McOmie, Ed., Plenum Press, New York, N.Y., 1973, and T. W. Greene, "Protective Groups in Organic Synthesis", John Wiley and Sons, New York, N.Y., 1981. The related term "protected carboxyl" defines a carboxyl group substituted with a carboxyl -protecting group as discussed above.

EXAMPLES**General Procedure A: Preparation of 1-Aryloxy-2-nitrobenzenes, 1-Arylsulfanyl-2-nitrobenzenes and 2-Aryloxy-3-nitropyridines**

To a solution of potassium *t*-butoxide (0.62 g, 5.5 mmol) in anhydrous DMF (10 ml) was added a phenol, arylmercaptan or 2-mercaptopyridine (5.5 mmol) at room temperature and the mixture was stirred for 30 min. A 1-fluoro-2-nitrobenzene derivative or 2-bromo-3-nitropyridine (5.0 mmol) was added and the contents were heated at 80°C for 12 h. The contents were poured into water and extracted with ethyl acetate. The organic layer was washed (dil. NaOH, water, brine), dried (Na₂SO₄) and concentrated. In general, the desired products were of >90% pure and were used as such for further manipulations.

General Procedure B: Preparation of 2-Aryloxyanilines, 2-Arylsulfanylanilines and 3-amino-2-aryloxypyridines

The crude 2-substituted-1-nitrobenzene from procedure A was dissolved in ethanol (10 ml). To this solution were added anhydrous tin(II) chloride (3.8 g, 20 mmol) and conc HCl (0.2 ml). The resulting mixture was heated at 80°C for 10 h, cooled and concentrated. The residue was diluted with water (100 ml), neutralized to pH 8-9. To the suspension, ethyl acetate (40 ml) was added, stirred for 5 min and filtered through celite. The layers were separated and the aqueous layer was extracted with ethyl acetate (2x10 ml). The combined organic layer was washed with brine, dried (Na₂SO₄) and concentrated under reduced pressure to obtain the desired aniline in 60-70% yield. In general, the desired anilines were of >85% pure (LC-MS) and were used as such for further manipulations.

General Procedure C for Preparation of 2-Aryloxyanilines and 2-Arylsulfanylanilines

The crude 2-substituted-1-nitrobenzene (~5 mmol) was dissolved in methanol (10 ml) in a 100 ml round-bottom flask. To this solution was added 10% palladium on charcoal (300 mg) and the flask was evacuated. The flask was filled with hydrogen with the aid of a balloon and the contents were stirred overnight. The mixture was filtered through celite and concentrated to obtain desired aniline (>90% purity by LC-MS).

General Procedure D: Preparation of urea

A mixture of 1,1'-carbonyldiimidazole (98 mg, 0.6 mmol), 2-aminoheteroarene (0.6 mmol) and 4-(*N,N*-dimethylamino)pyridine (5 mg) in dichloroethane (5 ml) was heated at 80°C for 2 h. The reaction mixture was cooled to room temperature and was added solution

of a substituted aniline (0.5 mmol) in dichloroethane (2 ml). The resulting suspension was heated at 80°C for 10 h and concentrated. The residue was purified by column chromatography (silica, CH₂Cl₂ then 10-30% ethyl acetate in CH₂Cl₂) to afford the desired urea in 60-80% yield.

5 General Procedure E: Preparation of urea

A mixture of isocyanate (0.5 mmol) and 2-aminoheteroarene (0.5 mmol) in dichloroethane (4 ml) was heated at 80°C for 12 h. The reaction mixture was concentrated under reduced pressure. The residue was purified by column chromatography (silica, CH₂Cl₂ then 10-30% ethyl acetate in CH₂Cl₂) to afford the desired urea in 60-80% yield.

10 General Procedure F: for Preparation of 2-Aryloxy-1-nitrobenzenes

To a solution of potassium *t*-butoxide (0.62 g, 5.5 mmol) in anhydrous THF (20 ml) was added a phenol (5.5 mmol) at -10°C and the mixture was stirred for 30 min. A fluoro-1-nitrobenzene derivative (5.0 mmol) was added at -10°C and stirred for 12 h at room temperature. The contents were poured into water (25 ml) and extracted with ethyl acetate (3x20 ml). The organic layer was washed (dil. NaOH, water, brine), dried (Na₂SO₄) and concentrated to give the desired products with >90% purity by LC-MS and were used as such in the next step.

General Procedure G: for Preparation of 1-Alkoxy-2-nitrobenzenes

To a suspension of NaH (60%, 0.20 g, 5.0 mmol) in anhydrous THF (10 ml) was added an alcohol (5.0 mmol) dropwise at room temperature and the mixture was stirred for 30 min. 1-Fluoro-2-nitrobenzene (5.0 mmol) was added and the contents were heated at 60°C for 12 h. The contents were poured into water and extracted with ethyl acetate. The organic layer was washed (dil. NaOH, water, brine), dried (Na₂SO₄) and concentrated. In general, the desired products were of >90% pure and were used as such in the further manipulations.

General procedure H: General procedure for preparation of compounds of general formula I that contain a urea moiety in the central core.

One equivalent of a mono- di- or tri.substituted aniline is dissolved in an organic solvent such as ethyl acetate, toluene, or dichloromethane and hydrochloride dissolved in an organic solvent such as ethyl acetate, toluene or dichloromethane is added. The mixture is concentrated *in vacuo* to give the hydrochloride of the aniline. The residue is dissolved or

suspended in a non protic organic solvent such as toluene or dichloromethane and excess (e.g. 2 to 5 equivalents) of diphosgene or another phosgene equivalent is added. The mixture is either run at room temperature or heated (up to reflux temperature of the solvent) for 5 to 20 hours. The reaction mixture is concentrated *in vacuo* and the intermediate residue
5 used in the next step without further purification.


The crude intermediate isocyanate is dissolved in an organic solvent such as ethyl acetate, toluene, dichloromethane, dioxane, DMSO or DMF and one equivalent of a heterocyclic amine is added. The reaction mixture is run at either room temperature or heated until the reaction is taking place. The reaction temperature will depend on the
10 reactivity of the isocyanate and the nucleophilicity of the amine and can be followed either using HPLC or TLC. The reaction mixture is diluted with an organic solvent such as ethyl acetate, toluene or dichloromethane and the mixture extracted with water. The product is purified using standard procedures as described in the art or as exemplified below.

General Procedure H1 : Preparation of amides from carboxylic acids prepared using
15 procedure H:

One equivalent of a N-substituted aminothiazol-4-ylcarboxylic acid or N-substituted aminothiazol-4-ylacetic acid prepared by using general procedure H is dissolved in an organic solvent such as 1,2-dichloropropane, dimethylformamide or a mixture of two organic solvents such as a mixture of 1,2-dichloropropane and dimethylformamide. One equivalent of
20 PyBOP (benzotriazole-1-yl-oxy-tris-pyrrolidino-phosphonium hexafluorophosphate) is added, the reaction mixture is left standing for 20 minutes followed by addition of two equivalents of an appropriate amine and DIPEA (diisopropylethylamine), and the mixture is left overnight.


The reaction mixture is diluted with ethylacetate and extracted using a general washing procedure such as washing twice with water, twice with 4N HCl, once with water,
25 twice with 50% saturated sodiumhydrogencarbonate, and three times with water. The organic solvent is evaporated *in vacuo* giving an amorphous product. The product is purified by either recrystallization in an organic solvent such as diethylether or by HPLC (e.g. a Waters Deltaprep 4000).

In case the isolated product contains a carboxylic acid ester functionality, the ester
30 group can be hydrolysed to the corresponding acid by dissolving the compound in ethanol 96% and adding 2N NaOH. The mixture is left standing for some time (e.g. 2 hours) whereafter the ethanol is evaporated *in vacuo*, water is added, and pH is adjusted to acidic with 2N HCl. The mixture is extracted with an organic solvent such as ethylacetate and the combined organic phases are evaporated *in vacuo* giving an amorphous product.




General procedure H2: Preparation of intermediate isocyanates:2-Benzylphenyl isocyanate

- 5 2-Benzylaniline (2.0 g, 11 mmol) was dissolved in ethylacetate (5 ml) and hydrochloride in ethylacetate (3N, 5 ml) was added. After 2 hrs the organic solvent was removed *in vacuo* giving a solid residue. Toluene (50 ml) was added, then diphosgene (2.2 g, 33 mmol) and the reaction-mixture was heated at 110°C for 16 hours. The solvent and excess diphosgene was removed *in vacuo* giving an residual oil that was used for the next
- 10 step without further purification.

The following isocyanates were prepared using the same procedure used for preparation of benzylphenyl isocyanate:

(5-Chloro-2-isocyanatophenyl)phenyl methanone

- 15 2-(2-Methylphenoxy)phenyl isocyanate

2-(4-Methoxyphenoxy)-5-(trifluoromethyl)phenyl isocyanate2-(Phenylsulfonyl)phenyl isocyanate

General Procedure I: Preparation of 2-acyl anilines

A solution of 1 M borontrichloride in dichloromethane (110 mL, 0.11 mol) was cooled to -20°C . To this solution was added a solution of aniline (0.1 mol) in dichloromethane (100 mL). The mixture was warmed to room temperature and was stirred for 3 h. The mixture was re-cooled to -20°C . Alkyl nitrile (0.1 mol) was added over 5 min, followed by 1 M solution of (anhydrous) GaCl_3 (100 mL, 0.1 mol) in dichloromethane. To this solution was added chlorobenzene (300 mL) and the mixture was heated to reflux for 24 h. After being cooled to room temperature, the mixture was poured into ice water (1 L) and the mixture was stirred for 3 h. The organic layer was separated and the aqueous layer was extracted with dichloromethane (4 x 400 mL). The combined organic layer was washed with water (4 x 500 mL), brine (2 x 500 mL), dried over anhydrous Na_2SO_4 . The aqueous layer was then basified to pH 7.5 with Na_2CO_3 and the mixture was extracted with dichloromethane (2 x 400 mL). The organic layer was washed with water (4 x 500 mL), brine (2 x 500 mL), dried over anhydrous Na_2SO_4 . Both organic layers were combined and concentrated in *vacuo*. The crude mixture was purified by column chromatography with hexanes-ethyl acetate (9:1) as eluent to give the 2-amino-alkylphenones in 10-50% yield.

General Procedure J: Preparation of acids from esters

The ester (1 mmol) was dissolved in 1:1 mixture of THF and methanol (5 mL). To this solution was added 2 M solution of LiOH (2 mL, 4 mmol). The mixture was stirred for 1 h and was concentrated. The residue was diluted with water (10 mL) and the aqueous layer was washed with ether (2 x 10 mL). The water layer was acidified with HCl to pH 6.0 and precipitated acid was extracted with ethyl acetate (2 x 50 mL). The organic layer was washed with water (2 x 20 mL), dried (Na_2SO_4) and concentrated in *vacuo* to furnish the desired acid in almost quantitative yield.

General Procedure K: Preparation of amides

A mixture of acid (0.5 mmol) and HBTU (0.5 mmol) was dissolved in anhydrous DMF (2 mL). To this solution was added DIEA (0.6 mmol) and stirred for 2-3 minutes. A solution of alkyl amine (0.5 mmol) in DMF (1 mL) was added and the mixture was stirred at room temperature for 30 min. The mixture was poured into water (20 mL) and was extracted with ethyl acetate (2 x 20 mL). The organic layer was washed with saturated solution of citric acid (5 mL), NaHCO_3 (2 x 10 mL) water (2 x 0 mL), brine (2 x 10 mL), dried (Na_2SO_4) and

concentrated in *vacuo* to give the desired amide. The crude mixture was purified by column chromatography (silica, CH₂Cl₂ then 10-50% ethyl acetate in CH₂Cl₂) to furnish amide in 50-75% yield.

5 General Procedure L: Preparation of sulfonamides/amides

To a solution of acid (1.0 mmol) and DIEA (1.5 mmol) in anhydrous THF (20 mL) was added diphenylphosphoroyl azide (1.5 mmol) and was heated to reflux for 8-12 hours. The reaction mixture was then concentrated in *vacuo* to give crude isocyanate. To this crude product was added dilute HCl (1.2 M, 20 mL) and the mixture was heated to reflux for 2
10 hours. The reaction mixture was neutralized with Na₂CO₃ and the aqueous layer was extracted with ethyl acetate (3 x 30 mL). The organic layer was washed with water (2 x 30 mL), brine (1 x 30 mL) and dried (anhydrous Na₂SO₄) and concentrated in *vacuo* to give the desired amine. This crude amine (1.0 mmol) was reacted with aryl/alkylsulfonyl chloride (1 mmol) and Et₃N (2 mmol) to give the desired sulfonamides. The amides were prepared as
15 described in procedure K. The crude product was purified by silica gel chromatography [hexanes:EtOAc/MeOH (70:30:0 to 5:90:5)] to furnish the desired sulfonamides in 20-30% yield.

General Procedure M: Preparation of bis-ureas or carbamates

20 A mixture of acid (1.0 mmol) and DIEA (1.5 mmol) was dissolved in anhydrous THF or CH₃CN (30 mL). Then diphenylphosphoroyl azide (1.5 mmol) added to the reaction mixture. Reaction mixture was refluxed for 8-12 hours. The reaction mixture was then concentrated in *vacuo* to give crude isocyanate. To this crude product desired amines or alcohols (2.0 mmol) were added and stirred at rt for 2h. The reaction mixture was then concentrated in *vacuo*.
25 The crude reaction mixture was then purified by silica gel chromatography (hexanes:EtOAc 70:30 to 10:90) to furnish the desired bis-ureas or carbamates in 30-45% yield.

General Procedure N: Preparation of alcohols

To a solution of ethyl-2-amino-4-thiazolyl acetate or ethyl-2-amino-4thiazoyl carboxylate (100 mmol) in anhydrous THF (100 mL) was added lithiumborohydride (200 mmol, 2.0 M solution in THF) at -10 °C, and the mixture was allowed to warm up to ambient temperature and stirred for 8-10 h. The mixture was then concentrated *in vacuo*. Methanol (200 mL) was added to quench excess lithiumborohydride and filtered through with a plug of silica gel to afford the amino alcohol.

To this crude amino alcohol (100 mmol) and imidazole (500 mmol) in anhydrous DMF (50 ml) was added *tert*-butyldimethylsilyl chloride (500 mmol) and stirred at rt for 6 h. The reaction mixture was then washed with water (5x100 mL) and brine (2x100 mL) and extracted with ethylacetate (3x200 mL), dried over Na₂SO₄, and concentrated *in vacuo* to give TBS-protected amino alcohol.

TBS-protected amino alcohol (50 mmol) was subjected to urea formation following general procedure D to give desired urea. This crude urea (25 mmol) was then treated with TBAF (50 mmol, 1.0 M solution in THF) and stirred at rt for 4 h. The reaction mixture was poured in to water and extracted with ethyl acetate. The organic extracts were combined, washed (water), dried (Na₂SO₄) and concentrated *in vacuo*. The crude mixture was purified by silica gel chromatography [hexanes:EtOAc (70:30 to 10:90)] to afford the desired alcohol in 70-80 % yield.

General Procedure O: Preparation of Amines by Reductive Amination

To the aldehyde (0.11mmol) in dichloroethane or THF (5 mL) was added the respective amine (0.11mmol) and stirred at room temperature for 15 min. To this solution was added sodium triacetoxyborohydride (0.16 mmol). After stirring at room temperature overnight, the mixture was concentrated *in vacuo* and purified by column chromatography (silica, 2-8% MeOH in DCM) to obtain the desired product in 30-50 % yield.

General Procedure P: Preparation of Arylethers by Mitsunobu Reaction

To a solution of 1-[2-(cyclopentanecarbonyl-4-methyl-phenyl)-3-[4-(2-hydroxy-ethyl)-thiazol-2-yl]-urea (0.268 mmol), phenol (0.536 mmol) and triphenylphosphine (0.268mmol) in THF (2 mL) was added diisopropyl azodicarboxylate (0.268mmol) at 0°C. The resulting solution was stirred at rt overnight, The mixture was concentrated and purified by flash chromatography on silica gel (10-50% EtOAc/ hexane) to give the desired product in 28-40% yield.

General procedure Q: Synthesis of 1-(2-cyclopentanoyl-4-methyl-phenyl)-3-(5-alkylthio-2-thiazolyl)ureas

A mixture of 1-(2-cyclopentanoyl-4-methyl-phenyl)-3-(5-bromol-2-thiazolyl)urea (1 mmol), alkylthiol (2 mmol) and DIEA (2 mmol) in DMF (5 mL) was heated at 80°C for 3 h.

- 5 The mixture was poured into water (20 mL) and was extracted with ethyl acetate (3 x 25 mL). The organic layer was washed with water (2 x 30 mL), brine (1 x 30 mL), dried (anhydrous Na₂SO₄) and concentrated in *vacuo* to furnish a residue containing 1-(2-cyclopentanoyl-4-methyl-phenyl)-3-(5-alkylthio-2-thiazolyl)urea along with 1-(2-cyclopentanoyl-4-methyl-phenyl)-3-(thiazol-2-yl)urea. The crude product was purified by column chromatography (silica, CH₂Cl₂ then 5-20% ethyl acetate in CH₂Cl₂) to afford the desired product in 25-35 % yield

- Same procedure was adopted for the synthesis of 1-(2-cyclopentanoyl-4-methyl-phenyl)-3-(5-arylthio-2-thiazolyl)ureas. The crude products were purified by column chromatography (silica, CH₂Cl₂ then 5-20% ethyl acetate in CH₂Cl₂ and 2% MeOH in CH₂Cl₂) to afford the desired product in 25-35 % yield.

General Procedure R: Oxidation of alkyl and arylthio substituted thiazolyl ureas

- Alkyl or arylthio substituted thiazolyl urea (0.5 mmol) was dissolved in CH₂Cl₂ (5 mL) and was cooled to 0° C in an ice bath. To this solution was added *m*-cpba (133 mg, 0.75 mmol) in CH₂Cl₂ (3 mL). The mixture was stirred at 0°C for 4 h and was diluted with CH₂Cl₂ (30 mL). The organic layer was washed with saturated solution of NaHCO₃ (2 x 20 mL), water (3 x 20 mL), brine (1 x 20 mL), dried (anhydrous Na₂SO₄) and concentrated in *vacuo*. The crude mixture was purified by column chromatography (silica, CH₂Cl₂ then 5-20% ethyl acetate in CH₂Cl₂ and 2% MeOH in CH₂Cl₂) to give the desired alkyl or aryl sulfone in 60-80% yield.

General Procedure S: Preparation of 2-Amino Arylphenones

- To a solution of 2-amino benzoic acid (10 mmol) in THF was added benzoyl chloride (2.8 g, 20 mmol) followed by pyridine (1.58 g, 20 mmol). The mixture was stirred for 1h at room temperature. The 2-phenyl-benzo[d][1,3]oxazin-4-one formed was filtered and the residue was washed with water and dried in vacuum desiccator.

- 30 To a solution of 2-phenyl-benzo[d][1,3]oxazin-4-one (5 mmol) in dry CH₂Cl₂ (20 mL) was added 1N solution of aryl magnesium bromide in THF (5 mL) at 0° C. The mixture was stirred at room temperature for 2 h and poured into water (30 mL). The aqueous layer was extracted with ethyl acetate (3 x 30 mL) and was washed with water (3 x 50 mL), brine (1 x 50 mL), dried (anhydrous Na₂SO₄) and concentrated in *vacuo* to afford N-(2-benzoyl-phenyl)-benzamide in 60-70% yield.

To a solution of the crude N-(2-benzoyl-phenyl)-benzamide (2 mmol) in THF (10 mL) was added 10 N solution of NaOH (5 mL) and was heated to reflux for 18 h. The mixture was poured into water (50 mL) and was extracted with ethyl acetate (3 x 30 mL). The organic layer was washed with water (3 x 50 mL), brine (1 x 50 mL), dried (anhydrous Na₂SO₄) and concentrated in *vacuo* to afford 2-amino arylphenone. The crude product was purified by column chromatography (silica, hexanes then 5-20% ethyl acetate) to furnish the desired product in 28-40 % yield.

General Procedure T: Preparation of Amides/Sulfonamides

To a solution of amine (0.5 mmol) in DCM (5 mL) was added triethylamine (1 mmol) and cooled the reaction mixture to 0 °C. Acid chloride or sulfonyl chloride (0.5 mmol) was added drop-wise and stirred for overnight. The reaction mixture was concentrated in *vacuo* and the residue was purified by column chromatography [silica, DCM:ethyl acetate (80:20 to 20:80)] to yield desired amides or sulfonamides respectively.

General Procedure U: Preparation of Hydantoins from Amino Acids:

To a solution of Boc-Gly-Merrifield resin (1.2 g, 0.96 mmol) was added trifluoroacetic acid (5 ml, 20% in DCM), then the resin was washed with three cycles of DMF, methanol, and DCM. To this resin in DCM (20 mL) was slowly added phosgene (10 mL, 20% in toluene, 2.0 mmol) and triethylamine (0.56 ml, 4.0 mmol) at -20 °C. The reaction mixture was allowed to warm up to room temperature. The excess phosgene was washed away with three cycles of DCM. To this resin in DCM (20 mL) was added 1-(4-aminomethyl-thiazol-2-yl)-3-(2-cyclopentanecarbonyl-4-methyl-phenyl)-urea (0.9 g, 2.5 mmol) in DCM (10 mL) and the reaction mixture was placed in a shaker and reacted for 4 h to give the corresponding urea. The resin was then washed with three cycles of DMF, methanol and DCM and dried over 2 h. To the resin was added triethylamine (10 mL, 20% solution in THF) and the reaction mixture was heated for 16 h. The mixture was filtered and the filtrate was concentrated *in vacuo* to afford hydantoins in 60-75% overall yields.

General Procedure V: Preparation of Specific 2-Aminothiazole Analogs:

To a solution of 1,3-dichloroacetone, 1,3-dibromoacetone, 1-acetoxy-3-chloroacetone, bromomalonaldehyde or 1,4-dibromobutan-2,3-dione (100 mmol) in methanol (100 ml) was added thiourea (7.6 g, 100 mmol) and the mixture was stirred at rt. for 3h. The reaction mixture was concentrated *in vacuo* to give the desired products in almost quantitative yields.

4-Chloromethyl-thiazole-2-ylamine (172 mg, 1.0 mmol) was reacted with arylthiols (2 mmol) and DIEA (2 mmol) in THF (5 mL) following the general procedure Z. These intermediates were coupled with CDI and 2-amino-5-methyl-phenyl)-cyclopentyl-methanone (203 mg, 1.0 mmol) following the general procedure D.

5

General Procedure W: Preparation of alkylamino nitrobenzenes

1-Fluoro-2-nitrobenzene derivative (5.0 mmol) and an amine (10 mmol) in THF (25 mL) were heated at 60 °C for 12 h. The contents were poured into water and extracted with ethyl acetate. The organic layer was washed (water, brine), dried (Na₂SO₄) and concentrated. The residue was dissolved in methanol (25 mL) and subjected reduction following the general procedure C. In general, the desired products were of >90% pure and were used as such in the further manipulations.

10

General Procedure X: Preparation of Alkenes by Wittig Reaction

15 The aldehyde (0.10g, 0.28mmol) and (carbethoxymethylene)-triphenylphosphorane (0.12g, 0.34mmol) were stirred at room temperature in benzene overnight. The reaction mixture was concentrated under vacuum and purified by column chromatography (silica, 15% EtOAc/hexanes) to obtain the product in 80% yield.

15

20 General procedure Y: Synthesis of 1-(2-cyclopentanoyl-4-methyl-phenyl)-3-(5-arylthio-2-thiazolyl)ureas

A mixture of 1-(2-cyclopentanoyl-4-methyl-phenyl)-3-(5-bromol-2-thiazolyl)urea (1 mmol), arylthiol (2 mmol) and tert. BuOK (2 mmol; 4 equivalent of tert.BuOK was used for arylthio carboxylic acids) in DMF (5 mL) was heated at 80°C for 3 h. The mixture was poured into water (20 mL). Urea containing arylthio carboxylic acid was neutralized with saturated NaHCO₃ solution. The aqueous layer was extracted with ethyl acetate (3 x 25 mL). The organic layer was washed with water (2 x 30 mL), brine (1 x 30 mL), dried (anhydrous Na₂SO₄) and concentrated in *vacuo* to furnish a residue containing 1-(2-cyclopentanoyl-4-methyl-phenyl)-3-(5-arylthio-2-thiazolyl)urea along with 1-(2-cyclopentanoyl-4-methyl-phenyl)-3-(2-thiazolyl)urea. The crude product was purified by column chromatography (silica, CH₂Cl₂ then 5-20% ethyl acetate in CH₂Cl₂ and 2% MeOH in CH₂Cl₂) to afford the desired product in 25-35 % yield.

25

30

General procedure Z: Synthesis of 1-(2-cyclopentanecarbonyl-4-methyl-phenyl)-3-[4-(aryl-sulfanylmethyl)-thiazol-2-yl]-ureas and 1-(2-cyclopentanecarbonyl-4-methyl-phenyl)-3-[4-(arylsulfanyl)-ethyl]-thiazol-2-yl]-ureas

- 5 A mixture of 1-(4-chloromethyl-thiazol-2-yl)-3-(2-cyclopentanecarbonyl-4-methyl-phenyl)-urea (1 mmol), arylthiol (2 mmol) and DIEA (2 mmol) in THF (5 mL) was heated at 80°C for 3 h. The mixture was poured into water (20 mL) and was extracted with ethyl acetate (3 x 25 mL). The organic layer was washed with water (2 x 30 mL), brine (1 x 30 mL), dried (anhydrous Na₂SO₄) and concentrated in *vacuo* to furnish a residue containing 1-(2-
- 10 cyclopentanecarbonyl-4-methyl-phenyl)-3-[4-(arylsulfanylmethyl)-thiazol-2-yl]-urea. The crude product was purified by column chromatography (silica, CH₂Cl₂ then 5-20% ethyl acetate in CH₂Cl₂ and 2% MeOH in CH₂Cl₂) to afford the desired product in 79-85 % yield

- Similarly, synthesis of 1-(2-cyclopentanecarbonyl-4-methyl-phenyl)-3-[4-(arylsulfanyl)-ethyl]-thiazol-2-yl]-urea was carried out by reacting methanesulfonic acid 2-{2-[3-(2-
- 15 cyclopentanecarbonyl-4-methyl-phenyl)-ureido]-thiazol-4-yl}-ethyl ester with arylthiol and Et₃N. This afforded the desired product in 60-80 % yield.

General procedure AA: Preparation of urea.

- A mixture of 1,1'-carbonyldiimidazole (98 mg, 0.6 mmol), (2-aminothiazol-4-yl)acetic acid ethyl ester (0.6 mmol) and 4-(*N,N*-dimethylamino)pyridine (2 mg) in dichloromethane (5
- 20 ml) was stirred at room temperature for 2 h. A solution of a substituted aniline derivative (0.6 mmol) in dichloromethane (1 ml) was added and stirring was continued at room temperature for 24 h. The reaction mixture was concentrated and the residue was purified by column chromatography (silica, CH₂Cl₂ then 10-30% ethyl acetate in CH₂Cl₂) to afford the desired urea.

25 **HPLC-MS (Method A)**

The following instrumentation is used:

- Hewlett Packard series 1100 G1312A Bin Pump
- Hewlett Packard series 1100 Column compartment
- Hewlett Packard series 1100 G1315A DAD diode array detector
- 30 • Hewlett Packard series 1100 MSD
- Sedere 75 Evaporative Light Scattering detector

The instrument is controlled by HP Chemstation software.

The HPLC pump is connected to two eluent reservoirs containing:

A: 0.01% TFA in water

B: 0.01% TFA in acetonitrile

The analysis is performed at 40°C by injecting an appropriate volume of the sample (preferably 1 µl) onto the column which is eluted with a gradient of acetonitrile.

5 The HPLC conditions, detector settings and mass spectrometer settings used are giving in the following table.

Column: Waters Xterra MS C-18 X 3 mm id 5 µm

Gradient: 5% - 100% acetonitrile linear during 7.5 min at 1.5ml/min

Detection: 210 nm (analogue output from DAD (diode array detector))

ELS (analogue output from ELS)


10 MS ionisation mode API-ES

Scan 100-1000 amu step 0.1 amu

After the DAD the flow is divided yielding approx 1 ml/min to the ELS and 0.5 ml/min to the MS.

Example 1


15 N-(2-Phenoxyphenyl)-N'-(thiazol-2-yl)urea



N-(2-Phenoxyphenyl)-N'-(thiazol-2-yl)urea (0.59 g, 94.9%) was prepared from 2-phenoxyaniline (0.37 g, 2.00 mmol) and 2-aminothiazole (0.20 g, 2.00 mmol) following the general procedure D.

20 LC-MS (m/z): 313 ($M+1$)⁺.


¹H NMR (400 MHz, acetone-*d*₆): δ 6.76 (d, J = 13.8 Hz, 1H), 6.98-7.04 (m, 4H), 7.11 (t, J = 4.8 Hz, 2H), 7.32 (t, J = 8.0 Hz, 2H), and 8.35 (dd, J = 1.6, 8.0 Hz, 1H), 10.2 (br, 2H).

Example 2**N-[2-(2,3-Dimethoxyphenoxy)-5-Fluorophenyl]-N'-(thiazol-2-yl)sulfamide**

To a solution of sulfuryl chloride (2.0 ml, 2.0 mmol, 1.0 M solution in dichloro-
5 methane) were added *p*-nitrophenol (0.55 g, 4.0 mmol) in dichloromethane and *N,N*-diisopro-
pylamine (0.71 ml, 4.0 mmol) at -78°C . The reaction mixture was stirred for 1 hour at $-$
 78°C and 2-(2,3-dimethoxyphenoxy)-5-fluoroaniline (0.52 g, 2.0 mmol) in dichloromethane (5
ml) was added. The reaction mixture was stirred for 10 min and 2-aminothiazole (0.2 g, 2.0
mmol) was added. The reaction mixture was allowed to warm up slowly to ambient
10 temperature. The mixture was concentrated under reduced pressure. The residue was
dissolved in ethyl acetate and was washed (dil. NaOH, water, brine), dried (Na_2SO_4) and
concentrated under reduced pressure. The crude product was treated with Dowex-50 acidic
resin in ethyl acetate-methanol (1:1) to remove unreacted 2-aminothiazole. The filtrate was
concentrated and the residue was purified by silica gel column chromatography
15 (ethylacetate:hexanes, from 50:50 to 90:10 as eluent system) to afford the title compound
(0.42 g, 49%).

LC-MS (m/z): 427 ($M+1$)⁺.


^1H NMR (400 MHz, acetone- d_6): δ 3.12 (br, 2H), 3.71 (s, 3H), 3.86 (s, 3H), 6.56 (dd, $J = 7.6$,
1.6 Hz, 1H), 6.66 (m, 1 H), 6.77-6.87 (m, 3H), 6.99 (t, $J = 3.6$ Hz, 1H), 7.26 (d, $J = 4.8$ Hz,
20 1H), and 7.38 (dd, $J = 10.8$, 3.2 Hz, 1H)

Example 3**1-(2-phenoxyphenyl)-5-(thiazol-2-yl)biuret**

To a solution of 2-phenoxyaniline (0.46 g, 2.50 mmol) in tetrahydrofuran (20 ml) was
5 added diisopropylethylamine (0.89 ml, 5.00 mmol) and the solution was cooled to -30°C ,
then *N*-(chlorocarbonyl)isocyanate (0.3 ml, 3.75 mmol) was slowly added. The mixture was
then allowed to warm up to the room temperature during 30 min. 2-Aminothiazole (0.375 g,
3.75 mmol) was added to the reaction mixture and stirred at room temperature for 6 hours.
The reaction mixture was concentrated under reduced pressure to afford crude product,
10 which was purified by silica gel chromatography (hexanes:ethyl acetate from 80:20 to 30:70)
to afford title compound (0.49 g, 55%) as pale yellow solid.

LC-MS (m/z): 356 ($M+1$).

^1H NMR (400 MHz, acetone- d_6): δ 6.92 (dd, $J = 1.6, 7.6$ Hz, 1H), 7.06 (m, 5H), 7.40 (m, 3H),
8.32 (d, $J = 4.8$ Hz, 1H), 8.45 (d, $J = 5.6$ Hz, 1H), 9.02 (br, 1H), 9.78 (br, 1H), and 10.46 (br,
15 1H).

Example 4**2-[[[(2-Phenoxyanilino)sulfonyl]amino]carbonyl]amino]thiazole**

To a solution of chlorosulfonyl isocyanate (0.22 ml, 2.5 mmol) in tetrahydrofuran (25
20 ml) were added 2-phenoxyaniline (0.37 g, 2.0 mmol) and DIEA (0.89 ml, 5.0 mmol) at -78°C .
The solution was stirred and slowly allowed to warmed up to 0°C . To this reaction mixture,
was added 2-amino thiazole (0.20 g, 2.0 mmol) and continued stirring at room temperature
for 3 h. The reaction mixture was concentrated under reduced pressure and the crude


product was purified by column chromatography (ethylacetate:hexanes, from 50:50 to 90:10 as eluent system) to afford the title compound (0.52 g, 66%).

LC-MS (m/z): 392 ($M+1$)⁺.

¹H NMR (400 MHz, acetone-*d*₆): δ 6.85 (dd, J = 8.8, 1.6 Hz, 1H), 6.88 (dd, J = 1.6, 8.8 Hz, 1H), 7.00-7.16 (m, 5H), 7.29-7.39 (m, 4H), 10.33 (br, 1H), 10.88 (br, 2H).

Example 5

N-(2-Phenylsulfanylphenyl)-N'-(thiazol-2-yl)urea




N-(2-Phenylsulfanylphenyl)-N'-(thiazol-2-yl)urea (116 mg, 71%) was prepared from 2-phenylsulfanylaniline (100 mg, 0.5 mmol) and 2-aminothiazole (60 mg, 0.6 mmol) following the general procedure D.

LC-MS (m/z): 329 ($M+1$)⁺.

¹H NMR (400 MHz, acetone-*d*₆): δ 7.03 (d, J = 3.6 Hz, 1H), 7.10-7.18 (m, 4H), 7.25-7.31 (m, 3H), 7.48 (m, 1H), 7.59 (dd, J = 3.6, 1.6 Hz, 1H), 8.44 (dd, J = 8.4, 1.2 Hz, 1H), 9.00 (br, 1H), 10.44 (br, 1H).

Example 6

N-(2-Phenylsulfonylphenyl)-N'-(thiazol-2-yl)urea




N-(2-Phenylsulfonylphenyl)-N'-(thiazol-2-yl)urea (98 mg, 55%) was prepared from 2-phenylsulfonylaniline (116 mg, 0.5 mmol) and 2-aminothiazole (60 mg, 0.6 mmol) following the general procedure D.

LC-MS (m/z): 361 ($M+1$)⁺.

^1H NMR (400 MHz, acetone- d_6): δ 7.07 (d, J = 3.6 Hz, 1H), 7.33-7.37 (m, 1H), 7.40 (d, J = 3.6 Hz, 1H), 7.55-7.60 (m, 2H), 7.63-7.71 (m, 2H), 7.98-8.01 (m, 2H), 8.12 (d, J = 8.0 Hz, 1H), 8.22-8.25 (m, 1H), 9.26 (br, 1H), 10.99 (br, 1H).

Example 7

5 N-(2-Benzylphenyl)-N'-(thiazol-2-yl)urea




N-(2-Benzylphenyl)-N'-(thiazol-2-yl) urea (111 mg, 72 %) was prepared from commercially available 2-benzylaniline (91 mg, 0.5 mmol) and 2-aminothiazole (60 mg, 0.6 mmol) following the general procedure D.

10 LC-MS (m/z): 311 ($M+1$) $^+$.

^1H NMR (400 MHz, acetone- d_6): δ 4.08 (s, 2H), 7.02 (d, J = 3.2 Hz, 1H), 7.06-7.14 (m, 2H), 7.14-7.19 (m, 4H), 7.23-7.29 (m, 4H), 7.96 (d, J = 8.4 Hz, 1H), 8.75 (br, 1H), 10.01 (br, 1H).

Example 8

15 N-(2-Benzoylphenyl)-N'-(thiazol-2-yl)urea




N-(2-Benzoylphenyl)-N'-(thiazol-2-yl)urea (100 mg, 63%) was prepared from 2-aminobenzophenone (97 mg, 0.5 mmol) and 2-aminothiazole (60 mg, 0.6 mmol) following the general procedure D.

LC-MS (m/z): 325 ($M+1$) $^+$.

20 ^1H NMR (400 MHz, acetone- d_6): δ 7.05 (s, 1H), 7.16 (s, 1H), 7.35 (d, J = 3.6 Hz, 1H), 7.54 (d, J = 7.2 Hz, 2H), 7.65 (s, 1H), 7.75 (s, 1H), 8.45 (s, 1H), 10.18 (br, 1H), 10.85 (br, 1H).

Example 9

N-[2-(Phenylamino)phenyl]-N'-(thiazol-2-yl)urea




5 N-[2-(Phenylamino)phenyl]-N'-(thiazol-2-yl)urea (75 mg, 49%) was prepared from 2-(N-Phenylamino)aniline (92 mg, 0.5 mmol) and 2-aminothiazole (60 mg, 0.6 mmol) following the general procedure D.

LC-MS (*m/z*): 312 (*M*+1)⁺.

¹H NMR (400 MHz, acetone-*d*₆): δ 6.75-6.79 (m, 3H), 6.86 (br, 1H), 7.01 (d, *J* = 3.6 Hz, 1H), 7.05-7.09 (m, 1H), 7.15-7.19 (m, 3H), 7.24-7.28 (m, 2H), 8.16 (d, *J* = 8.0, 1.2 Hz, 1H), 8.65
10 (br, 1H), 10.15 (br, 1H).

Example 10

N-[2-Fluoro-6-(4-methoxyphenoxy)benzyl]-N'-(thiazol-2-yl)urea




15 N-[2-Fluoro-6-(4-methoxyphenoxy)benzyl]-N'-(thiazol-2-yl)urea (0.61 g, 81%) was prepared from 2-fluoro-6-(4-methoxyphenoxy)benzylamine (0.494 g, 2.00 mmol) and 2-aminothiazole (0.20 g, 2.00 mmol) following the general procedure D.

LC-MS (*m/z*): 327 (*M*+1)⁺.

¹H NMR (400 MHz, acetone-*d*₆): δ 3.06 (br, 1H), 3.78 (s, 3H), 4.67 (d, *J* = 4.8 Hz, 2H), 6.52 (d, *J* = 8.4 Hz, 1H), 6.85 (t, *J* = 9.6 Hz, 1H), 6.95 (m, 3H), 7.05 (m, 2H), 7.24 (m, 2H), 10.06
20 (br, 1H).

Example 11

N-(2-Benzyloxyphenyl)-N'-(thiazol-2-yl)urea




N-(2-Benzyloxyphenyl)-N'-(thiazol-2-yl)urea (0.56 g, 86%) was prepared from 2-benzyloxyaniline (0.40 g, 2.00 mmol) and 2-aminothiazole (0.20 g, 2.00 mmol) following the general procedure D.

LC-MS (m/z): 327 ($M+1$).

^1H NMR (400 MHz, CDCl_3): δ 4.00 (br, 2H), 5.15 (s, 2H), 6.85 (dd, $J = 1.6, 8.8$ Hz, 2H), 6.97 (dd, $J = 1.6, 7.2$ Hz, 1H), 6.98 (dd, $J = 2.0, 6.0$ Hz, 1H), 7.37 (m, 2H), 7.46 (t, $J = 7.2$ Hz, 1H), 7.52 (d, $J = 5.6$ Hz, 2H), and 7.56 (d, $J = 6.8$ Hz, 2H).

Example 12

N-[2-(2,3,4-Trimethoxybenzyloxy)phenyl]-N'-(thiazol-2-yl)urea



2-(2,3,4-Trimethoxybenzyloxy)-1-nitrobenzene (0.46 g, 72 %) was prepared from 2,3,4-trimethoxybenzyl alcohol (0.35 ml, 2.0 mmol) and 1-fluoro-2-nitrobenzene (0.21 ml, 2.0 mmol) following the general procedure G. This was reduced to 2-(2,3,4-trimethoxybenzyloxy)aniline (0.26 g, 65 %) following the general procedure B. N-[2-(2,3,4-trimethoxybenzyloxy)phenyl]-N'-(thiazol-2-yl)urea (240 mg, 65 %) was prepared from 2-(2,3,4-trimethoxy-


benzyloxy)aniline (0.26 g, 0.9 mmol) and 2-aminothiazole (140 mg, 1.4 mmol) following the general procedure D.

LC-MS (m/z): 417 ($M+1$)⁺.

¹H NMR (400 MHz, acetone-*d*₆): δ 3.72 (s, 3H), 3.73 (s, 3H), 3.81 (s, 3H), 3.88 (s, 2H), 6.73-7.36 (m, 7H), 8.15 (t, J = 8.4 Hz, 1H), 8.90 (br, 1H), 10.10 (br, 1H).

Example 13

N-(2-Ethoxyphenyl)-N'-(thiazol-2-yl)urea




N-(2-Ethoxyphenyl)-N'-(thiazol-2-yl)urea (95 mg, 72%) was prepared from commercially available 2-ethoxyaniline (68 mg, 0.5 mmol) and 2-aminothiazole (60 mg, 0.6 mmol) following the general procedure D.

LC-MS (m/z): 265 ($M+1$)⁺.

¹H NMR (400 MHz, acetone-*d*₆): δ 1.29 (t, J = 7.0 Hz, 3H), 3.94-3.98 (q, J = 7.0 Hz, 2H), 6.74-6.88 (m, 4H), 7.27 (d, J = 5.2 Hz, 1H), 8.17 (dd, J = 1.6, 8.0 Hz, 1H), 8.42 (br, 1H), 10.92 (br, 1H).

Example 14

N-(2-Phenoxyphenyl)-N'-(pyridin-2-yl)urea




N-(2-Phenoxyphenyl)-N'-(pyridin-2-yl)urea (109 mg, 72%) was prepared from 2-phenoxyphenylisocyanate (106 mg, 0.5 mmol) and 2-aminopyridine (60 mg, 0.6 mmol) following the general procedure E.

LC-MS (m/z): 307 ($M+1$)⁺.

^1H NMR (400 MHz, $\text{DMSO}-d_6$): δ 6.88-6.91 (m, 1H), 6.96-7.08 (m, 4H), 7.15-7.19 (m, 2H), 7.32-7.36 (m, 2H), 7.65-7.69 (m, 1H), 7.88 (d, $J = 4.0$ Hz, 1H), 8.36 (d, $J = 8.0$ Hz, 1H), 9.84 (s, 1H), 11.5 (br, 1H).

Example 15

5 N-(2-Phenoxyphenyl)-N'-[(4-methoxycarbonylmethyl)thiazol-2-yl]urea




N-(2-Phenoxyphenyl)-N'-[(4-methoxycarbonylmethyl)thiazol-2-yl]urea (130 mg, 65%) was prepared from 2-phenoxyphenylisocyanate (106 mg, 0.5 mmol) and methyl 2-aminothiazole-4-acetate (104 mg, 0.6 mmol) following the general procedure E.

10 LC-MS (m/z): 401 ($M+1$) $^+$.

^1H NMR (400 MHz, $\text{acetone}-d_6$): δ 3.58 (s, 2H), 3.61 (s, 3H), 6.84-6.89 (m, 2H), 7.01-7.05 (m, 3H), 7.13-7.18 (m, 2H), 7.38-8.42 (m, 2H), 8.40 (d, $J = 8.0$ Hz, 1H), 8.91 (br, 1H), 10.12 (br, 1H).

Example 16

15 N-Methyl-N-(2-phenoxyphenyl)-N'-(thiazol-2-yl)urea



2-Phenoxyaniline (0.93 g, 5.00 mmol) and di-*tert*-butyl dicarbonate (2.18 g, 10.0 mmol) were dissolved in anhydrous dioxane (50 ml), then the reaction mixture was refluxed for 3 h. The mixture was concentrated under reduced pressure to quantitatively give 2-

phenoxy-*N*-(*t*-butoxycarbonyl)aniline (1.43 g). The product was confirmed by LC-MS and subjected to next reaction without further purification.

To a solution of 2-phenoxy-*N*-(*t*-butoxycarbonyl)aniline (1.43 g, 5.0 mmol) in anhydrous tetrahydrofuran (50 ml) was added lithium aluminumhydride (10 ml, 10.0 mmol, 1.0 M solution in tetrahydrofuran) at -10°C . The mixture was then refluxed at 65°C overnight. The reaction mixture was quenched with slow addition of MeOH (10 ml) and concentrated under reduced pressure. The reaction mixture was then poured into water (50 ml) and extracted with ethyl acetate (3x100 ml). Organic extracts were combined and washed with brine (2x100 ml) and dried over (Na_2SO_4), concentrated under reduced pressure to give *N*-methyl-2-phenoxyaniline (0.94 g, 94.0%) as a pale yellow oil. The product was confirmed by LC-MS and subjected to next reaction without further purification.


To a solution of 2-aminothiazole (0.20 g, 2.00 mmol) in dichloroethane (20 ml) was added 1,1'-carbonyldiimidazole (0.40 g, 2.5 mmol) and *N,N*-dimethylaminopyridine (0.05 g, 0.4 mmol) then the solution was refluxed at 80°C for 1 h. *N*-Methyl-2-phenoxyaniline (0.40 g, 2.00 mmol) was added to the solution. The reaction mixture was then stirred overnight at 80°C . The reaction was monitored by LC-MS and TLC, then concentrated under reduced pressure to afford crude product which subsequently was subjected to silica gel chromatography (hexanes:ethyl acetate from 80:20 to 50:50 as eluent system) to afford title product (0.48 g, 73%) as orange solid.

LC-MS (m/z): 327 ($\text{M}+1$)⁺.

^1H NMR (400 MHz, CDCl_3): δ 3.30 (s, 3H), 6.84 (d, $J = 4.8$ Hz, 1H), 6.97 (m, 3H), 7.16 (m, 2H), 7.29 (m, 5H), and 7.74 (br, 1H).

Example 17

N-isopropyl-*N*-(2-phenoxyphenyl)-*N'*-(thiazol-2-yl)urea



To a solution of 2-phenoxyaniline (0.93 g, 5.0 mmol) in dichloroethane (50 ml) was added anhydrous acetone (0.73 ml, 10.0 mmol) and acetic acid (0.1 ml, 2.0 mmol). The mixture was stirred for 30 min, and sodium triacetoxymethylborohydride (3.18 g, 15.0 mmol) was added in one portion. The reaction mixture was stirred overnight at ambient temperature. The

reaction mixture was quenched by slow addition of MeOH (10 ml) and concentrated under reduced pressure. The residue was poured into water (50 ml) and extracted with ethyl acetate (3x100 ml). Organic extracts were combined and washed with brine (2x100 ml) and dried (Na₂SO₄), concentrated under reduced pressure to give *N*-isopropyl-2-phenoxyaniline (1.05 g, 91%) as colorless oil.


To a solution of 2-aminothiazole (0.20 g, 2.00 mmol) in dichloroethane (20 ml) was added 1,1'-carbonyldiimidazole (0.40 g, 2.5 mmol) and *N,N*-dimethylaminopyridine (0.05 g, 0.4 mmol) then the solution was refluxed at 80°C for 1 h. *N*-Isopropyl-2-phenoxyaniline (0.46 g, 2.0 mmol) was added and the reaction mixture was then stirred overnight at 80°C. The reaction mixture was concentrated under reduced pressure to afford crude product, which was purified by silica gel chromatography (hexanes:ethyl acetate from 80:20 to 50:50 as eluent system) to afford title product (0.56 g, 79%).

LC-MS (*m/z*): 355 (M+1)⁺.

¹H NMR (400 MHz, acetone-*d*₆): δ 1.22 (d, *J* = 6.4 Hz, 6H), 4.81 (m, 1H), 6.91 (m, 2H), 7.10-7.24 (m, 5H), 7.39 (m, 4H), and 9.00 (br, 1H).

Example 18


N-[2-(4-Methoxyphenoxy)phenyl]-*N'*-(thiazol-2-yl)urea



2-(4-Methoxyphenoxy)-1-nitrobenzene (0.98 g, 80 %) was prepared from 4-methoxyphenol (0.62 g, 5.0 mmol) and 1-fluoro-2-nitrobenzene (0.71 g, 5.0 mmol) following the general procedure A. This was reduced to 2-(4-methoxyphenoxy)aniline (0.32 g, 60%, 2.5 mmol scale) following the general procedure B. *N*-[2-(4-Methoxyphenoxy)phenyl]-*N'*-(thiazol-2-yl)urea (256 mg, 75%) was prepared from 2-(4-methoxyphenoxy)aniline (215 mg, 1.0 mmol) and 2-aminothiazole (100 mg, 1.0 mmol) following the general procedure D.

LC-MS (*m/z*): 343 (M+1)⁺.


¹H NMR (400 MHz, acetone-*d*₆): δ 3.80 (s, 3H), 6.80 (d, *J* = 8.0 Hz, 1H), 6.98-7.05 (m, 6H), 7.10 (m, 1H), 7.32 (d, *J* = 3.6 Hz, 1H), 8.40 (d, *J* = 8.0 Hz, 1H), 8.90 (br, 1H), 10.23 (br, 1H).

Example 19**N-[2-(4-Fluorophenoxy)phenyl]-N'-(thiazol-2-yl)urea**

2-(4-Fluorophenoxy)-1-nitrobenzene (0.87 g, 75%) was prepared from 4-fluorophenol (0.62 g, 5.5 mmol) and 1-fluoro-2-nitrobenzene (0.71 g, 5.0 mmol) following the general procedure A. This was reduced to 2-(4-fluorophenoxy)aniline (0.51 g, 68%) following general procedure B. N-[2-(4-Fluorophenoxy)phenyl]-N'-(thiazol-2-yl)urea (118 mg, 72%) was prepared from 2-(4-fluorophenoxy)aniline (102 mg, 0.5 mmol) and 2-aminothiazole (60 mg, 0.6 mmol) following the general procedure D.

LC-MS (m/z): 331 ($M+1$)⁺.

¹H NMR (400 MHz, acetone-*d*₆): δ 6.89 (dd, J = 8.4, 1.2 Hz, 1H), 7.01-7.11 (m, 4H), 7.14-7.29 (m, 3H), 7.31 (d, J = 3.6 Hz, 1H), 8.42 (dd, J = 8.0, 1.6 Hz, 1H), 9.12 (br, 1H), 10.19 (br, 1H).

Example 20**N-[2-(4-Chlorophenoxy)phenyl]-N'-(thiazol-2-yl)urea**


2-(4-Chlorophenoxy)-1-nitrobenzene (0.88 g, 71%) was prepared from 4-chlorophenol (0.70 g, 5.5 mmol) and 1-fluoro-2-nitrobenzene (0.71 g, 5.0 mmol) following the general procedure A. This was reduced to 2-(4-chlorophenoxy)aniline (0.50 g, 65%) following general procedure B. N-[2-(4-Chlorophenoxy)phenyl]-N'-(thiazol-2-yl)urea (106 mg, 62%) was prepared from 2-(4-chlorophenoxy)aniline (109 mg, 0.5 mmol) and 2-aminothiazole (60 mg, 0.6 mmol) following the general procedure D.

LC-MS (m/z): 347 ($M+1$)⁺.

¹H NMR (400 MHz, acetone-*d*₆): δ 6.98 (d, *J* = 7.6 Hz, 1H), 7.04-7.09 (m, 4H), 7.19-7.23 (m, 1H), 7.31 (d, *J* = 3.6 Hz, 1H), 7.39-7.43 (m, 2H), 8.44 (dd, *J* = 8.4, 1.6 Hz, 1H), 8.90 (br, 1H), 10.13 (br, 1H).

Example 21

5 N-[2-(4-Cyanophenoxy)phenyl]-N'-thiazolylurea




2-(4-Cyanophenoxy)-1-nitrobenzene (0.82 g, 69%) was prepared from 4-cyanophenol (0.66 g, 5.5 mmol) and 1-fluoro-2-nitrobenzene (0.71 g, 5.0 mmol) following the general procedure A. This was reduced to 2-(4-cyanophenoxy)aniline (0.47 g, 65%) following
10 general procedure B. N-[2-(4-Cyanophenoxy)phenyl]-N'-thiazolylurea (110 mg, 65%) was prepared from 2-(4-cyanophenoxy)aniline (105 mg, 0.5 mmol) and 2-aminothiazole (60 mg, 0.6 mmol) following the general procedure D.

LC-MS (*m/z*): 338 (*M*+1)⁺.

¹H NMR (400 MHz, acetone-*d*₆): δ 6.04 (d, *J* = 8.0 Hz, 1H), 7.13-7.18 (m, 4H), 7.28-7.33 (m,
15 2H), 7.78-7.81 (m, 2H), 8.47 (d, *J* = 8.0 Hz, 1H), 8.95 (br, 1H), 10.43 (br, 1H).

Example 22

N-[2-(4-Methoxycarbonylphenoxy)phenyl]-N'-(thiazol-2-yl)urea



2-(4-Methoxycarbonylphenoxy)-1-nitrobenzene (0.79 g, 58%) was prepared from
20 methyl 4-hydroxybenzoate (0.84 g, 5.5 mmol) and 1-fluoro-2-nitrobenzene (0.71 g, 5.0 mmol) following the general procedure A. This was reduced to 2-(4-methoxycarbonylphenoxy)-


aniline (0.46 g, 66%) following general procedure B. N-[2-(4-Methoxycarbonylphenoxy)-phenyl]-N'-(thiazol-2-yl)urea (100 mg, 55%) was prepared from 2-(4-methoxycarbonylphenoxy)aniline (122 mg, 0.5 mmol) and 2-aminothiazole (60 mg, 0.6 mmol) following the general procedure D.

5 LC-MS (m/z): 371 ($M+1$)⁺.

¹H NMR (400 MHz, acetone- d_6): δ 3.85 (s, 3H), 7.03 (d, J = 3.6 Hz, 1H), 7.07–7.14 (m, 4H), 7.25–7.29 (m, 2H), 8.01–4.04 (m, 2H), 8.46 (dd, J = 8.1, 1.2 Hz, 1H), 8.76 (br, 1H), 10.07 (br, 1H).

Example 23


10 N-[2-(4-Isopropylphenoxy)phenyl]-N'-(thiazol-2-yl)urea



2-(4-Isopropylphenoxy)-1-nitrobenzene (0.95 g, 75 %) was prepared from 4-isopropylphenol (0.68 g, 5.0 mmol) and 1-fluoro-2-nitrobenzene (0.71 g, 5.0 mmol) following the general procedure A. This was reduced to 2-(4-isopropylphenoxy)aniline (0.34 g, 60 %, 2.5 mmol scale) following general procedure B. N-[2-(4-Isopropylphenoxy)phenyl]-N'-(thiazol-2-yl)urea (247 mg, 70 %) was prepared from 2-(4-isopropylphenoxy)aniline (227 mg, 1.0 mmol) and 2-aminothiazole (100 mg, 1.0 mmol) following the general procedure D.

LC-MS (m/z): 355 ($M+1$)⁺.


¹H NMR (400 MHz, acetone- d_6): δ 1.17 (d, J = 7.2 Hz, 3H), 1.22 (d, J = 6.8 Hz, 3H), 2.88 (m, 1H), 6.74 (d, J = 6.6 Hz, 1H), 6.88 (d, J = 8.0 Hz, 1H), 6.94–7.15 (m, 4H), 7.28 (m, 2H), 8.43 (d, J = 8.2 Hz, 1H), 10.15 (br, 2H).

Example 24**N-[2-(3,4-Difluorophenoxy)phenyl]-N'-(thiazol-2-yl)urea**

2-(3,4-Difluorophenoxy)-1-nitrobenzene (0.76 g, 60 %) was prepared from 3,4-difluorophenol (0.65 g, 5.5 mmol) and 1-fluoro-2-nitrobenzene (0.71 g, 5.0 mmol) following the general procedure A. This was reduced to 2-(3,4-difluorophenoxy)aniline (0.33 g, 60 %, 2.5 mmol scale) following general procedure B. N-[2-(3,4-Difluorophenoxy)phenyl]-N'-(thiazol-2-yl)urea (312 mg, 60 %) was prepared from 2-(3,4-difluorophenoxy)aniline (330 mg, 1.5 mmol) and 2-aminothiazole (150 mg, 1.5 mmol) following the general procedure D.

LC-MS (*m/z*): 349 (*M*+1)⁺.

¹H NMR (400 MHz, acetone-*d*₆): δ 6.88 (m, 1H), 7.01 (d, *J* = 8.0 Hz, 1H), 7.04-7.12 (m, 4H), 7.22 (m, 1H), 7.30-7.42 (m, 2H), 8.43 (d, *J* = 8.2 Hz, 1H), 10.16 (br, 2H).

Example 25**N-[2-(3,4-Dichlorophenoxy)phenyl]-N'-(thiazol-2-yl)urea**


2-(3,4-Dichlorophenoxy)-1-nitrobenzene (1.15 g, 81%) was prepared from 3,4-dichlorophenol (0.9 g, 5.5 mmol) and 1-fluoro-2-nitrobenzene (0.71 g, 5.0 mmol) following the general procedure A. This was reduced to 2-(3,4-dichlorophenoxy)aniline (0.69 g, 68%) following general procedure B. N-[2-(3,4-Dichlorophenoxy)phenyl]-N'-(thiazol-2-yl)urea (129 mg, 68%) was prepared from 2-(3,4-dichlorophenoxy)aniline (127 mg, 0.5 mmol) and 2-aminothiazole (60 mg, 0.6 mmol) following the general procedure D.

LC-MS (*m/z*): 382 (*M*+1)⁺.

¹H NMR (400 MHz, acetone-*d*₆): δ 6.99-7.13 (m, 4H), 7.24-7.28 (m, 2H), 7.30 (d, *J* = 3.6 Hz, 1H), 7.57 (d, *J* = 9.2 Hz, 1H), 8.44 (dd, *J* = 8.4, 1.2 Hz, 1H), 9.20 (br, 1H), 10.09 (br, 1H).

Example 26

N-[2-(4-Chloro-3-methylphenoxy)phenyl]-N'-(thiazol-2-yl)urea




5 N-[2-(4-Chloro-3-methylphenoxy)phenyl]-N'-(thiazol-2-yl)urea (233 mg, 65 %) was prepared from 2-(4-chloro-3-methylphenoxy)aniline (233 mg, 1.0 mmol) and 2-aminothiazole (100 mg, 1.0 mmol) following the general procedure D.

LC-MS (*m/z*): 361 (*M*+1)⁺.

¹H NMR (400 MHz, acetone-*d*₆): δ 2.36 (s, 3H), 6.94-6.98 (m, 2H), 7.05 (m, 4H), 7.19 (m, 1H), 7.31 (d, *J* = 4.0 Hz, 1H), 7.37 (d, *J* = 8.4 Hz, 1H), 8.43 (d, *J* = 8.4 Hz, 1H), 10.23 (br, 2H).

Example 27

N-[2-(3,4-Dimethoxyphenoxy)phenyl]-N'-(thiazol-2-yl)urea



15 2-(3,4-Dimethoxyphenoxy)-1-nitrobenzene (0.93 g, 68%) was prepared from 3,4-dimethoxyphenol (0.85 g, 5.5 mmol) and 1-fluoro-2-nitrobenzene (0.71 g, 5.0 mmol) following the general procedure A. This was reduced to 2-(3,4-dimethoxyphenoxy)aniline (0.51 g, 62%) following general procedure B. N-[2-(3,4-Dimethoxyphenoxy)phenyl]-N'-(thiazol-2-yl)urea (128 mg, 69%) was prepared from 2-(3,4-dimethoxyphenoxy)aniline (123 mg, 0.5 mmol) and 2-aminothiazole (60 mg, 0.6 mmol) following the general procedure D.


20 LC-MS (*m/z*): 373 (*M*+1)⁺.

¹H NMR (400 MHz, acetone-*d*₆): δ 3.80 (s, 6H), 6.54 (dd, *J* = 8.4, 2.4 Hz, 1H), 6.78 (d, *J* = 2.4 Hz, 1H), 6.82 (d, *J* = 8.0 Hz, 1H), 6.97 (d, *J* = 8.0 Hz, 1H), 6.99-7.01 (m, 1H), 7.04 (d, *J* = 3.6

Hz, 1H), 7.07-7.12 (m, 1H), 7.33-7.34 (d, $J = 3.6$ Hz, 1H), 8.39 (dd, $J = 8.0, 1.2$ Hz, 1H), 8.90 (br, 1H), 10.23 (br, 1H).

Example 28

N-[2-(3,4-Methylenedioxyphenoxy)phenyl]-N'-(thiazol-2-yl)urea



5

2-(3,4-Methylenedioxyphenoxy)-1-nitrobenzene (0.75 g, 58%) was prepared from 3,4-methylenedioxyphenol (0.76 g, 5.5 mmol) and 1-fluoro-2-nitrobenzene (0.71 g, 5.0 mmol) following the general procedure A. This was reduced to 2-(3,4-methylenedioxyphenoxy)-aniline (0.47 g, 71%) following general procedure B. N-[2-(3,4-Methylenedioxyphenoxy)-phenyl]-N'-(thiazol-2-yl)urea (120 mg, 68%) was prepared from 2-(3,4-methylenedioxyphenoxy)aniline (115 mg, 0.5 mmol) and 2-aminothiazole (60 mg, 0.6 mmol) following the general procedure D.


LC-MS (m/z): 357 ($M+1$)⁺.

¹H NMR (400 MHz, acetone- d_6): δ 6.06 (s, 2 H), 6.51 (dd, $J = 8.4, 2.4$ Hz, 1H), 6.67 (d, $J = 2.4$ Hz, 1H), 6.84-6.87 (m, 2H), 6.99-7.05 (m, 2H), 7.10-7.14 (m, 1H), 7.32-7.33 (d, $J = 3.6$ Hz, 1H), 8.39 (dd, $J = 8.0, 1.2$ Hz, 1H), 8.95 (br, 1H), 10.22 (br, 1H).

15

Example 29

N-[2-(2,4-Dichlorophenoxy)phenyl]-N'-(thiazol-2-yl)urea



N-[2-(2,4-Dichlorophenoxy)phenyl]-N'-(thiazol-2-yl)urea (125 mg, 67%) was prepared from 2-(2,4-dichlorophenoxy)aniline (127 mg, 0.5 mmol) and 2-aminothiazole (60 mg, 0.6 mmol) following the general procedure D.


20

LC-MS (m/z): 382 ($M+1$)⁺.

¹H NMR (400 MHz, acetone-*d*₆): δ 6.86 (d, J = 8.0 Hz, 1H) 7.03-7.08 (m, 3H), 7.19-7.23 (m, 1H), 7.30 (d, J = 4.0 Hz, 1H), 7.38 (dd, J = 8.8, 2.8 Hz, 1H), 7.65 (d, J = 2.4 Hz, 1H), 8.43 (dd, J = 8.0, 1.2 Hz, 1H), 8.87 (br, 1H), 10.16 (br, 1H).

5 Example 30

N-[2-(2,4-Difluorophenoxy)phenyl]-N'-(thiazol-2-yl)urea




2-(2,4-Difluorophenoxy)-1-nitrobenzene (0.95 g, 76%) was prepared from 2,4-difluorophenol (0.72 g, 5.5 mmol) and 1-fluoro-2-nitrobenzene (0.71 g, 5.0 mmol) following the
10 general procedure A. This was reduced to 2-(2,4-difluorophenoxy)aniline (0.53 g, 63%) following general procedure B. N-[2-(2,4-Difluorophenoxy)phenyl]-N'-(thiazol-2-yl)urea (120 mg, 69%) was prepared from 2-(2,4-difluorophenoxy)aniline (110 mg, 0.5 mmol) and 2-aminothiazole (60 mg, 0.6 mmol) following the general procedure D.

LC-MS (m/z): 349 ($M+1$)⁺.

15 ¹H NMR (400 MHz, acetone-*d*₆): δ 6.83 (m, 1H), 7.00-7.30 (m, 6H), 7.32 (d, J = 3.6 Hz, 1H), 8.42 (dd, J = 8.0, 2.0 Hz, 1H), 9.00 (br, 1H), 10.19 (br, 1H).

Example 31

N-[2-(4-Fluoro-2-methoxyphenoxy)phenyl]-N'-(thiazol-2-yl)urea



20 2-(4-Fluoro-2-methoxyphenoxy)-1-nitrobenzene (0.88 g, 67%) was prepared from 4-fluoro-2-methoxyphenol (0.78 g, 5.5 mmol) and 1-fluoro-2-nitrobenzene (0.71 g, 5.0 mmol) following the general procedure A. This was reduced to 2-(4-fluoro-2-methoxyphenoxy)-aniline (0.60 g, 78%) following general procedure C. N-[2-(4-Fluoro-2-methoxyphenoxy)-


phenyl]-N'-(thiazol-2-yl)urea (127 mg, 71%) was prepared from 2-(4-fluoro-2-methoxyphenoxy)aniline (116 mg, 0.5 mmol) and 2-aminothiazole (60 mg, 0.6 mmol) following the general procedure D.

LC-MS (*m/z*): 361 (*M*+1)⁺.

- 5 ¹H NMR (400 MHz, acetone-*d*₆): δ 3.82 (s, 3H), 6.61 (dd, *J* = 8.0, 2.8 Hz, 1H), 6.74-6.79 (m, 1H), 6.89-6.94 (m, 1H), 7.00-7.06 (m, 3H), 7.16 (dd, *J* = 8.8, 5.6 Hz, 1H), 7.33 (d, *J* = 3.6 Hz, 1H), 8.43 (dd, *J* = 8.4, 1.6 Hz, 1H), 8.85 (br, 1H), 10.29 (br, 1H).

Example 32

N-[2-(4-Methoxy-2-methoxycarbonylphenoxy)phenyl]-N'-(thiazol-2-yl)urea



10

2-[4-Methoxy-2-methoxycarbonylphenoxy]-1-nitrobenzene (0.78 g, 52%) was prepared from methyl 5-methoxysalicylate (1.0 g, 5.5 mmol) and 1-fluoro-2-nitrobenzene (0.71 g, 5.0 mmol) following the general procedure A. This was reduced to 2-[4-methoxy-2-(methoxycarbonyl)phenoxy]aniline (0.47 g, 68%) following general procedure B.


- 15 N-[2-(4-Methoxy-2-methoxycarbonylphenoxy)phenyl]-N'-(thiazol-2-yl)urea (130 mg, 66%) was prepared from 2-[4-methoxy-2-(methoxycarbonyl)phenoxy]aniline (136 mg, 0.5 mmol) and 2-aminothiazole (60 mg, 0.6 mmol) following the general procedure D.

LC-MS (*m/z*): 401 (*M*+1)⁺.

- 20 ¹H NMR (400 MHz, acetone-*d*₆): δ 3.69 (s, 3H), 3.84 (s, 3H), 6.61 (dd, *J* = 8.0, 1.6 Hz, 1H), 6.90-6.95 (m, 1H), 7.03-7.09 (m, 2H), 7.16-7.25 (m, 2H), 7.25-7.39 (m, 1H), 7.43-7.44 (m, 1H), 8.37 (dd, *J* = 8.4, 1.6 Hz, 1H), 8.96 (br, 1H), 10.26 (br, 1H).

Example 33

N-[2-(3-Methoxyphenoxy)phenyl]-N'-(thiazol-2-yl)urea




2-(3-Methoxyphenoxy)-1-nitrobenzene (0.84 g, 69%) was prepared from 3-methoxy-
5 phenol (0.68 g, 5.5 mmol) and 1-fluoro-2-nitrobenzene (0.71 g, 5.0 mmol) following the
general procedure A. This was reduced to 2-(3-methoxyphenoxy)aniline (0.51 g, 70%)
following general procedure B. N-[2-(3-Methoxyphenoxy)phenyl]-N'-(thiazol-2-yl)urea (136
mg, 68%) was prepared from 2-(3-methoxyphenoxy)aniline (108 mg, 0.5 mmol) and 2-amino-
thiazole (60 mg, 0.6 mmol) following the general procedure D.

10 LC-MS (*m/z*): 343 (*M*+1)⁺.

¹H NMR (400 MHz, acetone-*d*₆): δ 3.78 (s, 3H), 6.55-6.62 (m, 2H), 6.71-6.73 (m, 1H), 6.95
(dd, *J* = 8.0, 1.6 Hz, 1H), 7.03-7.07 (m, 2H), 7.16-7.20 (m, 1H), 7.27 (d, *J* = 8.4 Hz, 1H), 7.31
(d, *J* = 4.0 Hz, 1H), 8.43 (dd, *J* = 8.4, 1.6 Hz, 1H), 8.84 (br, 1H), 10.37 (br, 1H).

Example 34

15 N-[2-(3-Fluorophenoxy)phenyl]-N'-(thiazol-2-yl)urea



2-(3-Fluorophenoxy)-1-nitrobenzene (0.85 g, 73%) was prepared from 3-fluoro-
phenol (0.62 g, 5.5 mmol) and 1-fluoro-2-nitrobenzene (0.71 g, 5.0 mmol) following the
general procedure A. This was reduced to 2-(3-fluorophenoxy)aniline (0.50 g, 68%) following
20 general procedure B. N-[2-(3-Fluorophenoxy)phenyl]-N'-(thiazol-2-yl)urea (110 mg, 68%)


was prepared from 2-(3-fluorophenoxy)aniline (102 mg, 0.5 mmol) and 2-aminothiazole (60 mg, 0.6 mmol) following the general procedure D.

LC-MS (m/z): 331 ($M+1$)⁺.

¹H NMR (400 MHz, acetone-*d*₆): δ 6.80-6.84 (m, 2H), 6.85-6.93 (m, 1H), 7.03-7.07 (m, 2H),
5 7.08-7.13 (m, 1H), 7.30 (d, J = 3.6 Hz, 1H), 7.38-7.45 (m, 1H), 8.45 (dd, J = 8.0, 1.2 Hz, 1H),
9.06 (br, 1H), 10.1 (br 1H).

Example 35

N-[2-(3-Trifluoromethylphenoxy)phenyl]-N'-(thiazol-2-yl)urea




10 2-[3-(Trifluoromethyl)phenoxy]-1-nitrobenzene (0.92 g, 65%) was prepared from 3-hydroxybenzotrifluoride (0.89 g, 5.5 mmol) and 1-fluoro-2-nitrobenzene (0.71 g, 5.0 mmol) following the general procedure A. This was reduced to 2-[3-(trifluoromethyl)phenoxy]aniline (0.56 g, 68%) following general procedure B. N-[2-(3-Trifluoromethylphenoxy)phenyl]-N'-(thiazol-2-yl)urea (120 mg, 62%) was prepared from 2-[3-(trifluoromethyl)phenoxy]aniline
15 (127 mg, 0.5 mmol) and 2-aminothiazole (60 mg, 0.6 mmol) following the general procedure D.

LC-MS (m/z): 381 ($M+1$)⁺.

¹H NMR (400 MHz, acetone-*d*₆): δ 7.04-7.13 (m, 3H), 7.24-7.29 (m, 3H), 7.36 (s, 1H), 7.47
(d, J = 7.6 Hz, 1H), 7.63 (t, J = 8.0 Hz, 1H), 8.46 (dd, J = 8.0, 1.2 Hz, 1H), 8.95 (br, 1H),
20 10.08 (br, 1H).

Example 36

N-[2-(2-Methylphenoxy)phenyl]-N'-(thiazol-2-yl)urea




N-[2-(2-Methylphenoxy)phenyl]-N'-(thiazol-2-yl)urea (110 mg, 68%) was prepared from 2-(2-methylphenoxy)aniline (100 mg, 0.5 mmol) and 2-aminothiazole (60 mg, 0.6 mmol) following the general procedure D.

LC-MS (*m/z*): 327 (*M*+1)⁺.

- 5 ¹H NMR (400 MHz, acetone-*d*₆): δ 2.25 (s, 3H), 6.68 (d, *J* = 8.4 Hz, 1H), 6.89 (d, *J* = 8.0 Hz, 1H), 6.96-7.00 (m, 1H), 7.04-7.13 (m, 3H), 7.20-7.34 (m, 3H), 8.42 (dd, *J* = 8.0, 1.6 Hz, 1H), 8.95 (br, 1H), 10.25 (br, 1H).

Example 37

N-[2-(2-Methoxyphenoxy)phenyl]-N'-(thiazol-2-yl)urea



10

2-(2-Methoxyphenoxy)-1-nitrobenzene (0.99 g, 81%) was prepared from 2-methoxyphenol (0.68 g, 5.5 mmol) and 1-fluoro-2-nitrobenzene (0.71 g, 5.0 mmol) following the general procedure A. This was reduced to 2-(2-methoxyphenoxy)aniline (0.63 g, 73%) following general procedure B. N-2-(2-Methoxyphenoxy)phenyl-N'-(thiazol-2-yl)urea (110 g, 65%) was prepared from 2-(2-methoxyphenoxy)aniline (108 mg, 0.5 mmol) and 2-amino-

15


thiazole (60 mg, 0.6 mmol) following the general procedure D.

LC-MS (*m/z*): 343 (*M*+1)⁺.

¹H NMR (400 MHz, acetone-*d*₆): δ 3.78 (s, 3H), 6.61 (d, *J* = 8.0 Hz, 1H), 6.91-7.33 (m, 8H), 8.37 (d, *J* = 8.0 Hz, 1H), 9.13 (br, 1H), 10.31 (br, 1H).

20 Example 38

N-[2-(2-Isopropoxyphenoxy)phenyl]-N'-(thiazol-2-yl)urea




2-(2-Isopropoxyphenoxy)-1-nitrobenzene (0.90 g, 69%) was prepared from 2-isopropoxyphenol (0.84 g, 5.5 mmol) and 1-fluoro-2-nitrobenzene (0.71 g, 5.0 mmol) following the general procedure A. This was reduced to 2-(2-methoxyphenoxy)aniline (0.65 g, 81%) following general procedure B. N-[2-(2-Isopropoxyphenoxy)phenyl]-N'-(thiazol-2-yl)urea (120 mg, 65%) was prepared from 2-(2-isopropoxyphenoxy)aniline (122 mg, 0.5 mmol) and 2-aminothiazole (60 mg, 0.6 mmol) following the general procedure D.

LC-MS (*m/z*): 371 (*M*+1)⁺.

¹H NMR (400 MHz, acetone-*d*₆): δ 1.07 (d, *J* = 6.0 Hz, 6H), 4.54-4.59 (m, 1H), 6.64 (dd, *J* = 8.0, 1.6 Hz, 1H), 6.88-6.92 (m, 1H), 6.97-7.06 (m, 3H), 7.12-7.22 (m, 3H), 7.33 (d, *J* = 3.6 Hz, 1H), 8.38 (dd, *J* = 8.0, 1.6 Hz, 1H), 8.86 (br, 1H), 10.3 (br, 1H).

Example 39

N-[2-(2-Fluorophenoxy)phenyl]-N'-(thiazol-2-yl)urea




2-(2-Fluorophenoxy)-1-nitrobenzene (0.94 g, 81%) was prepared from 2-fluorophenol (0.62 g, 5.5 mmol) and 1-fluoro-2-nitrobenzene (0.71 g, 5.0 mmol) following the general procedure A. This was reduced to 2-(2-fluorophenoxy)aniline (0.59 g, 72%) following general procedure B. N-[2-(2-Fluorophenoxy)phenyl]-N'-(thiazol-2-yl)urea (110 mg, 68%) was prepared from 2-(2-fluorophenoxy)aniline (102 mg, 0.5 mmol) and 2-aminothiazole (60 mg, 0.6 mmol) following the general procedure D.

LC-MS: 331 (*M*+1)⁺.

¹H NMR (400 MHz, acetone-*d*₆): δ 6.84 (d, *J* = 8.0 Hz, 1H), 7.05-7.08 (m, 1H), 7.01-7.06 (m, 2H), 7.12-7.17 (m, 2H), 7.22-7.25 (m, 2H), 7.29-7.36 (m, 2H), 8.43 (dd, *J* = 8.0, 1.6 Hz, 1H), 8.95 (br, 1H), 10.17 (br, 1H).

Example 40


N-[2-(2-Methylsulfonylphenoxy)phenyl]-N'-(thiazol-2-yl)urea



- 2-(2-Methylsulfonylphenoxy)-1-nitrobenzene (0.88 g, 68%) was prepared from 2-hydroxythioanisole (0.77 g, 5.5 mmol) and 1-fluoro-2-nitrobenzene (0.71 g, 5.0 mmol) following the general procedure A. This was reduced to 2-(2-methylsulfonylphenoxy)aniline (0.53 g, 68%) following general procedure B. N-[2-(2-Methylsulfonylphenoxy)phenyl]-N'-(thiazol-2-yl)urea (115 mg, 65%) was prepared from 2-(2-methylsulfonylphenoxy)aniline (115 mg, 0.5 mmol) and 2-aminothiazole (60 mg, 0.6 mmol) following the general procedure D.
- LC-MS (m/z): 359 ($M+1$)⁺.
¹H NMR (400 MHz, acetone-*d*₆): δ 2.42 (s, 3H), 6.68 (dd, J = 8.0, 1.2 Hz, 1H), 6.92 (d, J = 8.0 Hz, 1H), 6.95-6.99 (m, 1H), 7.04 (d, J = 3.6 Hz, 1H), 7.09-7.14 (m, 1H), 7.17-7.26 (m, 2H), 7.31 (d, J = 3.6 Hz, 1H), 7.38 (dd, J = 7.6, 1.6 Hz, 1H), 8.42 (dd, J = 8.0, 1.6 Hz, 1H), 8.95 (br, 1H), 10.26 (br, 1H).

Example 41

N-[2-(2-Methylsulfonylphenoxy)phenyl]-N'-thiazolylurea



- To a solution of 2-(2-methylsulfonylphenoxy)-1-nitrobenzene (1.3 g, 5.0 mmol) in dichloromethane (30 ml) at 0°C was added *m*CPBA (70%, 3.68 g, 15 mmol), in portions during 10 min. The contents were stirred for 2 h at room temperature. The precipitate was filtered off and the mother liquor was washed thrice with 10% aq Na₂S₂O₇. The organic layer was washed (dil NaOH, water, brine), dried and concentrated to obtain 2-(2-methylsulfonylphenoxy)-1-nitrobenzene (1.0 g, 68%). This was reduced to 2-(2-methylsulfonylphenoxy)-


aniline (0.68 g, 76%) following general procedure B. N-[2-(2-Methylsulfonylphenoxy)phenyl]-N'-(thiazol-2-yl)urea (105 mg, 55%) was prepared from 2-(2-methylsulfonylphenoxy)aniline (132 mg, 0.5 mmol) and 2-aminothiazole (60 mg, 0.6 mmol) following the general procedure D.

5 LC-MS (*m/z*): 391 (*M*+1)⁺.

¹H NMR (400 MHz, acetone-*d*₆): δ 3.44 (s, 3H), 7.02 (d, *J* = 3.6 Hz, 1H), 7.06 (d, *J* = 8.4 Hz, 1H), 7.12-7.20 (m, 2H), 7.27-7.36 (m, 2H), 7.38-7.40 (m, 1H), 7.67-7.71 (m, 1H), 8.02 (dd, *J* = 8.0, 1.6 Hz, 1H), 8.42 (dd, *J* = 8.4, 1.2 Hz, 1H), 9.20 (br, 1H), 9.78 (br, 1H).

Example 42


10 N-[2-(2-Trifluoromethylphenoxy)phenyl]-N'-(thiazol-2-yl)urea



2-[2-(Trifluoromethyl)phenoxy]-1-nitrobenzene (0.92 g, 65%) was prepared from 2-hydroxybenzotrifluoride (0.89 g, 5.5 mmol) and 1-fluoro-2-nitrobenzene (0.71 g, 5.0 mmol) following the general procedure A. This was reduced to 2-[2-(trifluoromethyl)phenoxy]aniline (0.56 g, 68%) following general procedure B. N-[2-(2-Trifluoromethylphenoxy)phenyl]-N'-(thiazol-2-yl)urea (115 mg, 61%) was prepared from 2-[2-(trifluoromethyl)phenoxy]aniline (127 mg, 0.5 mmol) and 2-aminothiazole (60 mg, 0.6 mmol) following the general procedure D.

LC-MS (*m/z*): 381 (*M*+1)⁺.

20 ¹H NMR (400 MHz, acetone-*d*₆): δ 6.91 (dd, *J* = 8.0, 1.6 Hz, 1H), 7.01-7.06 (m, 2H), 7.08-7.11 (m, 1H), 7.22-7.26 (m, 2H), 7.34 (t, *J* = 7.6 Hz, 1H), 7.65 (t, *J* = 7.6 Hz, 1H), 7.81 (dd, *J* = 8.0, 1.2 Hz, 1H), 8.45 (dd, *J* = 8.4, 1.6 Hz, 1H), 9.00 (br, 1H), 10.2 (br, 1H).

Example 43**N-[2-(2,6-Dimethoxyphenoxy)phenyl]-N'-(thiazol-2-yl)urea**


2-(2,6-Dimethoxyphenoxy)-1-nitrobenzene (0.85 g, 63%) was prepared from 2,6-dimethoxyphenol (0.85 g, 5.5 mmol) and 1-fluoro-2-nitrobenzene (0.71 g, 5.0 mmol) following the general procedure A. This was reduced to 2-(2,6-dimethoxyphenoxy)aniline (0.51 g, 68%) following general procedure C. N-[2-(2,6-Dimethoxyphenoxy)phenyl]-N'-(thiazol-2-yl)-urea (117 g, 63%) was prepared from 2-(2,6-dimethoxyphenoxy)aniline (123 mg, 0.5 mmol) and 2-aminothiazole (60 mg, 0.6 mmol) following the general procedure D.

LC-MS (m/z): 373 ($M+1$)⁺.

¹H NMR (400 MHz, acetone-*d*₆): δ 3.76 (s, 3 H), 6.49 (dd, J = 8.4, 1.2 Hz, 1H), 6.80 (d, J = 8.4 Hz, 2H), 6.84-6.88 (m, 1H), 6.95-6.99 (m, 1H), 7.04 (d, J = 3.2 Hz, 1H), 7.22 (t, J = 8.4 Hz, 1H), 7.35 (d, J = 3.6 Hz, 1H), 8.33 (dd, J = 8.0, 1.6 Hz, 1H), 8.85 (br, 1H), 10.40 (br, 1H).

Example 44

N-[2-(2,6-difluorophenoxy)phenyl]-N'-(thiazol-2-yl)urea




2-(2,6-difluorophenoxy)-1-nitrobenzene (0.44 g, 70 %) was prepared from 2,6-difluorophenol (0.32 g, 2.5 mmol) and 1-fluoro-2-nitrobenzene (0.36 g, 2.5 mmol) following the general procedure A. This was reduced to 2-(2,6-difluorophenoxy)aniline (0.24 g, 60 %) following general procedure B. N-[2-(2,6-difluorophenoxy)phenyl]-N'-(thiazol-2-yl)urea (133 mg, 60 %) was prepared from 2-(2,6-difluorophenoxy)aniline (206 mg, 0.9 mmol) and 2-aminothiazole (405 mg, 2.5 mmol) following the general procedure D.

LC-MS (m/z): 349 ($M+1$)⁺.

^1H NMR (400 MHz, acetone- d_6): δ 6.73 (d, J = 8 Hz, 1H), 6.98 (m, 1H), 7.06 (d, J = 3.2 Hz, 1H), 7.12 (m, 1H), 7.20-7.28 (m, 2H), 7.25 (d, J = 3.6 Hz, 1H), 7.40 (m, 1H), 8.43 (d, J = 8.2 Hz, 1H), 9.05 (br, 1H), 10.12 (br, 1H).

Example 45

5 N-[2-(2-Fluoro-6-methoxyphenoxy)phenyl]-N'-(thiazol-2-yl)urea




2-(2-Fluoro-6-methoxyphenoxy)-1-nitrobenzene (0.84 g, 64%) was prepared from 2-fluoro-6-methoxyphenol (0.78 g, 5.5 mmol) and 1-fluoro-2-nitrobenzene (0.71 g, 5.0 mmol) following the general procedure A. This was reduced to 2-(2-fluoro-6-methoxyphenoxy)-aniline (0.63 g, 85%) following general procedure C. N-2-(2-Fluoro-6-methoxyphenoxy)-phenyl)-N'-(thiazol-2-yl)urea (114 mg, 63%) was prepared from 2-(2-fluoro-6-methoxyphenoxy)aniline (117 mg, 0.5 mmol) and 2-aminothiazole (60 mg, 0.6 mmol) following the general procedure D.

LC-MS (m/z): 361 ($M+1$) $^+$.

15 ^1H NMR (400 MHz, acetone- d_6): δ 3.82 (s, 3H), 6.59 (m, 1H), 6.89-6.96 (m, 2H), 7.02-7.07 (m, 3H), 7.27-7.33 (m, 1H), 7.34 (d, J = 3.6 Hz, 1H), 8.38 (dd, J = 8.4, 1.6 Hz, 1H), 8.85 (br, 1H), 10.31 (br, 1H).

Example 46

N-[2-(2-Methoxy-6-methoxycarbonylphenoxy)phenyl]-N'-(thiazol-2-yl)urea



20

2-(2-Methoxy-6-methoxycarbonylphenoxy)-1-nitrobenzene (1.24 g, 82%) was prepared from methyl 3-methoxysalicylate (1.0 g, 5.5 mmol) and 1-fluoro-2-nitrobenzene

(0.71 g, 5.0 mmol) following the general procedure A. This compound was reduced to 2-(2-methoxy-6-methoxycarbonylphenoxy)aniline (0.89 g, 80%) following the general procedure C. N-[2-(2-Methoxy-6-methoxycarbonylphenoxy)phenyl]-N'-(thiazol-2-yl)urea (143 mg, 72%)


5 was prepared from 2-(2-methoxy-6-methoxycarbonylphenoxy)aniline (136 mg, 0.5 mmol) and 2-aminothiazole (60 mg, 0.6 mmol) following the general procedure D.

LC-MS (*m/z*): 401 (*M*+1)⁺.

¹H NMR (400 MHz, acetone-*d*₆): δ 3.78 (s, 3H), 3.87 (s, 3H), 6.89-6.97 (m, 1H), 7.04-7.23 (m, 3H), 7.36-7.47 (m, 2H), 7.67-7.76 (m, 2H), 8.17 (dd, *J* = 1.6, 10.8 Hz, 1H), 8.78 (br, 1H), 10.64 (br, 1 H).

10 Example 47

N-[(3-methoxy-2-methoxycarbonylphenoxy)phenyl]-N'-(thiazol-2-yl)urea




2-(3-Methoxy-2-methoxycarbonylphenoxy)-1-nitrobenzene (1.21 g, 80%) was prepared from methyl 6-methoxysalicylate (1.09 g, 5.5 mmol) and 1-fluoro-2-nitrobenzene (0.71 g, 5.0 mmol) following the general procedure A. This compound was reduced to 2-(3-methoxy-2-methoxycarbonylphenoxy)aniline (0.82 g, 75%) following the general procedure C. N-[(3-methoxy-2-methoxycarbonylphenoxy)phenyl]-N'-(thiazol-2-yl)urea (130 mg, 65%) was prepared from 2-(3-methoxy-2-methoxycarbonylphenoxy)aniline (136 mg, 0.5 mmol) and 2-aminothiazole (60 mg, 0.6 mmol) following the general procedure D.

20 LC-MS (*m/z*): 401 (*M*+1)⁺.

¹H NMR (400 MHz, acetone-*d*₆): δ 3.66 (s, 3H), 3.80 (s, 3H), 6.44-6.46 (m, 1H), 6.83-6.87 (m, 1H), 6.96-6.99 (m, 1H), 7.03 (d, *J* = 3.6 Hz, 1H), 7.33 (d, *J* = 4 Hz, 1H), 7.37-7.46 (m, 2H), 7.46-7.48 (m, 1H), 8.32-8.35 (dd, *J* = 1.6, 8.0 Hz, 1H), 8.80 (br, 1H), 10.32 (br, 1H).

Example 48


N-[2-(2,3-Dimethoxyphenoxy)phenyl]-N'-(thiazol-2-yl)urea



- 2-(2,3-Dimethoxyphenoxy)-1-nitrobenzene (0.88 g, 64%) was prepared from 2,3-dimethoxyphenol (0.85 g, 5.5 mmol) and 1-fluoro-2-nitrobenzene (0.71 g, 5.0 mmol) following the general procedure A. This was reduced to 2-(2,3-dimethoxyphenoxy)aniline (0.57 g, 73%) following general procedure B. N-[2-(2,3-Dimethoxyphenoxy)phenyl]-N'-(thiazol-2-yl)urea (131 g, 71%) was prepared from 2-(2,3-dimethoxyphenoxy)aniline (123 mg, 0.5 mmol) and 2-aminothiazole (60 mg, 0.6 mmol) following the general procedure D.
- LC-MS (*m/z*): 373 (*M*+1)⁺.
¹H NMR (400 MHz, acetone-*d*₆): δ 3.66 (s, 3H), 3.88 (s, 3H), 6.66-6.74 (m, 2H), 6.91-6.98 (m, 2H), 7.04-7.16 (m, 3H), 7.32 (d, *J* = 3.6 Hz, 1H), 8.39 (dd, *J* = 8.0, 1.6 Hz, 1H), 8.90 (br, 1H), 10.26 (br, 1H).

Example 49

- N-[2-(2,3,4-Trichlorophenoxy)phenyl]-N'-(thiazol-2-yl)urea




- 2-(2,3,4-Trichlorophenoxy)-1-nitrobenzene (1.04 g, 65%) was prepared from 2,3,4-trichlorophenol (0.98 g, 5.0 mmol) and 1-fluoro-2-nitrobenzene (0.71 g, 5.0 mmol) following the general procedure A. This was reduced to 2-(2,3,4-trichlorophenoxy)aniline (0.42 g, 60%, 2.5 mmol scale) following general procedure B. N-[2-(2,3,4-Trichlorophenoxy)phenyl]-N'-(thiazol-2-yl)urea (131 g, 71%) was prepared from 2-(2,3,4-trichlorophenoxy)aniline (123 mg, 0.5 mmol) and 2-aminothiazole (60 mg, 0.6 mmol) following the general procedure D.

(thiazol-2-yl)urea (343 mg, 55%) was prepared from 2-(2,3,4-trichlorophenoxy)aniline (420 mg, 1.5 mmol) and 2-aminothiazole (150 mg, 1.5 mmol) following the general procedure D. LC-MS (m/z): 415 ($M+1$)⁺.

¹H NMR (400 MHz, acetone- d_6): δ 9.97-7.12 (m, 4H), 7.23-7.28 (m, 2H), 7.58 (d, J = 8.8 Hz, 1H), 8.45 (d, J = 8.0 Hz, 1H), 10.13 (br, 1H).

Example 50

N-[2-(2,4,6-Trifluorophenoxy)phenyl]-N'-(thiazol-2-yl)urea




2-(2,4,6-Trifluorophenoxy)-1-nitrobenzene (1.14 g, 85%) was prepared from 2,4,6-trifluorophenol (0.74 g, 5.0 mmol) and 1-fluoro-2-nitrobenzene (0.71 g, 5.0 mmol) following the general procedure A. This was reduced to 2-(2,4,6-trifluorophenoxy)aniline (0.42 g, 70%, 2.5 mmol scale) following general procedure B. N-[2-(2,4,6-Trifluorophenoxy)phenyl]-N'-(thiazol-2-yl)urea (219 mg, 60%) was prepared from 2-(2,4,6-trifluorophenoxy)aniline (237 mg, 1.0 mmol) and 2-aminothiazole (100 mg, 1.0 mmol) following the general procedure D. LC-MS (m/z): 367 ($M+1$)⁺.

¹H NMR (400 MHz, acetone- d_6): δ 6.78 (d, J = 8.4 Hz, 1H), 7.00-7.21 (m, 5H), 7.36 (d, J = 3.6 Hz, 1H), 8.42 (d, J = 8.4 Hz, 1H), 9.05 (br, 1H), 10.36 (br, 1H).

Example 51

N-[2-(2,4-dichloronaphth-1-oxy)phenyl]-N'-(thiazol-2-yl)urea



2-(2,4-Dichloronaphth-1-oxy)-1-nitrobenzene (1.17 g, 60%) was prepared from 2,4-dichloronaphth-1-ol (1.06 g, 5.0 mmol) and 1-fluoro-2-nitrobenzene (0.71 g, 5.0 mmol) following the general procedure A. This was reduced to 2-(2,4-dichloronaphth-1-oxy)aniline


(0.45 g, 60%, 2.5 mmol scale) following general procedure B. N-[2-(2,4-Dichloronaphth-1-oxy)phenyl]-N'-thiazolylurea (172 g, 40%) was prepared from 2-(2,4-dichloronaphth-1-oxy)aniline (303 mg, 1.0 mmol) and 2-aminothiazole (100 mg, 1.0 mmol) following the general procedure D.

5 LC-MS (m/z): 431 ($M+1$)⁺.

¹H NMR (400 MHz, acetone-*d*₆): δ 6.39 (d, J = 8.2 Hz, 1H), 6.86 (m, 1H), 7.07 (m, 2H), 7.34 (d, J = 3.6 Hz, 1H), 7.71-7.88 (m, 3H), 8.00 (d, J = 8.0 Hz, 1H), 8.32 (d, J = 8.0 Hz, 1H), 8.48 (d, J = 8.0 Hz, 1H), 9.15 (br, 1H), 10.35 (br, 1H).

Example 52

10 N-[2-(2-Methoxyphenoxy)-5-(methylsulfonyl)phenyl]-N'-(thiazol-2-yl)urea




2-(2-Methoxyphenoxy)-5-(methylsulfonyl)-1-nitrobenzene (1.21 g, 75%) was prepared from 2-methoxyphenol (0.68 g, 5.5 mmol) and 1-fluoro-4-methylsulfonyl-2-nitrobenzene (1.09 g, 5.0 mmol) following the general procedure A. This was reduced to 2-(2-methoxyphenoxy)-5-(methylsulfonyl)aniline (0.68 g, 62%) following general procedure B. N-[2-(2-Methoxyphenoxy)-5-(methylsulfonyl)phenyl]-N'-(thiazol-2-yl)urea (136 mg, 65%) was prepared from 2-(2-methoxyphenoxy)-5-(methylsulfonyl)aniline (146 mg, 0.5 mmol) and 2-aminothiazole (60 mg, 0.6 mmol) following the general procedure D.

LC-MS (m/z): 421 ($M+1$)⁺.

20 ¹H NMR (400 MHz, acetone-*d*₆): δ 2.82 (s, 3H), 3.77 (s, 3H), 6.73 (d, J = 8.4 Hz, 1H) 7.05-7.11 (m, 2H), 7.23-7.28 (m, 2H), 7.32-7.37 (m, 2H), 7.50 (dd, J = 8.4, 3.6 Hz, 1H), 9.02 (d, J = 2.4 Hz, 1H), 9.12 (br, 1H), 10.42 (br, 1H).

Example 53

N-[5-Cyano-2-(2-methoxyphenoxy)phenyl]-N'-(thiazol-2-yl)urea




4-(2-Methoxyphenoxy)-3-nitrobenzonitrile (1.10 g, 81%) was prepared from 2-methoxyphenol (0.68 g, 5.5 mmol) and 4-fluoro-3-nitrobenzonitrile (1.33 g, 5.0 mmol) following general procedure A. This was reduced to 4-(2-methoxyphenoxy)-3-aminobenzonitrile (0.62 g, 63%) following general procedure B. N-[5-Cyano-2-(2-methoxyphenoxy)-phenyl]-N'-(thiazol-2-yl)urea (130 mg, 71%) was prepared from 4-(2-methoxyphenoxy)-3-aminobenzonitrile (120 mg, 0.5 mmol) and 2-aminothiazole (60 mg, 0.6 mmol) following general procedure D.

LC-MS (*m/z*): 368 (*M*+1)⁺.

¹H NMR (400 MHz, acetone-*d*₆): δ 3.79 (s, 3H), 6.69 (d, *J* = 8.4 Hz, 1H) 7.05-7.09 (m, 1H), 7.10 (d, *J* = 3.6 Hz, 1H), 7.22-7.27 (m, 2H), 7.32-7.36 (m, 3H), 8.75 (d, *J* = 2.0 Hz, 1H), 9.12 (br, 1H), 10.38 (br, 1H).

Example 54

N-[5-Fluoro-2-(2-methylsulfanylphenoxy)phenyl]-N'-thiazolylurea



5-Fluoro-2-(2-methylsulfanylphenoxy)-1-nitrobenzene (0.95 g, 68%) was prepared from 2-hydroxythioanisole (0.77 g, 5.5 mmol) and 2,5-difluoro-1-nitrobenzene (0.80 g, 5.0 mmol) following the general procedure A. This was reduced to 5-fluoro-2-(2-methylsulfanyl-


phenoxy)aniline (0.52 g, 62%) following general procedure B. N-[5-Fluoro-2-(2-methylsulfanyphenoxy)phenyl]-N'-(thiazol-2-yl)urea (120 mg, 65%) was prepared from 5-fluoro-2-(2-methylsulfanyphenoxy)aniline (125 mg, 0.5 mmol) and 2-aminothiazole (60 mg, 0.6 mmol) following the general procedure D.

5 LC-MS (*m/z*): 377 (*M*+1)⁺.

¹H NMR (400 MHz, acetone-*d*₆): δ 2.48 (s, 3H), 6.61 (m, 2H), 6.92 (d, *J* = 7.6 Hz, 1H), 7.06 (d, *J* = 3.6 Hz, 1H), 7.16-7.25 (m, 2H), 7.31 (d, *J* = 3.6 Hz, 1H), 7.39 (dd, *J* = 7.6, 2.0 Hz, 1H), 8.26-8.29 (m, 1H), 9.00 (br, 1H), 10.32 (br, 1H).

Example 55

10 N-[5-Fluoro-2-(2-fluoro-6-methoxyphenoxy)phenyl]-N'-(thiazol-2-yl)urea



5-Fluoro-2-(2-fluoro-6-methoxyphenoxy)-1-nitrobenzene (0.91 g, 65%) was prepared from 2-fluoro-6-methoxyphenol (0.78 g, 5.5 mmol) and 2,5-difluoro-1-nitrobenzene (0.80 g, 5.0 mmol) following the general procedure A. This was reduced to 5-fluoro-2-(2-


15 fluoro-6-methoxyphenoxy)aniline (0.66 g, 81%) following general procedure C. N-[5-Fluoro-2-(2-fluoro-6-methoxyphenoxy)phenyl]-N'-(thiazol-2-yl)urea (130g, 69%) was prepared from 5-fluoro-2-(2-fluoro-6-methoxyphenoxy)aniline (126 mg, 0.5 mmol) and 2-aminothiazole (60 mg, 0.6 mmol) following the general procedure D.

LC-MS (*m/z*): 379 (*M*+1)⁺.

20 ¹H NMR (400 MHz, acetone-*d*₆): δ 3.83 (s, 3H), 6.60-6.70 (m, 2H), 6.92-6.97 (m, 1H), 7.03-7.05 (m, 1H), 7.09 (d, *J* = 4.0 Hz, 1H), 7.27-7.33 (m, 1H), 7.36 (d, *J* = 3.6 Hz, 1H), 8.23 (dd, *J* = 10.8, 3.2 Hz, 1H), 9.05 (br, 1H), 10.39 (br, 1H).

Example 56

N-[2-(2,3-Dimethoxyphenoxy)-5-fluorophenyl]-N'-(thiazol-2-yl)urea




2-(2,3-Dimethoxyphenoxy)-5-fluoro-1-nitrobenzene (1.0 g, 68%) was prepared from
5 2,3-dimethoxyphenol (0.85 g, 5.5 mmol) and 2,5-difluoro-1-nitrobenzene (0.80 g, 5.0 mmol)
following the general procedure A. This was reduced to 2-(2,3-dimethoxyphenoxy)-5-fluoro-
aniline (0.67 g, 75%) following general procedure C. N-[2-(2,3-Dimethoxyphenoxy)-5-fluoro-
phenyl]-N'-(thiazol-2-yl)urea (126 g, 65%) was prepared from 2-(2,3-dimethoxyphenoxy)-5-
fluoroaniline (132 mg, 0.5 mmol) and 2-aminothiazole (60 mg, 0.6 mmol) following the
10 general procedure D.

LC-MS (*m/z*): 391 (*M*+1)⁺.

¹H NMR (400 MHz, acetone-*d*₆): δ 3.68 (s, 3H), 3.88 (s, 3H), 6.67 (dd, *J* = 9.2, 1.2 Hz, 1H),
6.70-6.79 (m, 2H), 6.92 (dd, *J* = 8.0, 1.6 Hz, 1H), 7.05-7.09 (m, 2H), 7.32 (d, *J* = 3.6 Hz, 1H),
8.25 (dd, *J* = 11.2, 2.8 Hz, 1H), 9.03 (br, 1H), 10.33 (br, 1H).

Example 57

N-[2-(3,4-Difluorophenoxy)-5-fluorophenyl]-N'-(thiazol-2-yl)urea



2-(3,4-Difluorophenoxy)-5-fluoro-1-nitrobenzene (1.05 g, 78%) was prepared from
3,4-difluorophenol (0.72 g, 5.5 mmol) and 2,5-difluoro-1-nitrobenzene (0.8 g, 5.0 mmol)

following the general procedure A. This compound was reduced to 2-(3,4-difluorophenoxy)-5-fluoroaniline (0.63 g, 68%) following the general procedure B. N-[2-(3,4-Difluorophenoxy)-5-fluorophenyl]-N'-(thiazol-2-yl)urea (132 mg, 72%) was prepared from 2-(3,4-difluorophenoxy)-5-fluoroaniline (120 mg, 0.5 mmol) and 2-aminothiazole (60 mg, 0.6 mmol)


5 following the general procedure D.

LC-MS (m/z): 367 ($M+1$)⁺.

¹H NMR (400 MHz, acetone- d_6): δ 6.81-6.89 (m, 2H), 7.05-7.11 (m, 3H), 7.31-7.38 (m, 2H), 8.28 (dd, J = 2.8, 10.8 Hz, 1H), 9.10 (br, 1H), 10.34 (br, 1 H).

Example 58

10 N-[5-Fluoro-2-(4-fluorophenoxy)phenyl]-N'-(thiazol-2-yl)urea




5-Fluoro-2-(4-fluorophenoxy)-1-nitrobenzene (0.94 g, 75%) was prepared from 4-fluorophenol (0.62 g, 5.5 mmol) and 2,5-difluoro-1-nitrobenzene (0.71 g, 5.0 mmol) following the general procedure A. This compound was reduced to 2-(4-fluorophenoxy)-5-fluoroaniline (0.53 g, 64%) following the general procedure B. N-[5-Fluoro-2-(4-fluorophenoxy)phenyl]-N'-(thiazol-2-yl)urea (130 mg, 75%) was prepared from 5-fluoro-2-(4-fluorophenoxy)aniline (110 mg, 0.5 mmol) and 2-aminothiazole (60 mg, 0.6 mmol) following the general procedure D.

LC-MS (m/z): 349 ($M+1$)⁺.

¹H NMR (400 MHz, acetone- d_6): δ 6.78-6.82 (m, 1H), 6.93-6.97 (m, 1H), 7.06-7.10 (m, 3H), 7.14-7.18 (m, 2H), 7.3 (d, J = 3.2 Hz, 1H), 8.27 (dd, J = 2.8, 10.8 Hz, 1H), 8.82 (br, 1H), 10.28 (br, 1H).

Example 59

N-[2-(2,4-Dichlorophenoxy)-5-fluorophenyl]-N'-(thiazol-2-yl)urea




- 2-(2,4-Dichlorophenoxy)-5-fluoro-1-nitrobenzene (1.17 g, 78%) was prepared from
 5 2,4-dichlorophenol (0.89 g, 5.5 mmol) and 2,5-difluoro-1-nitrobenzene (0.8 g, 5.0 mmol)
 following the general procedure A. This compound was reduced to 2-(2,4-dichlorophenoxy)-
 5-fluoroaniline (0.68 g, 64%) following the general procedure B. N-[2-(2,4-Dichlorophenoxy)-
 5-fluorophenyl]-N'-(thiazol-2-yl)urea (150 mg, 76%) was prepared from 2-(2,4-dichloro-
 phenoxy)-5-fluoroaniline (135 mg, 0.5 mmol) and 2-aminothiazole (60 mg, 0.6 mmol)
 10 following the general procedure D.

LC-MS (m/z): 399 ($M+1$)⁺.

¹H NMR (400 MHz, acetone-*d*₆): δ 6.79-6.84 (m, 1 H), 6.93 (m, 1H), 7.03-7.07 (m, 2H), 7.30
 (d, J = 3.2 Hz, 1H), 7.38 (dd, 2.4, 8.8 Hz 1H), 7.64 (d, J = 2.4 Hz, 1H), 8.29 (dd, J = 2.4, 11.2
 Hz 1 H), 9.00 (br, 1H), 10.24 (bs, 1H).

Example 60

N-[5-Fluoro-2-(4-methoxyphenoxy)phenyl]-N'-(thiazol-2-yl)urea



- 5-Fluoro-2-(4-methoxyphenoxy)-1-nitrobenzene (1.05 g, 80%) was prepared from 4-
 methoxyphenol (0.68 g, 5.5 mmol) and 2,5-difluoro-1-nitrobenzene (0.8 g, 5.0 mmol)
 20 following the general procedure A. This compound was reduced to 5-fluoro-2-(4-methoxy-


phenoxy)aniline (0.62 g, 66%) following the general procedure B. N-[5-Fluoro-2-(4-methoxyphenoxy)phenyl]-N'-(thiazol-2-yl)urea (133 mg, 74%) was prepared from 2-(4-methoxyphenoxy)-5-fluoroaniline (117 mg, 0.5 mmol) and 2-aminothiazole (60 mg, 0.6 mmol) following the general procedure D.

5 LC-MS (m/z): 361 ($M+1$)⁺.

¹H NMR (400 MHz, acetone- d_6): δ 3.79 (s, 3H), 6.75-6.77 (m, 1H), 6.82-6.87 (m, 1H), 6.94-7.01 (m, 3H), 7.07 (d, J = 2.1 Hz, 1H), 7.32 (d, J = 3.6 Hz, 1H), 8.26 (dd, J = 3.2, 11.2 Hz, 1H), 9.05 (br, 1H), 10.32 (br, 1H).

Example 61

10 N-[5-Fluoro-2-(2-methoxyphenoxy)phenyl]-N'-(thiazol-2-yl)urea




5-Fluoro-2-(2-methoxyphenoxy)-1-nitrobenzene (1.07 g, 81%) was prepared from 2-methoxyphenol (0.62 g, 5.5 mmol) and 2,5-difluoro-1-nitrobenzene (0.8 g, 5.0 mmol) following the general procedure A. This compound was reduced to 2-(2-methoxyphenoxy)-5-fluoroaniline (0.58 g, 66%) following the general procedure B. 1-[5-Fluoro-2-(2-methoxyphenoxy)phenyl]-3-(thiazol-2-yl)urea (129 mg, 72%) was prepared from 5-fluoro-2-(2-methoxyphenoxy)aniline (117 mg, 0.5 mmol) and 2-aminothiazole (60 mg, 0.6 mmol) following the general procedure D.

LC-MS (m/z): 361 ($M+1$)⁺.

20 ¹H NMR (400 MHz, acetone- d_6): δ 3.79 (s, 3H), 6.65-6.7 (m, 2H), 6.98-7.01 (m, 1H), 7.05-7.22 (m, 4H), 7.32 (d, J = 3.6 Hz, 1H), 8.22 (dd, J = 2.8, 10.8 Hz, 1H), 9.12 (br, 1H), 10.42 (br, 1H).

Example 62


N-[5-Fluoro-2-(2-trifluoromethylphenoxy)phenyl]-N'-(thiazol-2-yl)urea



5-Fluoro-2-(2-trifluoromethylphenoxy)-1-nitrobenzene (1.17 g, 78%) was prepared from 2-hydroxybenzotrifluoride (0.89 g, 5.5 mmol) and 2,5-difluoro-1-nitrobenzene (0.8 g, 5.0 mmol) following the general procedure A. This compound was reduced to 5-fluoro-2-(2-trifluoromethylphenoxy)-aniline (0.65 g, 62%) following the general procedure B. N-[5-Fluoro-2-(2-trifluoromethylphenoxy)phenyl]-N'-(thiazol-2-yl)urea (146 mg, 74%) was prepared from 2-(2-trifluoromethylphenoxy)-5-fluoroaniline (135 mg, 0.5 mmol) and 2-aminothiazole (60 mg, 0.6 mmol) following the general procedure D.

LC-MS (*m/z*): 399 (*M*+1)⁺.¹H NMR (400 MHz, acetone-*d*₆): δ 6.81-6.87 (m, 1H), 6.95-7.05 (m, 3H), 7.21 (m, 1H), 7.31-7.36 (m, 1H), 7.6-7.65 (m, 1H), 7.79 (d, *J* = 7.6 Hz, 1H), 8.30 (dd, *J* = 3.2, 11.2 Hz, 1H), 8.72 (br, 1H), 10.24 (br, 1H).**Example 63**

N-[5-Fluoro-2-(naphth-2-oxy)phenyl]-N'-(thiazol-2-yl)urea



5-Fluoro-2-(naphth-2-oxy)-1-nitrobenzene (1.13 g, 80%) was prepared from 2-naphthol (0.79 g, 5.5 mmol) and 2,5-difluoro-1-nitrobenzene (0.8 g, 5.0 mmol) following the


general procedure A. This compound was reduced to 2-(2-naphth-2-oxy)-5-fluoroaniline (0.64 g, 64%) following the general procedure B. N-[5-Fluoro-2-(naphth-2-oxy)phenyl]-N'-(thiazol-2-yl)urea (123 mg, 65%) was prepared from 5-Fluoro-2-(naphth-2-oxy)-aniline (127 mg, 0.5 mmol) and 2-aminothiazole (60 mg, 0.6 mmol) following the general procedure D.

5 LC-MS (m/z): 381 ($M+1$)⁺.

¹H NMR (400 MHz, acetone-*d*₆): δ 6.83-6.87 (m, 1H), 7.01(bs, 1H), 7.08 (m, 1H), 7.33 (bs, 1H), 7.42-7.49 (m, 3H), 7.80 (d, J = 7.6 Hz, 1H), 7.90(d, J = 7.6 Hz, 1H), 7.97 (d, J = 8.8 Hz, 1H), 8.32 (d, J = 8.4 Hz, 1H), 8.62 (br, 1H), 10.22 (br, 1H)

Example 64

10 N-[2-(2,3-Dimethoxyphenoxy)-6-fluorophenyl]-N'-(thiazol-2-yl)urea




2-(2,3-Dimethoxyphenoxy)-6-fluoro-nitrobenzene (1.05 g, 72%) was prepared from 2,3-dimethoxyphenol (0.85 g, 5.5 mmol) and 2,6-difluoronitrobenzene (0.80 g, 5.0 mmol) following the general procedure A. This compound was reduced to 2-(2,3-dimethoxyphenoxy)-6-fluoroaniline (0.66 g, 70%) following the general procedure C N-[2-(2,3-Dimethoxyphenoxy)-6-fluorophenyl]-N'-(thiazol-2-yl)urea (136 mg, 70%) was prepared from 2-(2,3-dimethoxyphenoxy)-6-fluoroaniline (131 mg, 0.5 mmol) and 2-aminothiazole (60 mg, 0.6 mmol) following the general procedure D.

LC-MS (m/z): 391 ($M+1$)⁺.

20 ¹H NMR (400 MHz, acetone-*d*₆): δ 3.64 (s, 3H), 3.85 (s, 3H), 6.59 (d, J = 8.0 Hz, 1H), 6.68-6.69 (d, J = 8.0 Hz, 1H), 6.88-7.06 (m, 4H), 7.22-7.22 (m 1H), 7.30-7.31 (d, J = 3.2 Hz, 1H), 8.18 (br, 1H), 10.26 (br, 1H).

Example 65


N-[2-(4-Fluoro-2-methoxyphenoxy)phenyl]-N'-(thiazol-2-yl)urea



- 4-Fluoro-2-(2-methoxyphenoxy)-1-nitrobenzene (1.2 g, 91.6%) was prepared from 2-methoxyphenol (0.68 g, 5.5 mmol) and 2,4-difluoro-1-nitrobenzene (0.80 g, 5.0 mmol) following the general procedure F. The product was then reduced to 4-fluoro-2-(2-methoxyphenoxy)aniline (0.96 g, 73%) following general procedure B. N-[4-Fluoro-2-(2-methoxyphenoxy)phenyl]-N'-(thiazol-2-yl)urea (0.32 g, 86%) was prepared from 4-fluoro-2-(2-methoxyphenoxy)aniline (0.117 g, 0.5 mmol) and 2-aminothiazole (0.060 g, 0.6 mmol) following the general procedure D.

LC-MS (m/z): 361 ($M+1$)⁺.¹H NMR (400 MHz, CDCl₃): δ 3.78 (s, 3H), 6.32 (dd, J = 4.0, 14.0 Hz, 1H), 6.35 (dd, J = 2.8, 10.4 Hz, 1H), 6.61 (t, J = 4.0 Hz, 1H), 6.80 (m, 2H), 7.05 (m, 2H), 7.18 (m, 2H), 11.46 (br, 2H).**Example 66**

N-[2-(2,3-Dimethoxyphenoxy)-4-fluorophenyl]-N'-(thiazol-2-yl)urea



- 2-(2,3-Dimethoxyphenoxy)-4-fluoro-1-nitrobenzene (0.88 g, 60%) was prepared from 2,3-dimethoxyphenol (0.85 g, 5.5 mmol) and 2,4-difluoro-1-nitrobenzene (0.8 g, 5.0 mmol) following the general procedure F. This compound was reduced to 2-(2,3-dimethoxyphenoxy)-4-fluoroaniline (0.52 g, 66%) following the general procedure C. N-[2-(2,3-dimethoxyphenoxy)-4-fluorophenyl]-N'-(thiazol-2-yl)urea (116 mg, 60%) was prepared from 2-


(2,3-dimethoxyphenoxy)-4-fluoroaniline (132 mg, 0.5 mmol) and 2-aminothiazole (60 mg, 0.6 mmol) following the general procedure D.

LC-MS (m/z): 391 ($M+1$)⁺.

¹H NMR (400 MHz, acetone- d_6): δ 3.68 (s, 3H), 3.89 (s, 3H), 6.46 (dd, J = 2.4, 9.2 Hz, 1H),
5 6.76 (dd, J = 1.6, 8.0 Hz, 1H); 6.82-6.87 (m, 1H), 6.98 (dd, J = 2.3, 6.8 Hz, 1H), 7.04 (d, J =
3.6 Hz, 1H), 7.1-7.13 (t, J = 8.4 Hz, 1H), 7.31 (d, J = 4.0 Hz, 1H), 8.33-8.37 (dd, J = 6.4, 9.6
Hz, 1H), 8.42 (br, 1H), 10.23 (br, 1H).

Example 67

N-[4-Fluoro-2-(2-fluoro-6-methoxyphenoxy)phenyl]-N'-(thiazol-2-yl)urea



10


4-Fluoro-2-(2-fluoro-6-methoxyphenoxy)-1-nitrobenzene (0.91 g, 65%) was
prepared from 2-fluoro-6-methoxyphenol (0.78 g, 5.5 mmol) and 2,4-difluoro-1-nitrobenzene
(0.8 g, 5.0 mmol) following the general procedure F. This compound was reduced to 4-fluoro-
2-(2-fluoro-6-methoxyphenoxy)aniline (0.61 g, 75%) following the general procedure C. N-[4-
15 Fluoro-2-(2-fluoro-6-methoxyphenoxy)phenyl]-N'-(thiazol-2-yl)urea (122 mg, 65%) was
prepared from 4-fluoro-2-(2-fluoro-6-methoxyphenoxy)aniline (126 mg, 0.5 mmol) and 2-
aminothiazole (60 mg, 0.6 mmol) following the general procedure D.

LC-MS (m/z): 379 ($M+1$)⁺.

¹H NMR (400 MHz, acetone- d_6): δ 3.84 (s, 3H), 6.39 (dd, 2.8, 9.6 Hz 1H), 6.81-6.85 (m, 1H),
20 6.95-6.98 (m, 1H), 7.05-7.07 (m, 2 H) 7.29-7.35 (m, 2H), 8.32-8.36 (m, 1H), 8.85 (br, 1H),
10.26 (br, 1H)

Example 68

N-[2-(2-Fluoro-6-methoxyphenoxy)-4-methoxyphenyl]-N'-(thiazol-2-yl)urea




A mixture of 4-fluoro-2-(2-fluoro-6-methoxyphenoxy)-1-nitrobenzene (1.33 g, 5 mmol) and sodium methoxide (0.35g, 6 mmol) in DMF (10 ml) was heated at 80°C for 2 h. The mixture was poured into water and was extracted with ethyl acetate (3x20 ml). The combined organic layer was washed with water, brine and dried (Na₂SO₄). The solution was concentrated under reduced pressure to obtain 2-(2-fluoro-6-methoxyphenoxy)-4-methoxy-1-nitrobenzene (1.02 g, 70%). This compound was reduced to 2-(2-fluoro-6-methoxyphenoxy)-4-methoxyaniline (0.62 g, 73%) following the general procedure C. N-[2-(2-Fluoro-6-methoxyphenoxy)-4-methoxyphenyl]-N'-(thiazol-2-yl)urea (136 mg, 70%) was prepared from 2-(2-methoxy-6-fluorophenoxy)-4-methoxyaniline (132 mg, 0.5 mmol) and 2-aminothiazole (60 mg, 0.6 mmol) following the general procedure D.

LC-MS (*m/z*): 391 (*M*+1)⁺.

¹H NMR (400 MHz, CDCl₃): δ 3.74 (s, 3H), 6.19 (bs, 1H), 6.46 (dd, *J* = 2.4, 7.6 Hz, 1H), 6.61-6.84 (m, 3H), 7.04-7.19 (m, 1H), 7.32-7.44 (bs, 1H), 8.04-8.12 (m, 1 H), 8.20 (br, 1H), 10.22 (br, 1 H).

Example 69

N-[3-Fluoro-2-(2-fluoro-6-methoxyphenoxy)phenyl]-N'-(thiazol-2-yl)urea




Methyl 3-fluoro-2-(2-fluoro-6-methoxyphenoxy)benzoate (0.88 g, 60%) was prepared from methyl 2,3-difluorobenzoate (0.86 g, 5 mmol) and 2-fluoro-6-methoxyphenol (0.78 g, 5.5 mmol) as described in procedure A. Hydrolysis of this ester with LiOH in aqueous methanol furnished 2-(2-methoxy-6-fluorophenoxy)-3-fluorobenzoic acid (0.86 g, 90%). To a solution of 3-fluoro-2-(2-fluoro-6-methoxyphenoxy)benzoic acid (0.56 g, 2.0 mmol) in 1,2-dichloroethane (10 ml) was added oxalyl chloride (0.18 ml, 2.2 mmol). The solution was stirred at room temperature for 45 min. To this solution was added NaN₃ (390 mg, 6 mmol) and the mixture was heated to reflux for 3 h. 2-Aminothiazole (200 mg, 2 mmol) was then added to this mixture and was further refluxed for 3 h. The mixture was concentrated and the residue was purified by column chromatography (silica, CH₂Cl₂ then 10% ethyl acetate in CH₂Cl₂) to afford 1-[3-fluoro-2-(2-fluoro-6-methoxyphenoxy)phenyl]-3-(thiazol-2-yl)urea in (414 mg, 55%).

LC-MS (*m/z*): 379 (*M*+1)⁺.

¹H NMR (400 MHz, acetone-*d*₆): δ 3.72 (s, 3H), 6.44 (d, *J* = 8.4 Hz, 1H), 6.69-6.84 (m, 4H), 7.01-7.12 (m, 2H), 7.35-7.36 (d, *J* = 4.0 Hz, 1H), 8.28 (br, 1H), 10.28 (br, 1H).

Example 70

N-[2-(2,3-Dimethoxyphenoxy)-5-methoxyphenyl]-N'-thiazol-2-ylurea




2-(2,3-Dimethoxyphenoxy)-5-methoxy-1-nitrobenzene (0.85 g, 56%) was prepared from 2,3-dimethoxyphenol (0.85 g, 5.5 mmol) and 4-chloro-3-nitroanisole (0.94 g, 5.0 mmol) following the general procedure A. This was reduced to 2-(2,3-dimethoxyphenoxy)-5-methoxyaniline (0.65 g, 85%) following general procedure C. N-[2-(2,3-Dimethoxyphenoxy)-5-methoxyphenyl]-N'-(thiazol-2-yl)urea (124 g, 62%) was prepared from 2-(2,3-dimethoxyphenoxy)-5-methoxyaniline (133 mg, 0.5 mmol) and 2-aminothiazole (60 mg, 0.6 mmol) following the general procedure D.

LC-MS (m/z): 403 ($M+1$)⁺.

¹H NMR (400 MHz, acetone-*d*₆): δ 3.70 (s, 3H), 3.79 (s, 3H), 3.87 (s, 3H), 6.53-6.57 (m, 2H), 6.75 (d, J = 8.8 Hz, 1H), 6.85 (dd, J = 8.4, 1.2 Hz, 1H), 7.00 (d, J = 8.4 Hz, 1H), 7.04 (d, J = 3.6 Hz, 1H), 7.30 (d, J = 3.6 Hz, 1H), 8.10 (d, J = 3.6 Hz, 1H), 10.26 (br, 1H).

5 **Example 71**

N-[2-(2,3-Dimethoxyphenoxy)-4-methylphenyl]-N'-(thiazol-2-yl)urea




3-(2,3-Dimethoxyphenoxy)-4-nitrotoluene (0.92 g, 64%) was prepared from 2,3-dimethoxyphenol (0.85 g, 5.5 mmol) and 3-fluoro-4-nitrotoluene (0.78 g, 5.0 mmol) following the general procedure A. This was reduced to 2-(2,3-dimethoxyphenoxy)-4-methylaniline (0.65 g, 79%) following general procedure C. N-[2-(2,3-Dimethoxyphenoxy)-4-methylphenyl]-N'-(thiazol-2-yl)urea (120 mg, 63%) was prepared from 2-(2,3-dimethoxyphenoxy)-4-methylaniline (130 mg, 0.5 mmol) and 2-aminothiazole (60 mg, 0.6 mmol) following the general procedure D.

15 LC-MS (m/z): 387 ($M+1$)⁺.

¹H NMR (400 MHz, acetone-*d*₆): δ 2.20 (s, 3H), 3.67 (s, 3H), 3.88 (s, 3H) 6.57 (d, J = 1.2 Hz, 1H), 6.64 (dd, J = 8.4, 1.6 Hz, 1H), 6.89-6.92 (m, 2H), 7.01-7.08 (m, 2H), 7.31 (d, J = 3.6 Hz, 1H), 8.24 (d, J = 8.4 Hz, 1H), 8.60 (br, 1H), 10.23 (br, 1H).

Example 72

N-[2-(2,3-Dimethoxyphenoxy)-3-methylphenyl]-N'-(thiazol-2-yl)urea




2-(2,3-Dimethoxyphenoxy)-3-nitrotoluene (0.80 g, 56%) was prepared from 2,3-dimethoxyphenol (0.85 g, 5.5 mmol) and 2-chloro-3-nitrotoluene (0.85 g, 5.0 mmol) following the general procedure A. This was reduced to 2-(2,3-dimethoxyphenoxy)-3-methylaniline (0.54 g, 76%) following general procedure C. N-[2-(2,3-Dimethoxyphenoxy)-3-methylphenyl]-N'-(thiazol-2-yl)urea (116 mg, 61%) was prepared from 2-(2,3-dimethoxyphenoxy)-3-methylaniline (130 mg, 0.5 mmol) and 2-aminothiazole (60 mg, 0.6 mmol) following the general procedure D.

LC-MS (m/z): 387 ($M+1$)⁺.

¹H NMR (400 MHz, acetone-*d*₆): δ 2.09 (s, 3H), 3.88 (s, 6H), 6.07 (dd, J = 8.4, 1.2 Hz, 1H), 6.73 (dd, J = 8.4, 1.2 Hz, 1H), 6.85-6.89 (m, 1H), 6.98-7.00 (m, 2H), 7.16-7.20 (m, 1H), 7.24 (d, J = 3.6 Hz, 1H), 8.27 (d, J = 8.0 Hz, 1H), 8.95 (br, 1H), 10.09 (br, 1H).

Example 73

N-[2-(2-Chlorophenoxy)-5-chlorophenyl]-N'-(thiazol-2-yl)urea




N-[2-(2-Chlorophenoxy)-5-chlorophenyl]-N'-(thiazol-2-yl)urea (119 mg, 63%) was prepared from 2-(2-chlorophenoxy)-5-chloroaniline (127 mg, 0.5 mmol) and 2-aminothiazole (60 mg, 0.6 mmol) following the general procedure D.

LC-MS (m/z): 381 ($M+1$)⁺.

¹H NMR (400 MHz, acetone-*d*₆): δ 6.76 (d, *J* = 8.4 Hz, 1H), 7.04 (dd, *J* = 8.4, 2.4 Hz, 1H), 7.07 (d, *J* = 3.6 Hz, 1H), 7.17 (d, *J* = 8.4 Hz, 1H), 7.24-7.42 (m, 3H), 7.59 (dd, *J* = 8.0, 1.6 Hz, 1H), 8.55 (d, *J* = 2.4 Hz, 1H), 9.01 (br, 1H), 10.28 (br, 1H).

Example 74

- 5 N-[5-Chloro-2-(4-chloro-3-methylphenoxy)phenyl]-N'-(thiazol-2-yl)urea




- N-[5-Chloro-2-(4-chloro-3-methylphenoxy)phenyl]-N'-(thiazol-2-yl)urea (236 mg, 60%) was prepared from 2-(2-methyl-3-chlorophenoxy)-5-chloroaniline (commercially available, 268 mg, 1.0 mmol) and 2-aminothiazole (100 mg, 1.0 mmol) following the general procedure D.

LC-MS (*m/z*): 397 (*M*+1)⁺.

¹H NMR (400 MHz, acetone-*d*₆): δ 2.35 (s, 3H), 6.90-7.02 (m, 2H), 7.06-7.10 (m, 3H), 7.32 (d, *J* = 3.6 Hz, 1H), 7.40 (d, *J* = 8.8 Hz, 1H), 8.54 (d, *J* = 2.4 Hz, 1H), 10.22 (br, 2H).

Example 75

- 15 N-[2-(4-Chlorophenoxy)-5-(trifluoromethyl)phenyl]-N'-(thiazol-2-yl)urea




N-[2-(4-Chlorophenoxy)-5-(trifluoromethyl)phenyl]-N'-(thiazol-2-yl)urea (136 mg, 66%) was prepared from 2-(4-Chlorophenoxy)-5-(trifluoromethyl)aniline (144 mg, 0.5 mmol) and 2-aminothiazole (60 mg, 0.6 mmol) following the general procedure D.

LC-MS (m/z): 415 ($M+1$)⁺.

¹H NMR (400 MHz, acetone- d_6): δ 5.12 (br, 1H), 7.02 (d, J = 9 Hz, 1H), 7.09 (d, J = 3.6 Hz, 1H), 7.21 (d, J = 8.8 Hz, 2H), 7.34 (d, J = 3.6 Hz, 1H), 7.39 (d, J = 8.8 Hz, 1H), 7.49 (d, J = 8.8 Hz, 2H), 8.85 (d, J = 1.5 Hz, 1H), 8.95 (br, 1H), 10.29 (br, 1H).

5 Example 76

N-[4,5-difluoro-2-(2,3-dimethoxyphenoxy)phenyl]-N'-(thiazol-2-yl)urea




4,5-Difluoro-2-(2,3-dimethoxyphenoxy)-1-nitrobenzene (1.4 g, 90.3%) was prepared from 2,3-dimethoxyphenol (0.85 g, 5.5 mmol) and 1,3,4-trifluoronitrobenzene (0.885 g, 5.0 mmol) following the general procedure F. The product was then reduced to 2-(2-methoxyphenoxy)-4,5-difluoroaniline (1.35 g, 96.1%) following general procedure B. N-[4,5-Difluoro-2-(2,3-dimethoxyphenoxy)phenyl]-N'-(thiazol-2-yl)urea (0.181 g, 89%) was prepared from 2-(2,3-dimethoxyphenoxy)-4,5-difluoroaniline (0.140 g, 0.5 mmol) and 2-aminothiazole (0.060 g, 0.6 mmol) following the general procedure D.

LC-MS (m/z): 361 ($M+1$)⁺.

¹H NMR (400 MHz, acetone- d_6): δ 3.71 (s, 3H), 3.88 (s, 3H), 6.63 (dd, J = 1.6, 8.4 Hz, 1H), 6.89 (dd, J = 1.2, 6.4 Hz, 1H), 7.03-7.11 (m, 2H), 7.38 (d, J = 3.6 Hz, 1H), 8.26 (dd, J = 1.6 Hz, 7.6 Hz, 1H), 8.42 (dd, J = 1.2 Hz, 7.2 Hz, 1H), 9.13 (br, 1H), 10.22 (br, 1H).

Example 77

N-[4,5-Dichloro-2-(2,3-dimethoxyphenoxy)phenyl]-N'-(thiazol-2-yl)urea




- 4,5-Dichloro-2-(2,3-dimethoxyphenoxy)-1-nitrobenzene (1.12 g, 65%) was prepared from 2,3-dimethoxyphenol (0.85 g, 5.5 mmol) and 4,5-dichloro-2-fluoro-1-nitrobenzene (1.05 g, 5.0 mmol) following the general procedure A. This was reduced to 4,5-dichloro-2-(2,3-dimethoxyphenoxy)aniline (0.68 g, 67%) following general procedure B. N-[4,5-Dichloro-2-(2,3-dimethoxyphenoxy)phenyl]-N'-(thiazol-2-yl)urea (136 mg, 62%) was prepared from 4,5-dichloro-2-(2,3-dimethoxyphenoxy)aniline (157 mg, 0.5 mmol) and 2-aminothiazole (60 mg, 0.6 mmol) following the general procedure D.

LC-MS (m/z): 441 ($M+1$)⁺.

¹H NMR (400 MHz, acetone-*d*₆): δ 3.70 (s, 3H), 3.91 (s, 3H), 6.81-6.83 (m, 2H), 7.01 (dd, J = 8.4, 1.2 Hz, 1H), 7.09 (d, J = 3.6 Hz, 1H), 7.15 (t, J = 8.4 Hz, 1H), 7.34 (d, J = 3.6 Hz, 1H), 8.65 (s, 1H), 8.95 (br, 1H), 10.36 (br, 1H).

Example 78

N-[5-Chloro-2-(2,3-dimethoxyphenoxy)-4-dimethylaminophenyl]-N'-(thiazol-2-yl)urea




4,5-Dichloro-2-(2,3-dimethoxyphenoxy)-1-nitrobenzene (0.35 g, 1.0 mmol) was heated with dimethylamine in tetrahydrofuran (4 ml, 2 M) in a sealed vial at 80°C for 48 h. The reaction mixture was cooled to room temperature and dissolved in ethyl acetate (15 ml). The solution was washed (water, brine) and concentrated to obtain the desired nitrobenzene, which was reduced to 5-chloro-2-(2,3-dimethoxyphenoxy)-4-(dimethylamino)aniline (0.23 g, 73%) following general procedure B. N-[5-Chloro-2-(2,3-dimethoxyphenoxy)-4-dimethylaminophenyl]-N'-(thiazol-2-yl)urea (136 mg, 61%) was prepared from 5-chloro-2-(2,3-dimethoxyphenoxy)-4-(dimethylamino)aniline (161 mg, 0.5 mmol) and 2-aminothiazole (60 mg, 0.6 mmol) following the general procedure D.

LC-MS (*m/z*): 450 (*M*+1)⁺.

¹H NMR (400 MHz, acetone-*d*₆): δ 2.61 (s, 3H), 2.66 (s, 3H), 3.70 (s, 3H), 3.89 (s, 3H), 6.58 (s, 1H), 6.68 (dd, *J* = 8.4, 1.2 Hz, 1H), 6.92 (dd, *J* = 8.4, 1.6 Hz, 1H), 7.04-7.10 (m, 2H), 8.44 (s, 1H), 9.20 (br, 1H), 10.26 (br, 1H).

Example 79

N-[5-Chloro-2-(2,3-dimethoxyphenoxy)-4-(4-morpholino)phenyl]-N'-(thiazol-2-yl)urea




4,5-Dichloro-2-(2,3-dimethoxyphenoxy)-1-nitrobenzene (0.35 g, 1.0 mmol) was heated with morpholine (4 ml) in a sealed vial at 100°C for 48 h. The reaction mixture was cooled to rt and dissolved in ethyl acetate (15 ml). The solution was washed (water, brine) and concentrated to obtain the desired nitrobenzene, which was reduced to 5-chloro-2-(2,3-dimethoxyphenoxy)-4-(4-morpholino)aniline (0.22 g, 61%) following general procedure B. N-[5-Chloro-2-(2,3-dimethoxyphenoxy)-4-(4-morpholino)phenyl]-N'-(thiazol-2-yl)urea (127 mg, 52%) was prepared from 5-chloro-2-(2,3-dimethoxyphenoxy)-4-(4-morpholino)aniline (182 mg, 0.5 mmol) and 2-aminothiazole (60 mg, 0.6 mmol) following the general procedure D.

LC-MS (*m/z*): 492 (*M*+1)⁺.

^1H NMR (400 MHz, acetone- d_6): δ 2.83 (m, 4H), 3.69-3.72 (m, 7H), 3.89 (s, 3H), 6.58 (s, 1H), 6.69 (dd, J = 8.4, 1.6 Hz, 1H), 6.93 (dd, J = 8.4, 1.6 Hz, 1H), 7.05-7.10 (m, 2H); 7.31 (d, J = 3.2 Hz, 1H), 8.48 (s, 1H), 9.02 (br, 1H), 10.25 (s, 1H).

Example 80

5 N-[2-(2,4-Difluorophenoxy)pyridin-3-yl]-N'-(thiazol-2-yl)urea




N-2-(2,4-Difluorophenoxy)pyridin-3-yl-N'-(thiazol-2-yl)urea (112 mg, 65%) was prepared from 3-amino-2-(2,4-difluorophenoxy)pyridine (111 mg, 0.5 mmol) and 2-aminothiazole (60 mg, 0.6 mmol) following the general procedure D.

10 LC-MS (m/z): 350 ($M+1$) $^+$.

^1H NMR (400 MHz, acetone- d_6): δ 7.08-7.16 (m, 3H), 7.20-7.26 (m, 1H), 7.36 (d, J = 3.2 Hz, 1H), 7.40-7.46 (m, 1 H), 7.71 (dd, J = 4.8, 1.6 Hz, 1 H), 8.69 (dd, J = 8.0, 1.6 Hz, 1 H), 9.00 (br, 1H), 10.31 (br, 1H).

Example 81

15 N-[2-(2-Fluorophenoxy)pyridin-3-yl]-N'-(thiazol-2-yl)urea




N-2-(2-Fluorophenoxy)pyridin-3-yl-N'-(thiazol-2-yl)urea (112 mg, 68%) was prepared from 3-amino-2-(2-fluorophenoxy)pyridine (102 mg, 0.5 mmol) and 2-aminothiazole (60 mg, 0.6 mmol) following the general procedure D.

20 LC-MS (m/z): 332 ($M+1$) $^+$.

^1H NMR (400 MHz, acetone- d_6): δ 7.09 (d, J = 3.6 Hz, 1H) 7.13 (dd, J = 8.0, 4.8 Hz, 1H), 7.25-7.39 (m, 5H), 7.71 (dd, J = 4.8, 1.6 Hz, 1H), 8.69 (dd, J = 8.4, 1.6 Hz, 1H), 9.00 (br, 1H), 10.31 (br, 1H).

Example 82

N-[2-(2-Methoxyphenoxy)pyridin-3-yl]-N'-(thiazol-2-yl)urea




2-(2-Methoxy)phenoxy-3-nitropyridine (0.86 g, 70%) was prepared from 2-methoxy-
5 phenol (0.68 g, 5.5 mmol) and 2-bromo-3-nitropyridine (1.05 g, 5.0 mmol) following the
general procedure A. This compound was reduced to 2-(2-methoxyphenoxy)-3-amino-
pyridine (0.49 g, 65%) following general procedure B. N-[2-(2-Methoxyphenoxy)pyridin-3-yl]-
N'-(thiazol-2-yl)urea (126 g, 74%) was prepared from 2-(2-methoxyphenoxy)-3-aminopyridine
(108 mg, 0.5 mmol) and 2-aminothiazole (60 mg, 0.6 mmol) following the general procedure
10 D.

LC-MS (*m/z*): 344 (*M*+1)⁺.

¹H NMR (400 MHz, acetone-*d*₆): δ 3.69 (s, 3H), 6.74 (d, *J* = 3.6 Hz, 1H), 6.99-7.01 (m, 2H),
7.16-7.21 (m, 3H), 7.45 (d, *J* = 2.8 Hz, 1H), 7.71 (d, *J* = 4.0 Hz, 1H), 8.57-8.59 (d, *J* = 8.0 Hz,
1H), 8.89 (br, 1H), 10.44 (br, 1H).

Example 83

N-[2-(2,3-Dimethoxyphenoxy)pyridin-3-yl]-N-(thiazol-2-yl)urea



2-(2,3-Dimethoxy)phenoxy-3-nitropyridine (1.03 g, 75%) was prepared from 2,3-
dimethoxyphenol (0.85 g, 5.5 mmol) and 2-bromo-3-nitropyridine (1.05 g, 5.0 mmol) following
20 the general procedure A. This compound was reduced to 2-(2,3-dimethoxyphenoxy)-3-
aminopyridine (0.57 g, 62%) following the general procedure B. N-[2-(2,3-Dimethoxy-


phenoxy)pyridin-3-yl]-N'-(thiazol-2-yl)urea (131 mg, 70%) was prepared from 2-(2,3-dimethoxyphenoxy)-3-aminopyridine (123 mg, 0.5 mmol) and 2-aminothiazole (60 mg, 0.6 mmol) following the general procedure D.

LC-MS (*m/z*): 374 (*M*+1)⁺.

- 5 ¹H NMR (400 MHz, acetone-*d*₆): δ 3.58 (s, 3H), 3.87 (s, 3H), 6.81 (dd, *J* = 1.6, 8.0 Hz, 1H), 6.95 (dd, *J* = 1.6, 8.4 Hz, 1H), 7.05-7.07 (m, 3H), 7.35 (d, *J* = 3.2 Hz, 1H), 7.67-7.79 (m, 1H), 8.64-8.86 (dd, *J* = 2.4, 8.0 Hz, 1H), 8.89 (br, 1H), 10.26 (br, 1H).

Example 84

N-[4-Chloro-2-(2-chlorobenzoyl)phenyl]-N'-(thiazol-2-yl)urea




N-[4-Chloro-2-(2-chlorobenzoyl)phenyl]-N'-(thiazol-2-yl)urea (0.51 g, 86%) was prepared from 2-amino-2',5-dichlorobenzophenone (0.4 g, 1.50 mmol) and 2-aminothiazole (150 mg, 1.5 mmol) following the general procedure D.

- 15 LC-MS (*m/z*): 394 (*M*+1), ¹H NMR (400 MHz, CDCl₃): δ 6.45 (br, 1H), 6.79 (dd, *J* = 1.2, 7.6 Hz, 2H), 7.11 (d, *J* = 7.2 Hz, 1H), 7.25-7.49 (m, 1H), 6.91-7.33 (m, 3H), 8.65 (d, *J* = 8.8 Hz, 1H), 8.68 (d, *J* = 9.6 Hz, 1H), 8.93 (br, 1H).

Example 85

N-[4-Chloro-2-(2-fluorobenzoyl)phenyl]-N'-(thiazol-2-yl)urea




N-[4-Chloro-2-(2-fluorobenzoyl)phenyl]-N'-(thiazol-2-yl)urea (116 mg, 62%) was prepared from 2-amino-5-chloro-2'-fluorobenzophenone (125 mg, 0.5 mmol) and 2-aminothiazole (60 mg, 0.6 mmol) following the general procedure D.

LC-MS (m/z): 377 ($M+1$)⁺.

Example 86

N-[2-(2,4-Difluorophenylsulfanyl)phenyl]-N'-(thiazol-2-yl)urea




- 5 2-(2,4-Difluorophenylsulfanyl)-1-nitrobenzene (1.04 g, 78%) was prepared from 2,4-difluorothiophenol (0.73 g, 5.5 mmol) and 2-fluoronitrobenzene (0.71 g, 5.0 mmol) following the general procedure A. This compound was reduced to 2-(2,4-difluorophenylsulfanyl)-aniline (0.64 g, 70%) following the general procedure B. 1-[2-(2,4-Difluorophenylsulfanyl)-phenyl]-3-(thiazol-2-yl)urea (130 g, 72%) was prepared from 2-(2,4-difluorophenylsulfanyl)-aniline (118 mg, 0.5 mmol) and 2-aminothiazole (60 mg, 0.6 mmol) following the general procedure D.
- 10

LC-MS (m/z): 365 ($M+1$)⁺.

- ¹H NMR (400 MHz, acetone-*d*₆): δ 6.69-6.79 (m, 3H), 6.85 (d, J = 3.6 Hz, 1H), 7.04-7.09 (m, 1H), 7.36-7.40 (m 2H), 7.51-7.53 (dd, J = 1.6, 8.0 Hz, 1H), 8.42 (d, J = 8.0 Hz, 1 H), 8.88 (br, 1H), 10.2 (br, 1H).
- 15

Example 87

1-[2-(2-Fluorophenylsulfanyl)phenyl]-3-(thiazol-2-yl)urea



- 20 2-(2-Fluorophenylsulfanyl)nitrobenzene (1.03 g, 83%) was prepared from 2-fluorothiophenol (0.70 g, 5.5 mmol) and 2-fluoronitrobenzene (0.71 g, 5.0 mmol) following the general procedure A. This compound was reduced to 2-(2-fluorophenylsulfanyl)aniline (0.65 g, 72%) following general procedure B. N-[2-(2-Fluorophenylsulfanyl)phenyl]-N'-(thiazol-2-yl)-


urea (129 mg, 75%) was prepared from 2-(2-fluorophenylsulfanyl)aniline (107 mg, 0.5 mmol) and 2-aminothiazole (60 mg, 0.6 mmol) following the general procedure D.

LC-MS (m/z): 347 ($M+1$)⁺.

¹H NMR (400 MHz, acetone-*d*₆): δ 6.74-6.82 (m, 1H), 7.02-7.07 (m, 2H), 7.14-7.23 (m, 3H),
5 7.29 (d, J = 3.6 Hz, 1H), 7.51-7.58 (m, 1H), 7.60 (dd, J = 1.2, 7.6 Hz, 1H), 8.43-8.46 (d, J =
8.8 Hz, 1H), 9.08 (br, 1H), 10.39 (br, 1H).

Example 88

N-[2-(2-Chloro-4-fluorophenylsulfanyl)phenyl]-N'-(thiazol-2-yl)urea




10 2-(2-Chloro-4-fluorophenylsulfanyl)nitrobenzene (1.13 g, 80%) was prepared from 2-chloro-4-fluorothiophenol (0.89 g, 5.5 mmol) and 2-fluoronitrobenzene (0.71 g, 5.0 mmol) following the general procedure A. This compound was reduced to 2-(2-chloro-4-fluorophenylsulfanyl)aniline (0.76 g, 75%) following the general procedure B. N-[2-(2-Chloro-4-fluorophenylsulfanyl)phenyl]-N'-(thiazol-2-yl)urea (144 mg, 76%) was prepared from 2-(2-chloro-4-fluorophenylsulfanyl)aniline (126 mg, 0.5 mmol) and 2-aminothiazole (60 mg, 0.6
15 mmol) following the general procedure D.

LC-MS (m/z): 381 ($M+1$)⁺.

¹H NMR (400 MHz, acetone-*d*₆): δ 6.59-6.68 (m, 1H), 6.96-7.04 (m, 2H), 7.16-7.22 (m, 1H),
7.26 (d, J = 4.0 Hz, 1H), 7.32-7.38 (m, 1H), 7.50-7.61 (m, 2H), 8.47 (d, J = 8.0 Hz, 1H), 9.00
20 (br, 1H), 10.23 (br, 1H).

Example 89


N-[2-(2,3-Dichlorophenylsulfanyl)phenyl]-N'-(thiazol-2-yl)urea



2-(2,3-Dichlorophenylsulfanyl)nitrobenzene (1.25 g, 84%) was prepared from 2,3-dichlorothiophenol (0.97 g, 5.5 mmol) and 2-fluoronitrobenzene (0.71 g, 5.0 mmol) following the general procedure A. This compound was reduced to 2-(2,4-dichlorophenylsulfanyl)-aniline (0.81 g, 72%) following the general procedure B. N-[2-(2,3-Dichlorophenylsulfanyl)-phenyl]-N'-(thiazol-2-yl)urea (154 mg, 78%) was prepared from 2-(2,3-chlorophenylsulfanyl)-aniline (135 mg, 0.5 mmol) and 2-aminothiazole (60 mg, 0.6 mmol) following the general procedure D.

LC-MS (*m/z*): 397 (*M*+1)⁺.¹H NMR (400 MHz, CDCl₃): δ 6.34-6.44 (m, 1H), 6.72-6.78 (m, 1H), 6.88-6.96 (m, 1H), 7.11-7.22 (m, 3H), 7.50-7.57 (m, 2H), 8.51-8.53 (d, *J* = 6.0 Hz, 1H), 9.08 (br, 1H), 10.32 (br, 1H).**Example 90**

N-[2-(3,5-Dimethylphenylsulfanyl)phenyl]-N'-(thiazol-2-yl)urea



2-(3,5-Dimethylphenylsulfanyl)nitrobenzene (1.01 g, 78%) was prepared from 3,5-dimethylthiophenol (0.76 g, 5.5 mmol) and 2-fluoronitrobenzene (0.71 g, 5.0 mmol) following the general procedure A. This compound was reduced to 2-(3,5-dimethylphenylsulfanyl)-aniline (0.64 g, 72%) following the general procedure B. N-[2-(3,5-Dimethylphenylsulfanyl)-phenyl]-N'-(thiazol-2-yl)urea (131 mg, 74%) was prepared from 2-(3,5-dimethylphenyl-


sulfanyl)aniline (115 mg, 0.5 mmol) and 2-aminothiazole (60 mg, 0.6 mmol) following the general procedure D.

LC-MS (*m/z*): 357 (*M*+1)⁺.

¹H NMR (400 MHz, CDCl₃): δ 2.09 (s, 6 H), 6.54-6.86 (m, 4H), 7.04-7.10 (m, 1H), 7.27-7.34 (m, 1H), 7.41-7.46 (m, 1H), 7.48-7.56 (bs, 1H), 8.41 (d, *J* = 8.4 Hz, 1 H), 8.20 (br, 1H), 11.8 (br, 1H).

Example 91

N-[2-(2-Methoxycarbonylphenylsulfanyl)phenyl]-N'-(thiazol-2-yl)urea




2-(2-Methoxycarbonylphenylsulfanyl)nitrobenzene (1.15 g, 80%) was prepared from methyl thiosalicylate (0.92 g, 5.5 mmol) and 2-fluoronitrobenzene (0.71 g, 5.0 mmol) following the general procedure A. This compound was reduced to 2-(2-methoxycarbonylphenylsulfanyl)aniline (0.75 g, 73%) following the general procedure B. N-[2-(2-Methoxycarbonylphenylsulfanyl)phenyl]-N'-(thiazol-2-yl)urea (142 mg, 74%) was prepared from 2-(2-methoxycarbonylphenylsulfanyl)aniline (130 mg, 0.5 mmol) and 2-aminothiazole (60 mg, 0.6 mmol) following the general procedure D.

LC-MS (*m/z*): 387 (*M*+1)⁺.

¹H NMR (400 MHz, acetone-*d*₆): δ 3.88 (s, 3H), 6.62-6.82 (m, 2H), 7.02-7.24 (m, 4H), 7.44-7.50 (m, 1H), 7.55-7.64 (bs, 1H), 7.87 (d, *J* = 6.4, 1H), 8.42 (d, *J* = 8.0 Hz, 1H), 8.82 (br, 1H), 10.9 (br, 1H).

Example 92

N-[2-(2-Methoxyphenylsulfanyl)phenyl]-N'-(thiazol-2-yl)urea




2-(2-Methoxyphenylsulfanyl)nitrobenzene (1.07 g, 82%) was prepared from 2-methoxy thiophenol (0.76 g, 5.5 mmol) and 2-fluoronitrobenzene (0.71 g, 5.0 mmol) following the general procedure A. This compound was reduced to 2-(2-methoxyphenylsulfanyl)aniline (0.66 g, 70%) following general procedure B. N-[2-(2-Methoxyphenylsulfanyl)phenyl]-N'-(thiazol-2-yl)urea (110 mg, 62 %) was prepared from 2-(2-methoxyphenylsulfanyl)aniline (115 mg, 0.5 mmol) and 2-aminothiazole (60 mg, 0.6 mmol) following the general procedure D.

LC-MS (*m/z*): 359 (*M*+1)⁺.

¹H NMR (400 MHz, acetone-*d*₆): δ 3.48 (s, 3H), 6.77-6.89 (m, 2H), 6.90-7.00 (m, 2H), 7.05-7.24 (m, 2H), 7.29 (d, *J* = 3.6 Hz, 1H), 7.38-7.42 (m, 1 H), 7.55 (dd, *J* = 1.6, 8.0 Hz, 1H), 8.41-8.43 (d, *J* = 8.4 Hz, 1H), 8.89 (br, 1H), 10.22 (br, 1H).

Example 93

N-[2-(2-Pyridylsulfanyl)phenyl]-N'-(thiazol-2-yl)urea




2-(2-Pyridylsulfanyl)nitrobenzene (0.70 g, 60%) was prepared from 2-mercapto-pyridine (0.61g, 5.5 mmol) and 2-fluoronitrobenzene (0.71 g, 5.0 mmol) following the general procedure A. This compound was reduced to 2-(2-pyridylsulfanyl)aniline (0.37 g, 62%) following general procedure B. N-[2-(2-Pyridylsulfanyl)phenyl]-N'-(thiazol-2-yl)urea (98 mg, 60%) was prepared from 2-(2-pyridylsulfanyl)aniline (101 mg, 0.5 mmol) and 2-aminothiazole (60 mg, 0.6 mmol) following the general procedure D.

LC-MS (*m/z*): 330 (*M*+1)⁺.

^1H NMR (400 MHz, CDCl_3): δ 6.78-6.80 (d, J = 8.0 Hz, 1H), 6.96 (d, J = 3.6 Hz, 1H), 7.03-7.07 (m, 1H), 7.13-7.18 (m, 1H), 7.23 (d, J = 2.8 Hz, 1H), 7.51-7.57 (m, 2H), 7.64 (d, J = 1.6, 7.6 Hz, 1H), 8.35 (d, J = 4.0 Hz, 1H), 8.45 (d, J = 8.4 Hz, 1H), 8.13 (br, 1H), 10.44 (br, 1H).

Example 94

5 N-(2-Propyloxyphenyl)-N'-(thiazol-2-yl)urea




N-(2-Propyloxyphenyl)-N'-(thiazol-2-yl)urea (97 mg, 70%) was prepared from 2-propyloxyaniline (76 mg, 0.5 mmol) and 2-aminothiazole (60 mg, 0.6 mmol) following the general procedure D.

10 LC-MS (m/z): 279 ($M+1$) $^+$.

^1H NMR (400 MHz, acetone- d_6): δ 1.07 (t, J = 7.6 Hz, 3H), 1.88 (m, 2H), 4.05 (q, J = 7.6 Hz), 6.90-7.01 (m, 3H), 7.05 (d, J = 3.6 Hz), 7.36 (d, J = 3.6 Hz), 8.30 (dd, J = 7.6, 1.6 Hz, 1H), 8.80 (br, 1H), 10.24 (br, 1H).

Example 95


15 N-(2-Butyloxyphenyl)-N'-(thiazol-2-yl)urea



N-(2-Butyloxyphenyl)-N'-(thiazol-2-yl)urea (94 mg, 65%) was prepared from 2-butyloxyaniline (82 mg, 0.5 mmol) and 2-aminothiazole (60 mg, 0.6 mmol) following the general procedure D.

20 LC-MS (m/z): 293 ($M+1$) $^+$.

¹H NMR (400 MHz, acetone-*d*₆): δ 0.98 (t, *J* = 7.2 Hz, 3H), 1.54 (m, 2H), 1.84 (quint, *J* = 6.4 Hz, 2H), 4.10 (t, *J* = 6.4 Hz, 2H), 6.90-7.03 (m, 3H), 7.05 (d, *J* = 3.6 Hz, 1H), 7.25 (d, *J* = 3.6 Hz, 1H), 8.30 (dd, *J* = 8.0, 1.2 Hz, 1H), 9.00 (br, 1H), 10.20 (br, 1H).


Example 96**5 N-(2-(Cyclopentyloxyphenyl)-N'-(thiazol-2-yl)urea**

2-(Cyclopentyloxy)-1-nitrobenzene (1.04 g, 80%) was prepared from cyclopentanol (0.46 ml, 5.0 mmol) and 1-fluoro-2-nitrobenzene (0.71 g, 5.0 mmol) following the general procedure G. This was reduced to 2-(cyclopentyloxy)aniline (0.30 g, 68%, 2.5 mmol scale) following general procedure B. N-(2-Cyclopentyloxyphenyl)-N'-(thiazol-2-yl)urea (212 mg, 70%) was prepared from 2-(cyclopentyloxy)aniline (177 mg, 1.0 mmol) and 2-aminothiazole (100 mg, 1.0 mmol) following the general procedure D.

LC-MS (*m/z*): 305 (*M*+1)⁺.

¹H NMR (400 MHz, acetone-*d*₆): δ 1.60-2.05 (m, 8H), 4.94 (m, 1H), 6.90-7.05 (m, 4H), 7.35 (d, *J* = 4.0 Hz, 1H), 8.30 (d, *J* = 8.8 Hz, 1H), 10.20 (br, 2H).

15

Example 97**N-(2-Isopropoxyphenyl)-N'-(thiazol-2-yl)urea**

2-(Isopropoxy)-1-nitrobenzene (317 mg, 70%) was prepared from isopropyl alcohol (0.24 ml, 3.0 mmol) and 1-fluoro-2-nitrobenzene (0.36 g, 2.5 mmol) following the general procedure G. This was reduced to 2-(isopropoxy)aniline (198 mg, 75%) following general procedure B. N-(2-Isopropoxyphenyl)-N'-(thiazol-2-yl)urea (218 mg, 60%) was prepared from 2-(isopropoxy)aniline (198 mg, 1.3 mmol) and 2-aminothiazole (150 mg, 1.5 mmol) following the general procedure D.


20

LC-MS (m/z): 279 ($M+1$)⁺.

¹H NMR (400 MHz, acetone- d_6): δ 1.36 (d, J = 6.4 Hz, 6H), 4.71 (m, 1H), 6.90-7.07 (m, 4H), 7.36 (d, J = 3.6 Hz, 1H), 8.30 (d, J = 8.0 Hz, 1H), 10.22 (br, 2H)

Example 98

5 N-[2-(2-Methylpropoxy)phenyl]-N'-(thiazol-2-yl)urea




2-(2-Methylpropoxy)-1-nitrobenzene (0.73 g, 75%) was prepared from 2-methylpropanol (0.46 ml, 5.0 mmol) and 1-fluoro-2-nitrobenzene (0.71 g, 5.0 mmol) following the general procedure G. This was reduced to 2-(2-methylpropoxy)aniline (0.29 g, 70%, 2.5
10 mmol scale) following general procedure B. N-[2-(2-Methylpropoxy)phenyl]-N'-(thiazol-2-yl)-urea (189 mg, 65%) was prepared from 2-(2-methylpropoxy)aniline (165 mg, 1.0 mmol) and 2-aminothiazole (100 mg, 1.0 mmol) following the general procedure D.

LC-MS (m/z): 293 ($M+1$)⁺.

¹H NMR (400 MHz, acetone- d_6): δ 1.09 (d, 6H), 2.13 (m, 1H), 3.86 (d, J = 6.8 Hz, 1H), 6.90-
15 7.10 (m, 4H), 7.38 (d, J = 3.6 Hz, 1H), 8.29 (d, J = 7.2 Hz, 1H), 10.05 (br, 2H).

Example 99

N-[2-(Cyclopentylmethoxy)phenyl]-N'-(thiazol-2-yl)urea




2-(Cyclopentylmethoxy)-1-nitrobenzene (443 mg, 80%) was prepared from cyclopentanemethanol (0.32 ml, 3.0 mmol) and 1-fluoro-2-nitrobenzene (0.36 g, 2.5 mmol) following the general procedure G. This was reduced to 2-(cyclopentylmethoxy)aniline (268 mg, 70%) following general procedure B. N-[2-(Cyclopentylmethoxy)phenyl]-N'-(thiazol-2-yl)-
20 urea (286 mg, 60%) was prepared from 2-(cyclopentylmethoxy)aniline (0.25 g, 1.5 mmol) and 2-aminothiazole (150 mg, 1.5 mmol) following the general procedure D.

25 LC-MS (m/z): 319 ($M+1$)⁺.

^1H NMR (400 MHz, acetone- d_6): δ 1.40 (m, 2H), 1.65 (m, 4H), 1.91 (m, 2H), 2.45 (m, 1H), 3.98 (d, J = 7.6 Hz, 1H), 6.88-7.06 (m, 4H), 7.35 (d, J = 3.6 Hz, 1H), 8.29 (d, J = 7.6 Hz, 1H), 10.14, (br, 2H).

Example 100

5 N-[2-(3-pentoxo)phenyl]-N'-(thiazol-2-yl)urea




2-(3-Pentoxo)-1-nitrobenzene (366 mg, 70%) was prepared from 3-pentanol (0.27 ml, 2.5 mmol) and 1-fluoro-2-nitrobenzene (0.36g, 2.5 mmol) following the general procedure G. This was reduced to 2-(3-pentoxo)aniline (0.25 g, 80%) following general procedure B. N-[2-(3-pentoxo)phenyl]-N'-(thiazol-2-yl)urea (0.26 g, 57%) was prepared from 2-(3-pentoxo)-aniline (0.25 g, 1.5 mmol) and 2-aminothiazole (150 mg, 1.5 mmol) following the general procedure D.

LC-MS (m/z): 307 ($M+1$) $^+$.

^1H NMR (400 MHz, acetone- d_6): δ 0.92 (t, J = 7.2 Hz, 6H), 1.75 (m, 4H), 4.38 (m, 1H), 6.89-7.06 (m, 4H), 7.36 (d, J = 3.6 Hz, 1H), 8.32 (d, J = 8.0 Hz, 1H).

Example 101

N-[2-(2-pentoxo)phenyl]-N'-(thiazol-2-yl)urea



2-(2-Pentoxo)-1-nitrobenzene (387 mg, 74%) was prepared from 2-pentanol (0.27 ml, 2.5 mmol) and 1-fluoro-2-nitrobenzene (0.36 g, 2.5 mmol) following the general procedure G. This was reduced to 2-(2-pentoxo)aniline (0.26 g, 79%) following general procedure B. N-[2-(2-pentoxo)phenyl]-N'-(thiazol-2-yl)urea (280 mg, 62%) was prepared from


2-(2-pentoxy)aniline (0.25 g, 1.5 mmol) and 2-aminothiazole (150 mg, 1.5 mmol) following the general procedure D.

LC-MS (m/z): 307 ($M+1$)⁺.

¹H NMR (400 MHz, acetone-*d*₆): δ 0.93 (t, J = 7.2 Hz, 3H), 1.32 (d, J = 6.0 Hz, 3H), 1.40-1.90 (m, 4H), 4.55 (m, 1H), 6.85-7.07 (m, 4H), 7.36 (d, J = 3.6 Hz, 1H), 8.31 (d, J = 8.0 Hz, 1H), 10.15 (br, 1H).

Example 102

N-[2-(2-Methoxyethoxy)phenyl]-N'-(thiazol-2-yl)urea




2-(2-Methoxyethoxy)-1-nitrobenzene (0.25 g, 64%) was prepared from 2-methoxyethanol (0.16 ml, 2.0 mmol) and 1-fluoro-2-nitrobenzene (0.21 ml, 2.0 mmol) following the general procedure G. This was reduced to 2-(2-methoxyethoxy)aniline (0.13 g, 60%) following general procedure B. N-[2-(2-Methoxyethoxy)phenyl]-N'-(thiazol-2-yl)urea (115 mg, 55%) was prepared from 2-(2-methoxyethoxy)aniline (115 mg, 0.7 mmol) and 2-aminothiazole (140 mg, 1.4 mmol) following the general procedure D.

LC-MS (m/z): 295 ($M+1$)⁺.

¹H NMR (400 MHz, acetone-*d*₆): δ 3.37 (s, 3H), 3.78 (t, J = 4.8 Hz, 2H), 4.25 (t, J = 4.8 Hz, 2H), 6.95-7.06 (m, 4H), 7.37 (d, J = 3.6 Hz, 1H), 8.29 (d, J = 8.0 Hz, 1H), 8.80 (br, 1H), 10.20 (br, 1H).

20 Example 103 (General procedure (H))

(2-[3-(2-Benzylphenyl)ureido]thiazol-4-yl)acetic acid



The intermediate 2-benzylphenyl isocyanate (0.26 g, 1.2 mmol) (general procedure H2) was dissolved in DMF (5 ml) and 2-amino-4-thiazole acetic acid (0.20 g, 1.2 mmol) was

added. After 16 hours at 20°C ethylacetate (60 ml) was added and the mixture was extracted with water (5x20 ml). The solvent was removed *in vacuo* and the remaining oil was purified on Waters Deltaprep 4000 giving 150 mg of the title compound.

Preparative system: (gradient 20-90% CH₃CN 40min., 20 ml/min., Rt=35 min

5 Solvent A=water, solvent B=CH₃CN, solvent C=0,5% TFA/water).


¹H-NMR (DMSO-*d*₆): δ 10.90 (broad, 1H); 8.45 (s, 1H); 7.80 (s, 1H); 7.20 (multi, 9H); 6.85 (s, 1H); 3.98 (s, 2H); 3.53 (s, 2H).

HPLC-MS (Method A): m/z = 368 (M+1); R_t = 4.10 min

10 ¹H NMR (CDCl₃): δ 7.98 (br d, 2H), 7.88-7.70 (m, 4H), 7.50 (t, 3H), 7.18 (d, 1H), 6.84 (br s, 1H), 3.71 (br s, 2H), 2.70 (br s, 2H), 2.52 (m, 1H), 1.90-1.70 (m, 5H), 1.45-1.15 (m, 5H).

Example 104 (General procedure (H))

(2-[3-(2-Benzoyl-4-chlorophenyl)ureido]thiazol-4-yl)acetic acid




15 The title compound was prepared as in example 103 using (5-chloro-2-isocyanatophenyl)phenyl methanone (general procedure H2) as intermediate isocyanate.

¹H-NMR (DMSO-*d*₆): Selected data : δ 9.48 (broad, 1H); 8.12 (broad, 1H); (multi, 9H); 6.87 (s, 1H); 3.57 (s, 2H).

HPLC-MS (method A): m/z = 415 (M+1); R_t = 3.79 min

Example 105 (General procedure (H))

20 (2-[3-(2-(2-Methylphenoxy)phenyl)ureido]thiazol-4-yl)acetic acid



The title compound was prepared as in example 103 using 2-(2-methylphenoxy)-phenyl isocyanat (general procedure H2) as intermediate isocyanate.


¹H-NMR (DMSO-*d*₆): δ 11.09 (broad, 1H); 8.88 (broad, 1H); 8.25 (d, 1H); 7.37 (d, 1H); 7.22 (t, 1H); 7.11 (multi, 2H); 6.97 (t, 1H); 6.89 (d, 1H); 6.87 (s, 1H); 6.67 (d, 1H); 3.53 (s, 2H);

5 2.23 (s, 3H).

HPLC-MS (method A): *m/z* = 384 (M+1); *R*_t = 4.67 min

Example 106 (General procedure (H))

(2-[3-(2-(4-Methoxyphenoxy)-5-(trifluoromethyl)phenyl)ureido]thiazol-4-yl)acetic acid




10 The title compound was prepared as in example 103 using 2-(4-methoxyphenoxy)-5-(trifluoromethyl)phenyl isocyanate (general procedure H2) as intermediate isocyanate.

¹H-NMR (DMSO-*d*₆): δ 12.32 (broad, 1H); 11.20 (broad, 1H); 9.30 (d, 1H); 8.62 (d, 1H); 7.32 (d, 1H); 7.17 (d, 2H); 7.05 (d, 2H); 6.90 (s, 1H); 6.83 (d, 1H); 3.78 (s, 3H); 3.53 (s, 2H).

Example 107 (General procedure (H))

15 {2-[3-(2-phenoxyphenyl)ureido]thiazol-4-y}acetic acid




The title compound was prepared as in example 103 using 2-phenoxyphenyl isocyanate (general procedure H2) as intermediate isocyanate.

¹H NMR (CDCl₃) selected data: δ 3.47 (2H, s), 6.78 (1H, s), 6.90 (1H, d), 6.97-7.08 (3H, m), 7.09-7.19 (2H, m), 7.38 (2H, t), 8.16 (1H, br s), HPLC-MS (Method A): *m/z* = 370 (M+1); *R*_t = 3.50 min.

20

Example 108 (General procedure (H))

2-{3-[5-Fluoro-2-(2-fluoro-6-methoxyphenoxy)phenyl]ureido}thiazole-4-carboxylic acid




The title compound was prepared according to general procedure H.

- 5 ^1H NMR (DMSO): δ 12.76 (br, 1H), 11.21 (br 1H), 9.23 (s, 1H), 8.06 (dd, 1H), 7.94 (s, 1H), 7.33 (d, 1H), 7.10-7.01 (m, 2H), 6.74 (t, 1H), 6.55 (dd, 1H), 3.80 (s, 3H).
HPLC-MS (Method B): m/z = 422 ($M+1$); R_t = 3.90min.

Example 109 (General procedure (H))

2-{3-[2-(2,3-Dimethoxyphenoxy)-5-fluorophenyl]ureido}thiazole-4-carboxylic acid ethyl ester




10

- ^1H NMR (DMSO): δ 11.44 (s, 1H), 8.99 (s 1H), 8.06 (dd, 1H), 7.99 (s, 1H), 7.09 (t, 1H), 6.94 (d, 1H), 6.80 (t, 1H), 6.72 (dd, 1H), 6.67 (d, 1H), 4.25 (q, 2H), 3.84 (s, 3H), 3.67 (s, 3H), 1.28 (t, 3H).
HPLC-MS (Method B): m/z = 462 ($M+1$); R_t = 4.53 min.

Example 110 (General procedure (H))

(2-{3-[2-(2,3-Dimethoxyphenoxy)-5-fluorophenyl]ureido}thiazol-4-yl)acetic acid ethyl ester




¹H NMR (DMSO): δ 11.16 (br, 1H), 9.09 (br 1H), 8.08 (dd, 1H), 7.09 (t, 1H), 6.94 (d, 1H), 6.92 (s, 1H), 6.78 (t, 1H), 6.70 (dd, 1H), 6.65 (d, 1H), 4.06 (q, 2H), 3.84 (s, 3H), 3.66 (s, 2H), 3.62 (s, 3H), 1.18 (t, 3H).

HPLC-MS (Method B): m/z = 476 (M+1); R_t = 4.43 min.

Example 111 (General procedure (H))

(2-{3-[5-Fluoro-2-(2-fluoro-6-methoxyphenoxy)phenyl]ureido}thiazol-4-yl)acetic acid




10

¹H NMR (DMSO): δ 12.27 (br, 1H), 11.06 (br, 1H), 9.22 (br, 1H), 8.06 (dd, 1H), 7.32 (dd, 1H), 7.08 (d, 1H), 7.03 (t, 1H), 6.90 (s, 1H), 6.75-6.70 (m, 2H), 6.53 (dd, 1H), 3.79 (s, 3H), 3.55 (s, 2H).

HPLC-MS (Method B): m/z = 436 (M+1); R_t = 3.85 min.

Example 112 (General procedure (H))


2-{3-[2-(2,3-Dimethoxyphenoxy)-5-fluorophenyl]ureido}thiazole-4-carboxylic acid



¹H NMR (DMSO): δ 12.77 (br, 1H), 11.33 (s, 1H), 9.06 (s, 1H), 8.07 (dd, 1H), 7.93 (s, 1H),
5 7.08 (d, 1H), 7.09 (t, 1H), 6.95 (dd, 1H), 6.80 (t, 1H), 6.73-6.63 (m, 2H), 3.84 (s, 3H), 3.66 (s, 3H).

HPLC-MS (Method B): m/z = 434 (M+1); R_t = 3.92 min.**Example 113 (General procedure (H))**

(2-{3-[2-(2,3-Dimethoxyphenoxy)-5-fluorophenyl]ureido}thiazol-4-yl)acetic acid




10

¹H NMR (DMSO): δ 12.30 (br, 1H), 11.14 (br, 1H), 9.08 (br 1H), 8.08 (dd, 1H), 7.08 (t, 1H),
6.94 (d, 1H), 6.89 (s, 1H), 6.78 (t, 1H), 6.70 (dd, 1H), 6.66 (dd, 1H), 3.84 (s, 3H), 3.66 (s, 3H),
3.54 (s, 2H).

HPLC-MS (Method B): m/z = 448 (M+1); R_t = 3.76 min.

Example 114 (General procedure (H))

2-{3-[5-Fluoro-2-(2-fluoro-6-methoxyphenoxy)phenyl]ureido}-4-methylthiazole-5-carboxylic acid ethyl ester




- 5 ^1H NMR (DMSO): δ 11.35 (s, 1H), 9.38 (br 1H), 8.05 (dd, 1H), 7.32 (dd, 1H), 7.10-7.02 (m, 2H), 6.76 (t, 1H), 6.56 (dd, 1H), 4.24 (q, 2H), 3.79 (s, 3H), 2.52 (s, 3H), 1.29 (t, 3H).
HPLC-MS (Method B): m/z = 464 (M+1); R_t = 4.91 min.

Example 115 (General procedure (H))

2-{3-[5-Fluoro-2-(2-fluoro-6-methoxyphenoxy)phenyl]ureido}-4-methylthiazole-5-carboxylic acid


10 acid



- ^1H NMR (DMSO): δ 12.79 (br, 1H), 11.29 (br, 1H), 9.38 (br 1H), 8.05 (dd, 1H), 7.32 (dd, 1H), 7.10-7.01 (m, 2H), 6.75 (t, 1H), 6.54 (dd, 1H), 3.79 (s, 3H), 2.51 (s, 2H).
HPLC-MS (Method B): m/z = 436 (M+1); R_t = 4.19 min.

Example 116 (General procedure (H1))

N-Ethyl-2-(2-{3-[5-fluoro-2-(2-fluoro-6-methoxyphenoxy)phenyl]ureido}thiazol-4-yl)acetamide


¹H NMR (DMSO): δ 11.03 (s, 1H), 9.22 (br 1H), 8.06 (dd, 1H), 7.89 (t, 1H), 7.32 (dd, 1H),

5 7.10-7.00 (m, 2H), 6.82 (s, 1H), 6.72 (t, 1H), 6.53 (dd, 1H), 3.79 (s, 3H), 3.39 (s, 2H), 3.07 (q, 2H), 1.01 (t, 3H).

HPLC-MS (Method B): m/z = 463 (M+1); R_t = 4.06 min.**Example 117** (General procedure (H1))

2-(2-{3-[5-Fluoro-2-(2-fluoro-6-methoxyphenoxy)phenyl]ureido}thiazol-4-yl)-N-(2-methoxy-ethyl)acetamide

10


¹H NMR (DMSO): δ 11.03 (br, 1H), 9.22 (br 1H), 8.06 (dd, 1H), 7.99 (t, 1H), 7.32 (dd, 1H),

7.10-7.00 (m, 2H), 6.82 (s, 1H), 6.72 (t, 1H), 6.53 (dd, 1H), 3.79 (s, 3H), 3.43 (s, 2H), 3.34 (t, 2H), 3.24 (s, 3H), 3.23 (t, 2H).

15 HPLC-MS (Method B): m/z = 493 (M+1); R_t = 4.01 min.

Example 118 (General procedure (H1))

2-(2-{3-[5-Fluoro-2-(2-fluoro-6-methoxyphenoxy)phenyl]ureido}thiazol-4-yl)-N-(2-morpholin-4-ylethyl)acetamide




- 5 ¹H NMR (DMSO): selected data: δ 11.01 (br, 1H), 9.77 (br, 1H), 9.23 (br 1H), 8.19 (br, 1H), 8.06 (dd, 1H), 7.32 (dd, 1H), 7.10-7.00 (m, 2H), 6.88 (s, 1H), 6.72 (t, 1H), 6.53 (dd, 1H), 3.97 (m, 2H), 3.79 (s, 3H), 3.64 (m, 2H), 3.47 (s, 2H).

HPLC-MS (Method B): m/z = 548 (M+1); R_t = 3.24 min.

Example 119 (General procedure (H1))

- 10 2-(2-{3-[5-Fluoro-2-(2-fluoro-6-methoxyphenoxy)phenyl]ureido}thiazol-4-yl)acetyl amino]acetic acid methyl ester




- 15 ¹H NMR (DMSO): Selected data δ 11.06 (br, 1H), 9.25 (br 1H), 8.34 (t, 1H), 8.06 (dd, 1H), 7.31 (dd, 1H), 7.10-7.00 (m, 2H), 6.88 (s, 1H), 6.73 (t, 1H), 6.53 (dd, 1H), 3.85 (d, 2H), 3.79 (s, 3H), 3.63 (s, 2H), 3.49 (s, 2H).

HPLC-MS (Method B): m/z = 507 (M+1); R_t = 3.74 min.

Example 120 (General procedure (H1))

2-{3-[2-(2,3-Dimethoxyphenoxy)-5-fluorophenyl]ureido}thiazole-4-carboxylic acid (2-methoxyethyl)amide




- 5 ^1H NMR (DMSO): Selected data δ 11.29 (s, 1H), 9.13 (s, 1H), 8.08 (dd, 1H), 7.78 (t, 1H), 7.73 (s, 1H), 7.09 (t, 1H), 6.95 (dd, 1H), 6.80 (t, 1H), 6.71 (dd, 1H), 6.67 (dd, 1H), 3.84 (s, 3H), 3.67 (s, 3H), 3.26 (s, 3H).

HPLC-MS (Method B): m/z = 491 ($M+1$); R_t = 3.86 min.

Example 121 (General procedure (H1))

- 10 [2-(2-{3-[5-Fluoro-2-(2-fluoro-6-methoxyphenoxy)phenyl]ureido}thiazol-4-yl)acetamino]acetic acid




- 15 ^1H NMR (DMSO): δ 12.54 (br, 1H), 11.06 (br, 1H), 9.23 (br, 1H), 8.21 (t, 1H), 8.06 (dd, 1H), 7.31 (dd, 1H), 7.09-7.01 (m, 2H), 6.88 (s, 1H), 6.72 (t, 1H), 6.53 (dd, 1H), 3.79 (s, 3H), 3.76 (s, 2H), 3.49 (s, 2H).

HPLC-MS (Method B): m/z = 493 ($M+1$); R_t = 3.47 min.

Example 122 (General procedure (H))

2-{3-[2-(2,3-Dimethoxyphenoxy)-5-fluorophenyl]ureido}thiazole-4-carboxylic acid ethylamide



¹H NMR (DMSO): δ 11.21 (s, 1H), 9.19 (s, 1H), 8.08 (dd, 1H), 7.89 (t, 1H), 7.69 (s, 1H), 7.10 (t, 1H), 6.95 (dd, 1H), 6.79 (t, 1H), 6.73-6.66 (m, 2H), 3.84 (s, 3H), 3.66 (s, 3H), 3.25 (q, 2H), 1.08 (t, 3H).

HPLC-MS (Method B): m/z = 461 (M+1); R_t = 3.96 min.


Microanalysis:

Calculated for : C, 55.09%; H, 4.88%; N, 11.88%.

10 Found: C, 55.18%; H, 4.67%; N, 12.21%.

Example 123 (General procedure (H1))


[(2-{3-[2-(2,3-Dimethoxyphenoxy)-5-fluorophenyl]ureido}thiazole-4-carbonyl)amino]acetic acid methyl ester



15 ¹H NMR (DMSO): δ 11.26 (s, 1H), 9.20 (s, 1H), 8.26 (t, 1H), 8.09 (dd, 1H), 7.77 (s, 1H), 7.10 (t, 1H), 6.95 (dd, 1H), 6.79 (t, 1H), 6.72-6.67 (m, 2H), 4.01 (d, 2H), 3.84 (s, 3H), 3.66 (s, 3H), 3.65 (s, 3H).

Example 124 (General procedure (H))

(5-{3-[5-Fluoro-2-(2-fluoro-6-methoxyphenoxy)phenyl]ureido}-[1,3,4]thiadiazol-2-yl)acetic acid ethyl ester




- 5 ¹H NMR (DMSO): δ 11.35 (s, 1H), 9.31 (s, 1H), 8.03 (t, 1H), 8.04 (dd, 1H), 7.31 (dd, 1H), 7.10-7.02 (m, 2H), 6.75 (t, 1H), 6.55 (dd, 1H), 4.19-4.14 (m, 4H), 3.80 (s, 3H), 1.23 (t, 3H).
HPLC-MS (Method B): m/z = 465 (M+1); R_t = 4.14 min.

Example 125 (General procedure (H1))

[(2-{3-[2-(2,3-Dimethoxyphenoxy)-5-fluorophenyl]ureido}thiazole-4-carbonyl)amino]acetic acid


10 acid



- ¹H NMR (DMSO): δ 12.67 (br, 1H), 11.36 (s, 1H), 9.24 (s, 1H), 8.08 (dd, 1H), 7.78 (s, 1H), 7.10 (t, 1H), 6.95 (dd, 1H), 6.79 (t, 1H), 6.72-6.67 (m, 2H), 3.93 (d, 2H), 3.84 (s, 3H), 3.67 (s, 3H).
HPLC-MS (Method B): m/z = 491 (M+1); R_t = 3.58 min.

Example 126 (General procedure (H))

(5-{3-[5-Fluoro-2-(2-fluoro-6-methoxyphenoxy)phenyl]ureido}-[1,3,4]thiadiazol-2-yl)acetic acid




¹H NMR (DMSO): δ 12.96 (br, 1H), 11.33 (br, 1H), 9.31 (s, 1H), 8.04 (dd, 1H), 7.33 (dd, 1H),
 5 7.11-7.01 (m, 2H), 6.75 (t, 1H), 6.55 (dd, 1H), 4.10 (s, 2H), 3.80 (s, 3H).

HPLC-MS (Method B): m/z = 337 (M+1); R_t = 3.47 min.

Example 127 (General procedure (H))

5-{3-[2-(2,3-Dimethoxyphenoxy)-5-fluorophenyl]ureido}-[1,3,4]thiadiazole-2-carboxylic acid ethyl ester




10

¹H NMR (DMSO): δ 11.91 (s, 1H), 9.26 (s, 1H), 8.05 (dd, 1H), 7.10 (t, 1H), 6.95 (dd, 1H), 6.83
 (t, 1H), 6.75-6.67 (m, 2H), 4.40 (q, 2H), 3.84 (s, 3H), 3.66 (s, 3H), 1.35 (t, 3H).

HPLC-MS (Method B): m/z = 463 (M+1); R_t = 4.34 min.

Example 128 (General procedure (H))

(5-{3-[2-(2,3-Dimethoxyphenoxy)-5-fluorophenyl]ureido}-[1,3,4]thiadiazol-2-yl) acetic acid ethyl ester




- 5 ¹H NMR (DMSO): δ 11.42 (s, 1H), 9.17 (s 1H), 8.05 (dd, 1H), 7.09 (t,1H), 6.95 (dd,1H), 6.81 (t, 1H), 6.72 (dd, 1H), 6.67 (dd, 1H), 4.19-4.13 (m, 4H), 3.84 (s, 3H), 3.66 (s, 3H), 1.22 (t, 3H).

HPLC-MS (Method B): m/z = 477 (M+1); R_t = 4.10 min.

Example 129 (General procedure (H1))

- 10 3-[2-(2-{3-[2-(2,3-Dimethoxyphenoxy)-5-fluorophenyl]ureido}thiazol-4-yl)acetamino]propionic acid methyl ester




- 15 ¹H NMR (DMSO): δ 11.11 (s, 1H), 9.07 (br 1H), 8.08 (dd, 1H), 7.98 (t,1H), 7.08 (d,1H), 6.94 (dd,1H), 6.81 (s, 1H), 6.77 (dd, 1H), 6.70 (dd,1H), 6.65 (dd, 1H), 3.84 (s, 3H), 3.66 (s, 3H), 3.58 (s, 3H), 3.39 (s, 2H), 3.27 (q, 2H), 2.46 (t, 2H).

HPLC-MS (Method B): m/z = 533 (M+1); R_t = 3.71 min.

Example 130 (General procedure (H1))


2-(2-{3-[2-(2,3-Dimethoxyphenoxy)-5-fluorophenyl]ureido}thiazol-4-yl)-N-(2-morpholin-4-ylethyl)acetamide



- 5 ¹H NMR (DMSO): Selected data δ 11.11 (br, 1H), 9.08 (br 1H), 8.07 (dd, 1H), 7.09 (t, 1H), 6.94 (dd, 1H), 6.86 (s, 1H), 6.77 (t, 1H), 6.70 (dd, 1H), 6.66 (dd, 1H), 3.84 (s, 3H), 3.66 (s, 3H), 3.43 (s, 2H), 3.03-2.99 (m, 4H), 1.74-1.71 (m, 4H).
HPLC-MS (Method B): m/z = 560 (M+1); R_t = 2.75 min.

Example 131 (General procedure (H1))


- 10 [(2-{3-[5-Fluoro-2-(2-fluoro-6-methoxyphenoxy)phenyl]ureido}thiazole-4-carbonyl)amino]acetic acid methyl ester



- 15 ¹H NMR (DMSO): δ 11.27 (s, 1H), 9.36 (s 1H), 8.32 (t, 1H), 8.07 (dd, 1H), 7.79 (s, 1H), 7.33 (dd, 1H), 7.11-7.02 (m, 2H), 6.75 (t, 1H), 6.55 (dd, 1H), 4.03 (d, 2H), 3.80 (s, 3H), 3.66 (s, 3H).
HPLC-MS (Method B): m/z = 493 (M+1); R_t = 3.91 min.

Example 132 (General procedure (H1))


3-[2-(2-{3-[2-(2,3-Dimethoxyphenoxy)-5-fluorophenyl]ureido}thiazol-4-yl)acetyl]amino]propionic acid



- 5 ^1H NMR (DMSO): δ 12.21 (br, 1H), 11.09 (br, 1H), 9.06 (br 1H), 8.07 (dd, 1H), 7.98 (t, 1H), 7.09 (t, 1H), 6.94 (dd, 1H), 6.81 (s, 1H), 6.76 (t, 1H), 6.71 (t, 1H), 6.66 (dd, 1H), 3.84 (s, 3H), 3.66 (s, 3H), 3.39 (s, 2H), 3.24 (q, 2H), 2.38 (t, 2H).

Example 133 (General procedure (H1))


- 10 [(2-{3-[5-Fluoro-2-(2-fluoro-6-methoxyphenoxy)phenyl]ureido}thiazole-4-carbonyl)amino]acetic acid



- ^1H NMR (DMSO): δ 12.74 (br, 1H), 11.23 (s, 1H), 9.31 (s 1H), 8.17 (t, 1H), 8.07 (dd, 1H), 7.78 (s, 1H), 7.33 (dd, 1H), 7.11-7.01 (m, 2H), 6.75 (t, 1H), 6.54 (dd, 1H), 3.92 (d, 2H), 3.80 (s, 3H).

Example 134 (General procedure (H1))

(R) 3-[(2-{3-[5-Fluoro-2-(2-fluoro-6-methoxyphenoxy)phenyl]ureido}thiazole-4-carbonyl)amino]-2-hydroxy-propionic acid




- 5 ¹H NMR (DMSO): δ 12.62 (br, 1H), 11.32 (s, 1H), 9.23 (s 1H), 8.06 (dd, 1H), 7.79-7.76 (m, 2H), 7.33 (dd, 1H), 7.11-7.02 (m, 2H), 7.06 (dd, 1H), 6.75 (t, 1H), 6.54 (dd, 1H), 5.62 (br, 1H), 4.16 (t, 1H), 3.80 (s, 3H), 3.64-3.52 (m, 1H), 3.49-3.40 (m, 1H).


HPLC-MS (Method B): m/z = 509 (M+1); R_t = 3.49 min.

Example 135 (General procedure (H1))

- 10 2-[(2-{3-[5-Fluoro-2-(2-fluoro-6-methoxyphenoxy)phenyl]ureido}thiazole-4-carbonyl)amino]-3-hydroxy-propionic acid




- ¹H NMR (DMSO): δ 12.88 (br, 1H), 11.42 (s, 1H), 9.17 (s 1H), 8.06 (dd, 1H), 7.92 (d, 1H), 7.80 (s, 1H), 7.32 (dd, 1H), 7.11-7.01 (m, 2H), 6.75 (t, 1H), 6.54 (dd, 1H), 5.10 (br, 1H), 4.46 -
15 4.43 (m, 1H), 3.89 (m, 4H).

Example 136**1-(2-Cyclopentanecarbonyl-4-methylphenyl)-3-thiazol-2-yl-urea**

- 2-Cyclopentanecarbonyl-4-methylaniline (10.2 g) is prepared from p-toluidine (10.7 g, 0.1 mol) and cyclopentanecarbonitrile (9.5 g, 0.1 mol) following the general procedure I.
- 1-(2-Cyclopentanecarbonyl-4-methyl-phenyl)-3-thiazol-2-yl-urea (132 mg) is prepared from 2-cyclopentanecarbonyl-4-methylaniline (102 mg, 0.5 mmol) and 2-aminothiazole (0.060 g, 0.6 mmol) following the general procedure D.

LC-MS (m/z): 330 ($M+1$)⁺


- ¹H NMR (400 MHz, CDCl₃): δ 1.69 (m, 2H), 1.91 (m, 2H), 2.37 (s, 3H), 3.76 (m, 1H), 6.89 (d, 1H), 7.35 (d, 1H), 7.71 (s, 2H), 8.42 (d, 1H), 11.57 (br, 1H) ppm.

Example 137**1-(2-Isobutyryl-4-methylphenyl)-3-thiazol-2-yl-urea**

- 2-Isopropylcarbonyl-4-methylaniline (7.0 g) is prepared from p-toluidine (10.7 g, 0.1 mol) and isobutyronitrile (6.9 g, 0.1 mol) following the general procedure I. 1-(2-Isobutyryl-4-methylphenyl)-3-thiazol-2-yl-urea (113 mg) is prepared from 2-isopropylcarbonyl-4-methylaniline (88 mg, 0.5 mmol) and 2-aminothiazole (0.060 g, 0.6 mmol) following the general procedure D.

- LC-MS (m/z): 304 ($M+1$)⁺


¹H NMR (400 MHz, CDCl₃): δ 1.97 (d, 6H), 2.37 (s, 3H), 3.72 (m, 1H), 6.90 (d, 1H), 7.36 (d, 1H), 7.69 (s, 2H), 8.44 (s, 1H), 11.51 (br, 1H), 11.72 (br, 1H) ppm.

Example 138**1-[5-Fluoro-2-(3-methylbutyryl)phenyl]-3-thiazol-2-yl-urea**

- 2-[(2-Methyl)-propylcarbonyl]-5-fluoroaniline (6.82 g) is prepared from m-fluoro aniline (11.1 g, 0.1 mol) and 3-methylbutyronitrile (8.3 g, 0.1 mol) following the general procedure I. 1-[5-Fluoro-2-(3-methyl-butyl)-phenyl]-3-thiazol-2-yl-urea (128 mg) is prepared from 2-[(2-methyl)-propylcarbonyl]-5-fluoroaniline (97 mg, 0.5 mmol) and 2-aminothiazole (0.060 g, 0.6 mmol) following the general procedure D.


LC-MS (*m/z*): 322 (*M*+1)⁺

- ¹H NMR (400 MHz, CDCl₃): δ 0.99 (d, 6H), 2.23 (m, 1H), 2.83 (d, 2H), 6.75 (m, 1H), 6.93 (d, 1H), 7.62 (d, 1H), 7.92 (dd, 1H), 8.43 (dd, 1H), 11.60 (br, 1H), 12.04 (br, 1H) ppm.

Example 139**1-[5-Methyl-2-(3-methylbutyryl)phenyl]-3-thiazol-2-yl-urea**

- 2-[2-Methylpropylcarbonyl]-5-methylaniline (6.8 g) is prepared from m-toluidine (10.2 g, 0.1 mol) and isobutyronitrile (8.3 g, 0.1 mol) following the general procedure I. 1-[5-Methyl-2-(3-methylbutyryl)phenyl]-3-thiazol-2-yl-urea (114 mg) is prepared from 2-[2-methylpropylcarbonyl]-5-methylaniline (95 mg, 0.5 mmol) and 2-aminothiazole (0.060 g, 0.6 mmol) following the general procedure D.
- LC-MS (*m/z*): 318 (*M*+1)⁺


¹H NMR (400 MHz, CDCl₃): δ 0.98 (d, 6H), 2.31 (m, 1H), 2.37 (s, 3H), 2.44 (d, 2H), 6.91 (m, 1H), 7.27 (m, 1H), 7.51 (d, 1H), 7.68 (dd, 1H), 8.45 (d, 1H), 11.58 (br, 1H), 11.60 (br, 1H) ppm.

Example 140**5 [3-Cyclopentanecarbonyl-4-(3-thiazol-2-yl-ureido)phenyl]acetic acid ethyl ester**

(4-Amino-3-cyclopentanecarbonylphenyl)acetic acid ethyl ester (2.7 g) is prepared from (4-aminophenyl)acetic acid ethyl ester (18.0 g, 0.1 mol) and cyclopentanecarbonitrile (9.5 g, 0.1 mol) following the general procedure I. [3-Cyclopentanecarbonyl-4-(3-thiazol-2-yl-ureido)phenyl]acetic acid ethyl ester (140 mg) is prepared from (4-amino-3-cyclopentanecarbonylphenyl)acetic acid ethyl ester (138 mg, 0.5 mmol) and 2-aminothiazole (0.060 g, 0.6 mmol) following the general procedure D.

LC-MS (*m/z*): 402 (*M*+1)⁺

¹H NMR (400 MHz, CDCl₃): δ 1.12 (t, 3H), 1.73 (m, 2H), 1.82 (m, 2H), 3.27 (m, 1H), 3.62 (s, 2H), 4.15 (q, 2H), 6.89 (d, 1H), 7.44 (dd, 1H), 7.62 (d, 1H), 7.86 (s, 1H), 8.51 (s, 1H), 11.62 (br, 1H) ppm.

Example 141**[3-Cyclopentanecarbonyl-4-(3-thiazol-2-yl-ureido)phenyl]acetic acid**


[3-Cyclopentanecarbonyl-4-(3-thiazol-2-yl-ureido)phenyl]acetic acid (168 mg) is prepared from [3-cyclopentanecarbonyl-4-(3-thiazol-2-yl-ureido)phenyl]acetic acid ethyl ester (201 mg, 0.5 mmol) following the general procedure J.

LC-MS (*m/z*): 374 (*M*+1)⁺

¹H NMR (400 MHz, DMSO-d₆): δ 1.66 (m, 2H), 1.96 (m, 2H), 3.71 (s, 2H), 3.92 (m, 1H), 7.05 (d, 1H), 7.38 (d, 1H), 7.56 (dd, 1H), 8.08 (s, 1H), 8.55 (s, 1H), 11.26 (br, 1H) ppm.

Example 142

2-[3-Cyclopentanecarbonyl-4-(3-thiazol-2-yl-ureido)phenyl]-N-methylacetamide



5


2-[3-Cyclopentanecarbonyl-4-(3-thiazol-2-yl-ureido)-phenyl]-N-methylacetamide (154 mg) is prepared from [3-cyclopentanecarbonyl-4-(3-thiazol-2-yl-ureido)phenyl]acetic acid (186 mg, 0.5 mmol) and 1 M solution of methylamine in THF (0.5 ml, 0.5 mmol) following the general procedure K.

10 LC-MS (*m/z*): 387 (*M*+1)⁺

¹H NMR (400 MHz, CDCl₃/DMSO-d₆): δ 1.51 (m, 2H), 1.73 (m, 2H), 2.58 (s, 3H), 3.36 (s, 2H), 3.59 (m, 1H), 6.67 (m, 1H), 7.21 (m, 2H), 7.71 (s, 1H), 8.26 (bs, 1H), 10.95 (br, 1H) ppm.

Example 143

15 {2-[3-(2-Cyclopentanecarbonyl-4-methyl-phenyl)-ureido]-thiazol-4-yl}-acetic acid ethyl ester



{2-[3-(2-Cyclopentanecarbonyl-4-methyl-phenyl)-ureido]-thiazol-4-yl}-acetic acid ethyl ester (154 mg) is prepared from (2-amino-5-methylphenyl)(cyclopentyl)methanone (102 mg, 0.5 mmol) and ethyl-2-amino-4-thiazolyl acetate (93 mg, 0.5 mmol) following the general procedure D.


20

LC-MS (m/z): 416 ($M+1$)⁺

¹H NMR (400 MHz, CDCl₃): δ 1.23 (t, 3H), 1.67 (m, 2H), 1.87 (m, 2H), 2.34 (s, 3H), 3.70 (s, 2H), 4.18 (m, 3H), 6.68 (s, 1H), 7.25 (s, 2H), 7.32 (d, 1H), 7.67 (s, 1H), 8.45 (s, 1H), 9.75 (br, 1H), 11.51 (br, 1H) ppm.

5 Example 144

{2-[3-(2-Cyclopentanecarbonyl-4-methyl-phenyl)-ureido]-thiazol-4-yl}acetic acid




- 10 {2-[3-(2-Cyclopentanecarbonyl-4-methylphenyl)-ureido]-thiazol-4-yl}-acetic acid (198 mg) is prepared from {2-[3-(2-cyclopentanecarbonyl-4-methylphenyl)-ureido]-thiazol-4-yl}-acetic acid ethyl ester (208 mg, 0.5 mmol) following the general procedure J.

LC-MS (m/z): 388 ($M+1$)⁺

¹H NMR (400 MHz, DMSO- d_6): δ 1.73 (m, 2H), 1.86 (m, 2H), 2.32 (s, 3H), 3.55 (s, 2H), 3.86 (m, 1H), 6.84 (s, 1H), 7.36 (d, 1H), 7.84 (d, 1H), 8.15 (d, 1H), 10.62 (br, 1H), 11.92 (br, 1H), 12.24 (br, 1H) ppm.

15 Example 145

{3-Cyclopentanecarbonyl-4-{3-(4-ethoxycarbonylmethylthiazol-2-yl)-ureido}phenyl}acetic acid ethyl ester




{3-Cyclopentanecarbonyl-4-{3-(4-ethoxycarbonylmethyl-thiazol-2-yl)-ureido}-phenyl}-acetic acid ethyl ester (205 mg) is prepared from (4-amino-3-cyclopentanecarbonyl-phenyl)acetic acid ethyl ester (138 mg, 0.5 mmol) and ethyl-2-amino-4-thiazolyl acetate (93 mg, 0.5 mmol) following the general procedure D.

5 LC-MS (*m/z*): 488 (*M*+1)⁺

¹H NMR (400 MHz, CDCl₃): δ 1.24 (t, 6H), 1.67 (m, 2H), 1.88 (m, 2H), 2.34 (s, 3H), 3.61 (s, 2H), 3.71 (s, 2H), 3.73 (m, 1H), 4.17 (m, 4H), 4.18 (m, 3H), 6.69 (s, 1H), 7.42 (d, 2H), 7.85 (s, 1H), 8.53 (s, 1H), 11.64 (br, 1H) ppm.

Example 146

10 1-(2-Cyclopentanecarbonyl-4-methyl-phenyl)-3-(5-methanesulfonyl-thiazol-2-yl)-urea




1-(2-Cyclopentanecarbonyl-4-methyl-phenyl)-3-(5-methanesulfonyl-thiazol-2-yl)-urea (149 mg) is prepared from 2-cyclopentanecarbonyl-4-methylaniline (102 mg, 0.5 mmol) and 5-methanesulfonyl-thiazol-2-yl-amine (106 mg, 0.6 mmol) following the general procedure D.

15 LC-MS (*m/z*): 408 (*M*+1)⁺

¹H NMR (400 MHz, CDCl₃): δ 1.65-1.77 (m, 4H), 1.88-1.97 (m, 4H), 2.40 (s, 3H), 3.21 (s, 3H), 3.81 (p, 1H), 7.41 (dd, 1H), 7.78 (s, 1H), 8.31 (d, 1H), 8.43 (d, 1H), 11.57 (br, 1H), 11.95 (br, 1H) ppm.

Example 147

20 2-[3-(2-Cyclopentanecarbonyl-4-methylphenyl)-ureido]-thiazole-4-carboxylic acid ethyl ester




2-[3-(2-Cyclopentanecarbonyl-4-methylphenyl)-ureido]-thiazole-4-carboxylic acid ethyl ester (3.6 g) is prepared from 2-cyclopentanecarbonyl-4-methyl aniline (2.04 g, 10 mmol) and ethyl-2-amino-4-thiazole acetate (1.72 g, 10 mmol) following the general procedure D.

5 LC-MS (m/z): 402 ($M+1$)⁺

¹H NMR (400 MHz, CDCl₃): δ 1.35 (t, 3H), 1.62-1.78 (m, 4H), 1.83-1.96 (m, 4H), 2.37 (s, 3H), 3.74 (p, 1H), 4.33 (q, 2H), 7.36 (d, 1H), 7.72 (s, 1H), 7.81 (s, 1H), 8.44 (d, 1H), 9.25 (br, 1H), 11.84 (br, 1H) ppm.

Example 148


10 2-[3-(2-Cyclopentanecarbonyl-4-methylphenyl)-ureido]-thiazole-4-carboxylic acid



2-[3-(2-Cyclopentanecarbonyl-4-methylphenyl)-ureido]-thiazole-4-carboxylic acid (3.2 g) is prepared from 2-[3-(2-cyclopentanecarbonyl-4-methyl-phenyl)-ureido]-thiazole-4-carboxylic acid ethyl ester (3.6 g, 8.95 mmol) following the general procedure J.

15 LC-MS (m/z): 374 ($M+1$)⁺


¹H NMR (400 MHz, DMSO- d_6): δ 1.58 (m, 4H), 1.73 (m, 4H), 2.29 (s, 3H), 3.72 (m, 1H), 7.13 (s, 1H), 7.28 (d, 1H), 7.63 (d, 1H), 7.72 (br, 1H), 7.94 (s, 1H), 8.22 (br, 1H), 10.92 (br, 1H) ppm.

Example 149**2-[3-(2-Cyclopentanecarbonyl-4-methylphenyl)-ureido]-thiazole-4-carboxamide**

2-[3-(2-Cyclopentanecarbonyl-4-methylphenyl)-ureido]-thiazole-4-carboxamide (145 mg) is prepared from 2-[3-(2-cyclopentanecarbonyl-4-methylphenyl)-ureido]-thiazole-4-carboxylic acid (187 mg, 0.5 mmol) following the general procedure K.

LC-MS (m/z): 373 ($M+1$)⁺

¹H NMR (400 MHz, DMSO- d_6): δ 1.61 (m, 4H), 1.73 (m, 2H), 1.88 (m, 2H), 2.32 (s, 3H), 3.86 (m, 1H), 7.38 (d, 1H), 7.56 (s, 1H), 7.68 (s, 1H), 7.84 (s, 1H), 7.93 (s, 1H), 8.14 (d, 1H), 10.73 (br, 1H), 11.86 (br, 1H) ppm.

Example 150**2-{2-[3-(2-Cyclopentanecarbonyl-4-methylphenyl)-ureido]-thiazole-4-yl}-acetamide**


2-{2-[3-(2-Cyclopentanecarbonyl-4-methylphenyl)-ureido]-thiazole-4-yl}-acetamide (325 mg) is prepared from 2-[3-(2-cyclopentanecarbonyl-4-methylphenyl)-ureido]-thiazole-4-yl acetic acid (386 mg, 1.0 mmol) following the general procedure K.

LC-MS (m/z): 387 ($M+1$)⁺

¹H NMR (400 MHz, DMSO- d_6): δ 1.60 (m, 4H), 1.73 (m, 2H), 1.87 (m, 2H), 2.31 (s, 3H), 3.38 (s, 2H), 3.88 (m, 1H), 6.77 (s, 1H), 6.97 (s, 1H), 7.38 (d, 2H), 7.83 (s, 1H), 8.15 (d, 1H), 10.63 (br, 1H), 11.81 (br, 1H) ppm.

Example 151


2-{2-[3-(2-Cyclopentanecarbonyl-4-methylphenyl)-ureido]-thiazol-4-yl}-N-methyl-acetamide



- 2-{2-[3-(2-Cyclopentanecarbonyl-4-methylphenyl)-ureido]-thiazol-4-yl}-N-methyl-acetamide (346 mg) is prepared from 2-[3-(2-cyclopentanecarbonyl-4-methyl-phenyl)-ureido]-thiazol-4-yl acetic acid (386 mg, 1.0 mmol) following the general procedure K.
- LC-MS (m/z): 401 ($M+1$)⁺
- ¹H NMR (400 MHz, DMSO- d_6): δ 1.70 (m, 4H), 1.88 (m, 2H), 1.94 (m, 2H), 2.63 (s, 3H), 2.79 (d, 3H), 3.62 (s, 2H), 3.77 (p, 1H), 6.65 (s, 1H), 6.77 (br, 1H), 7.35 (dd, 1H), 7.71 (s, 1H), 8.43 (d, 1H), 9.15 (br, 1H), 11.70 (br, 1H) ppm.

Example 152

4-(2-{2-[3-(2-Cyclopentanecarbonyl-4-methylphenyl)-ureido]-thiazol-4-yl}-acetyl)-1-methyl-piperazinium chloride



- 4-(2-{2-[3-(2-Cyclopentanecarbonyl-4-methylphenyl)-ureido]-thiazol-4-yl}-acetyl)-1-methyl-piperazine (337 mg) is prepared from 2-[3-(2-cyclopentanecarbonyl-4-methyl-phenyl)-

ureido]-thiazol-4-yl)acetic acid (386 mg, 1.0 mmol) and N-methylpiperazine following the general procedure K.


4-(2-{2-[3-(2-Cyclopentanecarbonyl-4-methylphenyl)-ureido]-thiazol-4-yl}-acetyl)-1-methyl-piperazinium chloride (392 mg) is prepared from 4-(2-{2-[3-(2-cyclopentanecarbonyl-4-methylphenyl)-ureido]-thiazol-4-yl}-acetyl)-1-methyl-piperazine (386 mg, 1.0 mmol) by treatment with anhydrous hydrogen chloride (5 ml, 4.0 M solution in dioxane) followed by collection of the solid product.

LC-MS (*m/z*): 471 (*M*+1)⁺

¹H NMR (400 MHz, DMSO-*d*₆): δ 1.60 (m, 4H), 1.72 (m, 2H), 1.87 (m, 2H), 2.32 (s, 3H), 2.47 (s, 3H), 2.76 (d, 2H), 2.94 (m, 2H), 3.37 (m, 2H), 3.73 (d, 2H), 3.84 (m, 2H), 4.22 (d, 1H), 4.44 (d, 1H), 6.83 (s, 1H), 7.36 (d, 1H), 7.82 (s, 1H), 8.11 (d, 1H), 10.63 (br, 1H), 10.82 (br, 1H) ppm.

Example 153

1-[4-Methyl-2-(2-methylpropoxy)phenyl]-3-thiazol-2-yl-urea




3-(2-Methylpropoxy)-4-nitrotoluene (0.78 g) is prepared from 2-methylpropanol (0.46 ml, 5.0 mmol) and 3-fluoro-4-nitrotoluene (0.77 g, 5.0 mmol) following the general procedure G. This is reduced to afford 4-methyl-2-(2-methylpropoxy)aniline (0.47 g) following general procedure C. N-[4-Methyl-2-(2-methylpropoxy)phenyl]-N'-(thiazol-2-yl)urea (210 mg) is prepared from 4-methyl-2-(2-methylpropoxy)aniline (179 mg, 1.0 mmol) and 2-aminothiazole (100 mg, 1.0 mmol) following the general procedure D.

LC-MS (*m/z*): 306 (*M*+1)⁺

¹H NMR (400 MHz, CDCl₃): δ 1.01 (d, 6H), 2.16 (m, 1H), 2.32 (s, 3H), 3.78 (d, 2H), 6.70 (s, 1H), 6.75 (d, 1H), 6.86 (d, 1H), 7.40 (d, 1H), 8.07 (d, 1H), 9.30 (br, 1H), 10.72 (br, 1H) ppm.

Example 154

{2-[3-(4-Methyl-2-[2-methylpropoxy]phenyl)-ureido]-thiazol-4-yl}-acetic acid




- 5 {2-[3-(4-Methyl-2-[2-methylpropoxy]phenyl)-ureido]-thiazol-4-yl}-acetic acid ethyl ester (485 mg) is prepared from 4-methyl-2-(2-methylpropoxy)aniline (360 mg, 2.0 mmol) and ethyl 2-amino-4-thiazolylacetate (372 mg, 2.0 mmol) following the general procedure D. Hydrolysis of this ester following general procedure J gave {2-[3-(4-methyl-2-[2-methylpropoxy]phenyl)-ureido]-thiazol-4-yl}-acetic acid (400 mg).

LC-MS (m/z): 464 ($M+1$)⁺

- 10 ¹H NMR (400 MHz, DMSO- d_6): δ 1.01 (d, 6H), 2.07 (m, 1H), 2.23 (s, 3H), 3.53 (s, 2H), 3.77 (d, 2H), 6.67 (d, 1H), 6.81 (s, 2H), 7.91 (d, 1H), 8.01 (br, 1H), 11.45 (br, 1H), 12.35 (br, 1H) ppm.

Example 155

{2-[3-(4-Methyl-2-[2-methylpropoxy]phenyl)-ureido]-thiazol-4-yl}-N-methyl-acetamide



15


{2-[3-(4-Methyl-2-[2-methylpropoxy]phenyl)-ureido]-thiazol-4-yl}-N-methyl-acetamide (150 mg) is prepared from {2-[3-(4-methyl-2-[2-methylpropoxy]phenyl)-ureido]-thiazol-4-yl}-acetic acid (182 mg, 0.5 mmol) following the general procedure K.

LC-MS (m/z): 477 ($M+1$)⁺

¹H NMR (400 MHz, DMSO-d₆): δ 1.00 (d, 6H), 2.06 (m, 1H), 2.19 (s, 3H), 2.48 (s, 3H), 3.40 (s, 2H), 3.68 (d, 2H), 6.54 (s, 1H), 6.59 (d, 2H), 7.40 (br, 1H), 7.91 (d, 1H), 8.08 (br, 1H), 11.27 (br, 1H) ppm.

5 EXAMPLE 157

1-(2-Cyclopentanecarbonyl-4-methyl-phenyl)-3-(4-methyl-thiazol-2-yl)-urea




- 1-(2-Cyclopentanecarbonyl-4-methyl-phenyl)-3-(4-methyl-thiazol-2-yl)-urea (66 mg, 77%) was prepared from (2-amino-5-methyl-phenyl)-cyclopentyl-methanone (51 mg, 0.25 mmol) following the general procedure D.
- LC-MS (*m/z*): 344 (M+1)⁺; ¹H NMR (400 MHz, CDCl₃): δ 1.60-1.75 (m, 4H), 1.80-1.94 (m, 4H), 2.32 (s, 3H), 2.36 (s, 3H), 3.73 (p, 1H), 6.44 (s, 1H), 7.28 (d, 1H), 7.35 (d, 1H), 7.67 (s, 1H), 8.44 (br, 1H), and 11.66 (br, 1H).

15

Example 158

1-(2-Cyclopentanecarbonyl-4-methyl-phenyl)-3-(4-methoxymethyl-thiazol-2-yl)-urea



20


1-(2-Cyclopentanecarbonyl-4-methyl-phenyl)-3-(4-methoxymethyl-thiazol-2-yl)-urea (64 mg, 69%) was prepared from (2-amino-5-methyl-phenyl)-cyclopentyl-methanone (50 mg, 0.25 mmol) and 4-methoxymethyl-thiazol-2-ylamine (54 mg, 0.375 mmol) following the general procedure D to give title compound.

LC-MS (m/z): 373 ($M+1$)⁺; ¹H NMR (400 MHz, CDCl₃): δ 1.70 (m, 4H), 1.88 (m, 4H), 2.36 (s, 3H), 3.41 (s, 2H), 3.72 (m, 1H), 4.44 (s, 3H), 6.78 (dd, 1H), 7.32 (d, 1H), 7.70 (s, 1H), 8.42 (d, 1H), 10.52 (br, 1H), and 11.64 (br, 1H).

5

Example 159

1-(2-Cyclopentanecarbonyl-4-methyl-phenyl)-3-(4-formyl-thiazol-2-yl)-urea



10

To 1-(2-cyclopentanecarbonyl-4-methyl-phenyl)-3-[4-hydroxymethyl-thiazol-2-yl]-urea (1.8 g, 5 mmol) in CH₂Cl₂ (40 mL) at 0 °C was added Et₃N (2.88 mL, 20 mmol), dimethyl sulfoxide (10 mL) followed by sulfur trioxide-pyridine (2.38 g, 15.0 mmol). The reaction mixture was stirred for 1h and then poured into water (30 mL). The mixture was extracted with CH₂Cl₂ and the organic layers was washed with 1.0 N ammonium chloride (2 x 20 mL), water (2 x 20 mL), brine (1 x 20 mL), dried (Na₂SO₄) and concentrated to give a solid. The solid was purified by column chromatography (silica, Hexanes/ EtOAc, 20-50%) to obtain 1-(2-cyclopentanecarbonyl-4-methyl-phenyl)-3-(4-formyl-thiazol-2-yl)-urea (1.59 g, 86%) as a white solid.


15

LC-MS (m/z): 358 ($M+1$)⁺; ¹H NMR (400 MHz, DMSO-d₆): δ 1.64 (m, 4H), 1.74 (m, 2H), 1.90 (m, 2H), 2.35 (s, 3H), 3.89 (p, 1H), 7.41 (d, 1H), 7.88 (s, 1H), 8.17 (d, 1H), 8.24 (s, 1H), 9.76 (s, 1H), 10.75 (s, 1H), and 12.20 (br, 1H).

20

Example 160

1-(2-Cyclopentanecarbonyl-4-methyl-phenyl)-3-(4-hydroxymethyl-thiazol-2-yl)-urea




1-(2-Cyclopentanecarbonyl-4-methyl-phenyl)-3-(4-hydroxymethyl-thiazol-2-yl)-urea (2.66 g, 92%) was prepared from acetic acid 2-[3-(2-cyclopentanecarbonyl-4-methyl-phenyl)-ureido]-thiazol-4-ylmethyl ester (3.2 g, 8.0 mmol) following the general procedure J to give
5 title compound.

LC-MS (m/z): 360 ($M+1$)⁺; ¹H NMR (400 MHz, DMSO- d_6): δ 1.75 (m, 4H), 1.89 (m, 4H), 2.34 (s, 3H), 3.88 (m, 1H), 4.43 (d, 2H), 5.20 (t, 1H), 6.82 (s, 1H), 7.39 (d, 1H), 7.85 (s, 1H), 8.18 (d, 1H), 10.66 (br, 1H), and 11.72 (br, 1H).

10

Example 161

1-(4-Chloromethyl-thiazol-2-yl)-3-(2-cyclopentanecarbonyl-4-methyl-phenyl)-urea



15


1-(4-Chloromethyl-thiazol-2-yl)-3-(2-cyclopentanecarbonyl-4-methyl-phenyl)-urea (2.16 g, 57.1%) was prepared from (2-amino-5-methyl-phenyl)-cyclopentyl-methanone (2.03 g, 10.0 mmol) and 4-chloromethyl-thiazol-2-ylamine (1.72 g, 10.0 mmol) following the general procedure D.

LC-MS (m/z): 378 ($M+1$)⁺; ¹H NMR (400 MHz, CDCl₃): δ 1.68 (m, 4H), 1.88 (m, 4H), 2.36 (s, 3H), 3.76 (m, 1H), 4.52 (s, 2H), 6.88 (s, 1H), 7.34 (d, 1H), 7.73 (s, 1H), 8.42 (d, 1H), 11.36 (br, 1H), and 11.75 (br, 1H).

20

Example 162

1-(4-Aminomethyl-thiazol-2-yl)-3-(2-cyclopentanecarbonyl-4-methyl-phenyl)-urea



To a solution of thiourea (7.6 g, 100 mmol) in methanol (150 mL) was added 1,3-dichloroacetone (12.7 g, 100 mmol) and the reaction mixture stirred at rt. for 3-4 hours. The mixture was concentrated *in vacuo* to give the crude product as hydrochloride salt. This salt
 5 was then washed with Et₂O (3 x 200 mL) and concentrated *in vacuo* to afford 4-chloromethyl-thiazol-2-ylamine in 95-100% yield.

To a solution of 4-chloromethyl-thiazol-2-ylamine (7.45 g, 50.0 mmol) in dioxane:water mixture (8:2) was added sodium azide (4.87 g, 75.0 mmol) and the reaction mixture was refluxed for 2-3 h. The mixture was concentrated *in vacuo* to remove all dioxane
 10 and the residue was dissolved in EtOAc (200 mL). The organic layer was washed with water (2 x 200 mL), brine (2 x 200 mL), dried over (Na₂SO₄) and concentrated *in vacuo* to give 4-azidomethyl-thiazol-2-ylamine in 70-90% yield.


(4-Azidomethyl-thiazol-2-ylamine) was subjected to urea formation by following general procedure D to give the desired urea (1-(4-azidomethyl-thiazol-2-yl)-3-(2-cyclopentanecarbonyl-4-methyl-phenyl)-urea) in 65-85% yield.
 15

To a solution of (1-(4-azidomethyl-thiazol-2-yl)-3-(2-cyclopentanecarbonyl-4-methyl-phenyl)-urea) (7.7 g, 20 mmol) in EtOH (100 mL) was added catalytic amount of palladium on charcoal (200 mg) and the reaction mixture was hydrogenated (1 atmos) for 3-4 hours to give 1-(4-aminomethyl-thiazol-2-yl)-3-(2-cyclopentanecarbonyl-4-methyl-phenyl)-urea. This
 20 crude product was then purified by flash column chromatography with DCM:EtOAc (80:20 to 50:50) as eluent to give the desired amine (5.7 g, 80%).

LC-MS (*m/z*): 359 (M+1)⁺; ¹H NMR (400 MHz, CDCl₃): δ 1.48 (m, 2H), 1.64 (m, 4H), 1.82 (m, 2H), 2.30 (s, 3H), 3.46 (d, 2H), 3.54 (p, 1H), 3.76 (t, 2H), 6.84 (s, 1H), 7.07 (s, 1H), 7.76 (d, 1H), 8.32 (d, 1H), 9.64 (br, 1H), and 11.18 (br, 1H).

Example 163

1-(2-Cyclopentanecarbonyl-4-methyl-phenyl)-3-(4-dimethylaminomethyl-thiazol-2-yl)-urea




1-(2-Cyclopentanecarbonyl-4-methyl-phenyl)-3-(4-dimethylaminomethyl-thiazol-2-yl)-
urea (79 mg, 82%) was prepared from 2-[3-(2-1-(2-cyclopentanecarbonyl-4-methyl-phenyl)-
5 3-(4-formyl-thiazol-2-yl)-urea (89 mg, 0.25 mmol) and (30 mg, 0.3 mmol) and dimethylamine
(0.1 mL, 2.0 M. solution in THF) following the general procedure O to give title compound.

LC-MS (*m/z*): 387 (*M*+1)⁺; ¹H NMR (400 MHz, CDCl₃): δ 1.62-1.72 (m, 4H), 1.78-1.88
(m, 4H), 2.23 (s, 6H), 2.34 (s, 3H), 3.53 (s, 2H), 3.70 (p, 1H), 6.64 (s, 1H), 7.30 (d, 1H), 7.64
(s, 1H), 8.32 (d, 1H), 10.25 (br, 1H), and 11.35 (br, 1H).

10

Example 164

N-{2-[3-(2-Cyclopentanecarbonyl-4-methyl-phenyl)-ureido]-thiazol-4-ylmethyl}-
acetamide




N-{2-[3-(2-Cyclopentanecarbonyl-4-methyl-phenyl)-ureido]-thiazol-4-ylmethyl}-acetamide (88 mg, 88%) was prepared from 1-(4-aminomethyl-thiazol-2-yl)-3-(2-cyclopentanecarbonyl-4-methyl-phenyl)-urea (90 mg, 0.25 mmol) and acetyl chloride (0.02 mL, 0.25 mmol) following the general procedure T.

- 5 LC-MS (*m/z*): 401 (*M*+1)⁺; ¹H NMR (400 MHz, CDCl₃): δ 1.56 (m, 4H), 1.72 (m, 2H), 1.78 (m, 2H), 2.22 (s, 3H), 2.82 (s, 3H), 3.18 (t, 1H), 3.64 (m, 1H), 4.07 (s, 2H), 6.55 (s, 1H), 7.22 (d, 1H), 7.53 (s, 1H), 8.18 (d, 1H), 10.18 (br, 1H), and 11.58 (br, 1H).

Example 165

- 10 1-{4-[2-(tert-Butyl-dimethyl-silanyloxy)-ethyl]-thiazol-2-yl}-3-(2-cyclopentanecarbonyl-4-methyl-phenyl)-urea




- 15 1-{4-[2-(tert-Butyl-dimethyl-silanyloxy)-ethyl]-thiazol-2-yl}-3-(2-cyclopentanecarbonyl-4-methyl-phenyl)-urea (425 mg, 87%) was prepared from (2-amino-5-methyl-phenyl)-cyclopentyl-methanone (203 mg, 1.00 mmol) and 4-[2-(tert-butyl-dimethyl-silanyloxy)-ethyl]-thiazol-2-ylamine (310 mg, 1.20 mmol) following the general procedure D.

- 20 LC-MS (*m/z*): 488 (*M*+1)⁺; ¹H NMR (400 MHz, CDCl₃): δ 0.10 (s, 9H), 0.85 (s, 6H), 1.70-1.78 (m, 4H), 1.82-1.94 (m, 4H), 2.35 (s, 3H), 2.84 (t, 2H), 3.73 (p, 1H), 3.87 (t, 2H), 6.52 (s, 1H), 6.60 (d, 1H), 7.32 (d, 1H), 7.72 (s, 1H), 8.48 (br, 1H), 11.68 (br, 1H).

Example 166

Acetic acid 2-[3-(2-cyclopentanecarbonyl-4-methyl-phenyl)-ureido]-thiazol-4-ylmethyl ester



Acetic acid 2-[3-(2-cyclopentanecarbonyl-4-methyl-phenyl)-ureido]-thiazol-4-ylmethyl ester (3.66 g, 91%) was prepared from (2-amino-5-methyl-phenyl)-cyclopentyl-methanone (2.03 g, 10.0 mmol) and acetic acid 2-amino-thiazol-4-ylmethyl ester (2.05 g, 12.0 mmol)


5 following the general procedure D.

LC-MS (m/z): 402 ($M+1$)⁺; ¹H NMR (400 MHz, CDCl₃): δ 1.70 (m, 4H), 1.86 (m, 4H), 2.10 (s, 3H), 2.37 (s, 3H), 3.75 (m, 1H), 5.09 (s, 2H), 6.87 (s, 1H), 7.35 (d, 1H), 7.71 (s, 1H), 8.45 (d, 1H), 9.20 (br, 1H), and 11.75 (br, 1H).

10

Example 167

1-(2-Cyclopentanecarbonyl-4-methyl-phenyl)-1-methyl-3-thiazol-2-yl-urea




1-(2-Cyclopentanecarbonyl-4-methyl-phenyl)-1-methyl-3-thiazol-2-yl-urea (287 mg, 83%) was prepared from cyclopentyl-(5-methyl-2-methylamino-phenyl)-methanone (217 mg, 1.00 mmol) following the general procedure D.

15

LC-MS (m/z): 344 ($M+1$)⁺; ¹H NMR (400 MHz, CDCl₃): δ 1.72 (m, 4H), 1.88 (m, 4H), 2.37 (s, 3H), 3.42 (s, 3H), 3.84 (p, 1H), 6.85 (d, 1H), 6.94 (d, 1H), 7.17 (d, 1H), 7.43 (d, 1H), 7.46 (s, 1H), and 9.62 (br, 1H).

EXAMPLE 168

1-(5-Chloro-thiazol-2-yl)-3-(2-cyclopentanecarbonyl-4-methyl-phenyl)-urea




5 1-(5-Chloro-thiazol-2-yl)-3-(2-cyclopentanecarbonyl-4-methyl-phenyl)-urea (109 mg, 60 %) was prepared from (2-amino-5-methyl-phenyl)-cyclopentyl-methanone (102 mg, 0.5 mmol) and 5-chloro-thiazol-2-ylamine (0.06 mg, 0.6 mmol) following the general procedure D.

10 LC-MS (*m/z*): 364 (*M*+1)⁺; ¹H NMR (400 MHz, CDCl₃): δ 1.72 (m, 4H), 1.93 (m, 4H), 2.38 (s, 3H), 3.80 (m, 1H), 7.37 (d, 1H), 7.64 (s, 1H), 7.74 (s, 1H), 8.44 (d, 1H), 11.69 (br, 1H).

Example 169

1-(5-Bromo-thiazol-2-yl)-3-(2-cyclopentanecarbonyl-4-methyl-phenyl)-urea




15 1-(5-Bromo-thiazol-2-yl)-3-(2-cyclopentanecarbonyl-4-methyl-phenyl)-urea (127 mg, 62%) was prepared from (2-amino-5-methyl-phenyl)-cyclopentyl-methanone (102 mg, 0.5 mmol) and 5-bromo-thiazol-2-ylamine (108 mg, 0.6 mmol) following the general procedure D.

20 LC-MS (*m/z*): 409 (*M*+1)⁺; ¹H NMR (400 MHz, CDCl₃): δ 1.72 (m, 4H), 1.93 (m, 4H), 2.38 (s, 3H), 3.78 (m, 1H), 7.38 (d, 1H), 7.66 (s, 1H), 7.74 (s, 1H), 8.44 (d, 1H), 11.67 (br, 1H).

Example 170

1-(5-Bromo-thiazol-2-yl)-3-(2-cyclopentanecarbonyl-5-fluoro-phenyl)-urea




1-(5-Bromo-thiazol-2-yl)-3-(2-cyclopentanecarbonyl-5-fluoro-phenyl)-urea (132 mg, 64%) was prepared from (2-amino-4-fluoro-phenyl)-cyclopentyl-methanone (104 mg, 0.5 mmol) and 5-bromo-thiazol-2-ylamine (108 mg, 0.6 mmol) following the general procedure D.

- 5 LC-MS (m/z): 413 ($M+1$)⁺; ¹H NMR (400 MHz, CDCl₃): δ 1.72 (m, 4H), 1.93 (m, 4H), 2.38 (s, 3H), 3.74 (m, 1H), 6.81 (m, 1H), 7.61 (s, 1H), 8.01 (dd, 1H), 8.42 (dd, 1H), 11.02 (br, 1H), 12.16 (br, 1H).

EXAMPLE 171

- 10 2-[3-(2-Cyclopentanecarbonyl-4-methyl-phenyl)-ureido]-4-methyl-thiazol e-5-carboxylic acid ethyl ester




2-[3-(2-Cyclopentanecarbonyl-4-methyl-phenyl)-ureido]-4-methyl-thiazole-5-carboxylic acid ethyl ester (154 mg, 80 %) was prepared from (2-amino-5-methyl-phenyl)-cyclopentyl-

- 15 methanone (102 mg, 0.5 mmol) and 2-amino-4-methyl-thiazole-5-carboxylic acid ethyl ester (112 mg, 0.6 mmol) following the general procedure D.

LC-MS (m/z): 416 ($M+1$)⁺; ¹H NMR (400 MHz, CDCl₃): δ 1.33 (t, 3H), 1.66 (m, 4H), 1.87 (m, 4H), 2.37 (s, 3H), 2.59 (s, 3H), 3.72 (m, 1H), 4.28 (q, 2H), 7.37 (s, 1H), 7.70 (s, 1H), 8.48 (s, 1H), 10.35 (br, 1H), 11.81 (br, 1H).

Example 172

2-[3-(2-Cyclopentanecarbonyl-4-methyl-phenyl)-ureido]-4-methyl-thiazol e-5-carboxylic acid




2-[3-(2-Cyclopentanecarbonyl-4-methyl-phenyl)-ureido]-4-methyl-thiazol e-5-carboxylic acid (198 mg, 95 %) was prepared from 2-[3-(2-cyclopentanecarbonyl-4-methyl-phenyl)-ureido]-4-methyl-thiazole-5-carboxylic acid ethyl ester (208 mg, 0.5 mmol) following the general procedure J.

LC-MS (m/z): 388 ($M+1$)⁺; ¹H NMR (400 MHz, CDCl₃): δ 1.66 (m, 4H), 1.74 (m, 2H), 1.94 (m, 2H), 2.61 (s, 3H), 2.65 (s, 3H), 3.88 (m, 1H), 7.42 (d, 1 H), 7.88 (s, 1H), 8.20 (d, 1 H), 10.25 (br, 1H), 12.22 (br, 2H).

Example 173

1-(2'-Amino-[4,4']bithiazolyl-2-yl)-3-(2-cyclopentanecarbonyl-4-methylphenyl)-urea




1-(2'-Amino-[4,4']bithiazolyl-2-yl)-3-(2-cyclopentanecarbonyl-4-methylphenyl)-urea (85 mg, 79%) was prepared from (2-Amino-5-methyl-phenyl)-cyclopentyl-methanone (51 mg, 0.25 mmol) following the general procedure D.

LC-MS (m/z): 428 ($M+1$)⁺; ¹H NMR (400 MHz, CDCl₃): δ 1.25 (m, 4H), 1.73 (m, 4H), 2.37 (s, 3H), 3.77 (p, 1H), 5.24 (br, 2H), 6.99 (s, 1H), 7.02 (s, 1H), 7.53 (dd, 1H), 7.58 (s, 1H), 7.72 (s, 1H), 8.42 (br, 1H), and 11.78 (br, 1H).

Example 174

2-{2-[3-(2-Cyclopentanecarbonyl-4-methyl-phenyl)-ureido]-thiazol-4-yl}-N-methoxy-N-methyl-acetamide




2-{2-[3-(2-Cyclopentanecarbonyl-4-methyl-phenyl)-ureido]-thiazol-4-yl}-N-methoxy-N-methyl-acetamide (88 mg, 82.%) was prepared from 2-[3-(2-cyclopentanecarbonyl-4-methyl-phenyl)-ureido]-thiazol-4-yl acetic acid (97 mg, 0.25 mmol) following the general procedure K.

LC-MS (m/z): 431 ($M+1$)⁺; ¹H NMR (400 MHz, CDCl₃): δ 1.72 (m, 4H), 1.93 (m, 4H), 2.35 (s, 3H), 2.80 (s, 2H), 3.70 (s, 3H), 3.85 (s, 3H), 4.21 (p, 1H), 6.67 (s, 1H), 6.91 (d, 1H), 7.54 (dd, 1H), 7.70 (d, 1H), 8.40 (br, 1H), and 11.52 (br, 1H).

Example 175

1-(2-Cyclopentanecarbonyl-4-methyl-phenyl)-3-[4-(2-morpholin-4-yl-2-oxo-ethyl)-thiazol-2-yl]-urea




1-(2-Cyclopentanecarbonyl-4-methyl-phenyl)-3-[4-(2-morpholin-4-yl-2-oxo-ethyl)-thiazol-2-yl]-urea (92 mg, 80%) was prepared from 2-[3-(2-cyclopentanecarbonyl-4-methyl-phenyl)-ureido]-thiazol-4-yl acetic acid (97 mg, 0.25 mmol) following the general procedure K.

LC-MS (*m/z*): 457 (*M*+1)⁺; ¹H NMR (400 MHz, CDCl₃): δ 1.64 (m, 4H), 1.82 (m, 2H), 1.88 (m, 2H), 2.34 (s, 3H), 2.96-3.14 (dd, 2H), 3.49 (t (2H), 3.60 (m, 4H), 3.72 (p, 1H), 3.76 (s, 2H), 6.60 (s, 1H), 7.33 (d, 1H), 7.67 (s, 1H), 8.42 (d, 1H), 8.70 (br, 1H), and 11.28 (br, 1H).

Example 176

2-[2-[3-(2-Cyclopentanecarbonyl-4-methyl-phenyl)-ureido]-thiazol-4-yl]-N-(4-methyl-piperazin-1-yl)-acetamide




2-[2-[3-(2-Cyclopentanecarbonyl-4-methyl-phenyl)-ureido]-thiazol-4-yl]-N-(4-methyl-piperazin-1-yl)-acetamide (105 mg, 87%) was prepared from 2-[3-(2-cyclopentanecarbonyl-4-methyl-phenyl)-ureido]-thiazol-4-yl acetic acid (97 mg, 0.25 mmol) following the general procedure K.

LC-MS (*m/z*): 485 (*M*+1)⁺; ¹H NMR (400 MHz, CDCl₃): δ 1.66 (m, 4H), 1.82 (m, 2H), 1.94 (m, 2H), 2.35 (s, 3H), 3.02 (m, 2H), 3.46 (s, 3H), 3.53 (t (2H), 3.64 (m, 4H), 3.76 (p, 1H), 3.80 (s, 2H), 5.36 (br, 1H), 6.56 (s, 1H), 7.34 (d, 1H), 7.69 (s, 1H), 8.46 (d, 1H), 8.82 (br, 1H), and 11.46 (br, 1H).

Example 177

2-[2-[3-(2-Cyclopentanecarbonyl-4-methyl-phenyl)-ureido]-thiazol-4-yl]-N-(2-morpholin-4-yl-ethyl)-acetamide



2-{2-[3-(2-Cyclopentanecarbonyl-4-methyl-phenyl)-ureido]-thiazol-4-yl}-N-(2-morpholin-4-yl-ethyl)-acetamide (108 mg, 86%) was prepared from 2-[3-(2-cyclopentanecarbonyl-4-methyl-phenyl)-ureido]-thiazol-4-yl acetic acid (97 mg, 0.25 mmol)


5 following the general procedure K.

LC-MS (m/z): 500 ($M+1$)⁺; ¹H NMR (400 MHz, CDCl₃): δ 1.22 (t, 2H), 1.35 (t, 4H), 1.70 (m, 2H), 1.86 (m, 2H), 1.94 (m, 2H), 2.32 (s, 3H), 2.88 (s, 2H), 2.96 (m, 4H), 3.22 (t, 4H), 3.77 (p, 1H), 5.30 (br, 1H), 6.58 (s, 1H), 7.34 (d, 1H), 7.90 (s, 1H), 8.38 (d, 1H), 9.14 (br, 1H), and 11.32 (br, 1H).

10

Example 178

2-[3-(2-Cyclopentanecarbonyl-4-methyl-phenyl)-ureido]-thiazole-4-carboxylic acid methylamide




2-[3-(2-Cyclopentanecarbonyl-4-methyl-phenyl)-ureido]-thiazole-4-carboxylic acid methylamide (77 mg, 80%) was prepared from 2-[3-(2-cyclopentanecarbonyl-4-methyl-phenyl)-ureido]-thiazol-4-yl acetic acid (97 mg, 0.25 mmol) following the general procedure K.

LC-MS (m/z): 387 ($M+1$)⁺; ¹H NMR (400 MHz, CDCl₃): δ 1.74 (m, 4H), 1.82 (m, 2H),
5 1.94 (m, 2H), 2.38 (s, 3H), 3.00 (d, 3H), 3.13 (m, 1H), 3.78 (p, 1H), 6.94 (s, 1H), 7.34 (d, 1H),
7.51 (br, 1H), 7.72 (d, 1H), 8.40 (d, 1H), and 11.88 (br, 1H).

Example 179

10 1-(2-Cyclopentanecarbonyl-4-methyl-phenyl)-3-[4-(morpholine-4-carbonyl)-thiazol-2-yl]-urea




1-(2-Cyclopentanecarbonyl-4-methyl-phenyl)-3-[4-(morpholine-4-carbonyl)-thiazol-2-yl]-urea (92 mg, 80%) was prepared from 2-[3-(2-cyclopentanecarbonyl-4-methyl-phenyl)-ureido]-thiazol-4-yl acetic acid (97 mg, 0.25 mmol) following the general procedure K.

15 LC-MS (m/z): 443 ($M+1$)⁺; ¹H NMR (400 MHz, CDCl₃): δ 1.70 (m, 4H), 1.90 (m, 4H),
2.36 (s, 3H), 3.12 (m, 2H), 3.76 (m, 4H), 3.82 (m, 1H), 6.69 (s, 1H), 7.34 (d, 1H), 7.69 (s, 1H),
8.38 (d, 1H), 9.83 (br, 1H), and 11.86 (br, 1H).

Example 180

20 N-(2-{2-[3-(2-Cyclopentanecarbonyl-4-methyl-phenyl)-ureido]-thiazol-4-yl}-acetyl)-methanesulfonamide




N-(2-{2-[3-(2-Cyclopentanecarbonyl-4-methyl-phenyl)-ureido]-thiazol-4-yl}-acetyl)-methanesulfonamide (88 mg, 76%) was prepared from 2-[3-(2-cyclopentanecarbonyl-4-methyl-phenyl)-ureido]-thiazol-4-yl acetic acid (97 mg, 0.25 mmol) following the general
5 procedure K.

LC-MS (m/z): 465 ($M+1$)⁺; ¹H NMR (400 MHz, CDCl₃): δ 1.72 (m, 4H), 1.92 (m, 4H), 2.37 (s, 3H), 3.13 (s, 3H), 3.30 (s, 2H), 3.81 (p, 1H), 4.88 (br, 1H), 6.66 (s, 1H), 7.36 (d, 1H), 7.72 (d, 1H), 8.46 (d, 1H), 10.70 (br, 1H), and 11.72 (br, 1H).

10

Example 181

N-{2-[3-(2-Cyclopentanecarbonyl-4-methyl-phenyl)-ureido]-thiazole-4-carbonyl}-methanesulfonamide



15


N-{2-[3-(2-Cyclopentanecarbonyl-4-methyl-phenyl)-ureido]-thiazole-4-carbonyl}-methanesulfonamide (103 mg, 91%) was prepared from 2-[3-(2-cyclopentanecarbonyl-4-methyl-phenyl)-ureido]-thiazol-4-yl acetic acid (97 mg, 0.25 mmol) following the general procedure K.

LC-MS (m/z): 451 ($M+1$)⁺; ¹H NMR (400 MHz, CDCl₃): δ 1.74 (m, 4H), 1.94 (m, 4H), 2.32 (s, 3H), 3.11 (s, 3H), 3.32 (p, 1H), 4.66 (br, 1H), 6.62 (s, 1H), 7.32 (d, 1H), 7.62 (d, 1H), 8.22 (br, 1H), 8.40 (d, 1H), and 11.68 (br, 1H).

5

Example 182

1-(2-Cyclopentanecarbonyl-4-methyl-phenyl)-3-{4-[3-(2-dimethylamino-ethyl)-ureidomethyl]-thiazol-2-yl}-urea




10 1-(2-Cyclopentanecarbonyl-4-methyl-phenyl)-3-{4-[3-(2-dimethylamino-ethyl)-ureidomethyl]-thiazol-2-yl}-urea (86 mg, 73%) was prepared from 2-[3-(2-cyclopentanecarbonyl-4-methyl-phenyl)-ureido]-thiazol-4-yl acetic acid (97 mg, 0.25 mmol) following the general procedure M.

LC-MS (m/z): 473 ($M+1$)⁺; ¹H NMR (400 MHz, CDCl₃): δ 1.40 (t, 2H), 1.72 (m, 4H), 1.92 (m, 4H), 2.16 (s, 6H), 2.35 (s, 3H), 2.88 (m, 2H), 3.13 (s, 2H), 3.70 (p, 1H), 6.90 (s, 1H), 7.36 (d, 1H), 7.67 (s, 1H), 8.20 (d, 1H), 8.44 (br, 1H), 8.92 (br, 1H), 10.26 (br, 1H), and 11.90 (br, 1H).

15

Example 183

20 (3-{2-[3-(2-Cyclopentanecarbonyl-4-methyl-phenyl)-ureido]-thiazol-4-ylmethyl}-ureido)-acetic acid




(3-{2-[3-(2-Cyclopentanecarbonyl-4-methyl-phenyl)-ureido]-thiazol-4-ylmethyl}-ureido)-acetic acid (96 mg, 83%) was prepared from 2-[3-(2-cyclopentanecarbonyl-4-methyl-phenyl)-ureido]-thiazol-4-yl acetic acid (97 mg, 0.25 mmol) and glycine methyl ester (48 mg, 0.5 mmol) following the general procedure M followed by hydrolysis using the general procedure J.

LC-MS (m/z): 460 ($M+1$)⁺; ¹H NMR (400 MHz, CDCl₃): δ 1.70 (m, 4H), 1.90 (m, 4H), 2.38 (s, 3H), 3.49 (s, 2H), 3.71 (s, 2H), 3.84 (p, 1H), 6.78 (s, 1H), 7.34 (d, 1H), 7.73 (s, 1H), 8.36 (d, 1H), 8.42 (br, 1H), 8.94 (br, 1H), 9.86 (br, 1H), 10.26 (br, 1H), and 11.64 (br, 1H).

10

Example 184

{2-[3-(2-Cyclopentanecarbonyl-4-methyl-phenyl)-ureido]-thiazol-4-ylmethyl}-carbamic acid 2-dimethylamino-ethyl ester hydrochloride salt



{2-[3-(2-Cyclopentanecarbonyl-4-methyl-phenyl)-ureido]-thiazol-4-ylmethyl}-carbamic acid 2-dimethylamino-ethyl ester hydrochloride salt (103 mg, 87%) was prepared from 2-[3-(2-cyclopentanecarbonyl-4-methyl-phenyl)-ureido]-thiazol-4-yl acetic acid (97 mg, 0.25 mmol)


15

and 2-N,N-dimethylethanol (0.05 mL, 0.5 mmol) following the general procedure M followed by treatment with HCl in ether.

LC-MS (m/z): 474 ($M+1$)⁺; ¹H NMR (400 MHz, CDCl₃): δ 1.42 (t, 2H), 1.74 (m, 4H),
5 1.92 (m, 4H), 2.28 (s, 6H), 2.39 (s, 3H), 2.96 (m, 2H), 3.24 (s, 2H), 3.74 (p, 1H), 6.86 (s, 1H),
7.34 (d, 1H), 7.72 (s, 1H), 8.44 (d, 1H), 9.12 (br, 1H), 10.34 (br, 1H), and 11.22 (br, 1H).

Example 185

N-{2-[3-(2-Cyclopentanecarbonyl-4-methyl-phenyl)-ureido]-thiazol-4-ylmethyl}-N',N'-
10 dimethylsulfamide




N-{2-[3-(2-Cyclopentanecarbonyl-4-methyl-phenyl)-ureido]-thiazol-4-ylmethyl}-N',N'-
dimethylsulfamide (104 mg, 89%) was prepared from 1-(4-aminomethyl-thiazol-2-yl)-3-(2-
cyclopentanecarbonyl-4-methyl-phenyl)-urea (90 mg, 0.25 mmol) and dimethylsulfamoyl
15 chloride (0.06 mL, 0.5 mmol) following the general procedure T.

LC-MS (m/z): 466 ($M+1$)⁺; ¹H NMR (400 MHz, CDCl₃): δ 1.72 (m, 4H), 1.88 (m, 4H),
2.36 (s, 3H), 2.75 (s, 6H), 3.72 (p, 1H), 4.22 (d, 2H), 6.74 (s, 1H), 7.03 (br, 1H), 7.34 (d, 1H),
7.70 (s, 1H), 8.42 (d, 1H), 9.72 (br, 1H), and 11.66 (br, 1H).

Example 186

3-[[2-[3-(2-Cyclopentanecarbonyl-4-methyl-phenyl)-ureido]-thiazol-4-ylmethyl]-(2-
methoxycarbonyl-ethyl)-amino]-propionic acid methyl ester




3-[(2-{3-(2-Cyclopentanecarbonyl-4-methyl-phenyl)-ureido}-thiazol-4-ylmethyl)-(2-methoxycarbonyl-ethyl)-amino]-propionic acid methyl ester (114 mg, 86%) was prepared by heating 1-(4-aminomethyl-thiazol-2-yl)-3-(2-cyclopentanecarbonyl-4-methyl-phenyl)-urea (90
 5 mg, 0.25 mmol) with methyl acrylate (0.1 mL, 1.1 mmol) and Cs_2CO_3 (325 mg, 1.0 mmol) in THF (5 mL) at 60 °C for 2h followed by column purification [silica, DCM:ethyl acetate (80:20 to 20:80)].

LC-MS (m/z): 531 ($M+1$)⁺; ^1H NMR (400 MHz, CDCl_3): δ 1.68 (m, 4H), 1.84 (m, 2H),
 2.04 (m, 2H), 2.16 (s, 6H), 2.36 (s, 3H), 2.45 (t, 4H), 3.66 (m, 4H), 3.72 (p, 1H), 4.48 (s, 2H),
 10 6.78 (s, 1H), 7.34 (d, 1H), 7.66 (s, 1H), 8.44 (d, 1H), 10.54 (br, 1H), and 11.62 (br, 1H).

Example 187

((2-{3-(2-Cyclopentanecarbonyl-4-methyl-phenyl)-ureido}-thiazol-4-ylmethyl)-methoxycarbonylmethyl-amino)-acetic acid methyl ester




((2-[3-(2-Cyclopentanecarbonyl-4-methyl-phenyl)-ureido]-thiazol-4-ylmethyl)-methoxycarbonylmethyl-amino)-acetic acid methyl ester (99 mg, 79%) was prepared by treating 1-(4-aminomethyl-thiazol-2-yl)-3-(2-cyclopentanecarbonyl-4-methyl-phenyl)-urea (90 mg, 0.25 mmol) with methyl bromoacetate (0.04 mL, 0.5 mmol) and pyridine (0.082 mL) in DCM (5 mL) at 60 °C for 2h followed by column purification [silica, DCM:ethyl acetate (80:20 to 20:80)].

LC-MS (*m/z*): 503 (*M*+1)⁺; ¹H NMR (400 MHz, CDCl₃): δ 1.53 (m, 4H), 1.76 (m, 4H), 2.18 (s, 3H), 2.59-2.74, (m, 6H), 3.50 (s, 6H), 3.62 (m, 1H), 6.69 (s, 1H), 7.18 (d, 1H), 7.54 (s, 1H), 8.18 (d, 1H), 10.58 (br, 1H), and 11.28 (br, 1H).

10

Example 188

(Carboxymethyl-{2-[3-(2-cyclopentanecarbonyl-4-methyl-phenyl)-ureido]-thiazol-4-ylmethyl}-amino)-acetic acid




15 (Carboxymethyl-{2-[3-(2-cyclopentanecarbonyl-4-methyl-phenyl)-ureido]-thiazol-4-ylmethyl}-amino)-acetic acid (44 mg, 88%) was prepared from ((2-[3-(2-cyclopentanecarbonyl-4-methyl-phenyl)-ureido]-thiazol-4-ylmethyl)-methoxycarbonylmethyl-amino)-acetic acid methyl ester (50 mg, 0.1 mmol) following the general procedure J.

LC-MS (*m/z*): 474 (*M*+1)⁺; ¹H NMR (400 MHz, CD₃OD): δ 1.53 (m, 4H), 1.76 (m, 4H), 2.32 (s, 3H), 2.63 (s, 2H), 2.74 (s, 4H), 3.72 (m, 1H), 6.76 (s, 1H), 7.28 (d, 1H), 7.66 (s, 1H), 8.38 (d, 1H), 9.66 (br, 1H), 10.08 (br, 2H), and 11.46 (br, 1H).

20

Example 189

25 N-{2-[3-(2-Cyclopentanecarbonyl-4-methyl-phenyl)-ureido]-thiazol-4-ylmethyl}-2-dimethylamino-acetamide




N-{2-[3-(2-Cyclopentanecarbonyl-4-methyl-phenyl)-ureido]-thiazol-4-ylmethyl}-2-dimethylamino-acetamide (86 mg, 78%) was prepared from 1-(4-aminomethyl-thiazol-2-yl)-3-(2-cyclopentanecarbonyl-4-methyl-phenyl)-urea (90 mg, 0.25 mmol) and N,N-dimethylglycine (0.30 mg, 0.30 mmol) following the general procedure K.

LC-MS (m/z): 444 ($M+1$)⁺; ¹H NMR (400 MHz, CDCl₃): δ 1.70 (m, 4H), 1.92 (m, 4H), 2.30 (s, 3H), 2.82 (s, 3H), 2.96 (s, 3H), 3.75 (s, 2H), 3.82 (m, 1H), 4.28 (d, 2H), 6.98 (s, 1H), 7.38 (d, 1H), 7.67 (s, 1H), 8.02 (d, 1H), 8.44 (br, 1H), 9.22 (br, 1H), and 11.44 (br, 1H).

10

Example 190

1-(2-Cyclopentanecarbonyl-4-methyl-phenyl)-3-(4-guanidinomethyl-thiazol-2-yl)-urea




To a solution of 1-(4-aminomethyl-thiazol-2-yl)-3-(2-cyclopentanecarbonyl-4-methyl-phenyl)-urea (90 mg, 0.25 mmol) in THF (10 mL) was added pyrrole-1-carboxamidine (54 mg, 0.50 mmol) and DIEA (90 μ L). The reaction mixture was heated 60 °C for 3 h and concentrated. The crude product was purified by flash chromatography [silica, DCM:ethyl acetate (50:50 to 10:90)] to yield 1-(2-cyclopentanecarbonyl-4-methyl-phenyl)-3-(4-guanidinomethyl-thiazol-2-yl)-urea (88 mg, 88%).

LC-MS (m/z): 401 ($M+1$)⁺; ¹H NMR (400 MHz, CDCl₃): δ 1.50 (m, 4H), 1.72 (m, 4H), 2.23 (s, 3H), 3.08 (q, 1H), 3.62 (p, 1H), 4.26 (br, 2H), 4.32 (t, 2H), 6.71 (s, 1H), 7.16 (d, 1H), 7.48 (s, 1H), 7.96 (br, 1H), 8.12 (d, 1H), 8.70 (br, 1H), and 10.92 (br, 1H).

Example 191

1-(2-Cyclopentanecarbonyl-4-methyl-phenyl)-3-[4-(4-S-methyl-2,5-dioxo-imidazolidin-1-ylmethyl)-thiazol-2-yl]-urea




1-(2-Cyclopentanecarbonyl-4-methyl-phenyl)-3-[4-(4-S-methyl-2,5-dioxo-imidazolidin-1-ylmethyl)-thiazol-2-yl]-urea (78 mg, 69%) was prepared from 1-(4-aminomethyl-thiazol-2-yl)-3-(2-cyclopentanecarbonyl-4-methyl-phenyl)-urea (358 mg, 1.0 mmol) and t-Boc-L-alanine (100 mg, 0.80 mmol) following the general procedure U.

LC-MS (m/z): 456 ($M+1$)⁺; ¹H NMR (400 MHz, CDCl₃): δ 1.43 (d, 3H), 1.64 (m, 4H), 1.88 (m, 4H), 2.35 (s, 3H), 3.56 (q, 1H), 3.74 (p, 1H), 4.73 (s, 2H), 5.52 (br, 1H), 6.84 (s, 1H), 7.32 (d, 1H), 7.68 (s, 1H), 8.40 (d, 1H), 9.84 (br, 1H), and 11.44 (br, 1H).

Example 192

1-(4-[[Bis-(3H-imidazol-4-ylmethyl)-amino]-methyl]-thiazol-2-yl)-3-(2-cyclopentanecarbonyl-4-methyl-phenyl)-urea




1-(4-[[Bis-(3H-imidazol-4-ylmethyl)-amino]-methyl]-thiazol-2-yl)-3-(2-cyclopentanecarbonyl-4-methyl-phenyl)-urea (110 mg, 85%) was prepared from 1-(4-aminomethyl-thiazol-2-yl)-3-(2-cyclopentanecarbonyl-4-methyl-phenyl)-urea (90 mg, 0.25 mmol) and imidazole-2-carboxyaldehyde (48 mg, 0.50 mmol) following the general procedure O.

LC-MS (m/z): 519 ($M+1$)⁺; ¹H NMR (400 MHz, CDCl₃): δ 1.58 (m, 4H), 1.84 (m, 4H), 2.34 (s, 3H), 3.10 (s, 2H), 3.51 (s, 4H), 3.64 (p, 1H), 6.62 (s, 1H), 6.84 (s, 1H), 7.08 (s, 1H), 7.11 (d, 1H), 7.66 (d, 1H), 7.70 (s, 1H), 7.83 (d, 1H), 8.36 (d, 1H), 8.70 (br, 2H), 9.74 (br, 1H), and 11.30 (br, 1H).

Example 193

1-(2-Cyclopentanecarbonyl-4-methyl-phenyl)-3-[4-(N',N',N'',N''-tetramethyl-guanidinomethyl)-thiazol-2-yl]-urea



15

1-(4-Aminomethyl-thiazol-2-yl)-3-(2-cyclopentanecarbonyl-4-methyl-phenyl)-urea (90 mg, 0.25 mmol) in DMF were added HBTU (94 mg, 0.25 mmol) and DIEA (90 μ L). The

reaction mixture was stirred at rt for 2 h. The reaction mixture was poured into water (30 mL) and extracted with ethyl acetate (3 x 25 mL). The organic layer was washed (water, brine), dried (Na_2SO_4) and concentrated in vacuo. The crude product was purified by flash chromatography [silica, DCM:ethyl acetate (50:50 to 10:90)] to yield 1-(2-


5 cyclopentanecarbonyl-4-methyl-phenyl)-3-[4-(N',N',N'',N''-tetramethyl-guanidinomethyl)-thiazol-2-yl]-urea (92 mg, 81%).

LC-MS (m/z): 457 ($M+1$)⁺; ¹H NMR (400 MHz, CDCl_3): δ 1.66 (m, 4H), 1.82 (m, 2H), 1.94 (m, 2H), 2.37 (s, 3H), 2.96 (s, 12H), 3.76 (p, 1H), 4.32 (s, 2H), 6.80 (s, 1H), 7.38 (d, 1H), 7.70 (s, 1H), 8.38 (d, 1H), 10.54 (br, 1H), and 11.58 (br, 1H).

10

Example 194

2-Amino-ethanesulfonic acid {2-[3-(2-cyclopentanecarbonyl-4-methyl-phenyl)-ureido]-thiazol-4-ylmethyl}-amide




15 2-Amino-ethanesulfonic acid {2-[3-(2-cyclopentanecarbonyl-4-methyl-phenyl)-ureido]-thiazol-4-ylmethyl}-amide (104 mg, 83%) was prepared from 1-(4-aminomethyl-thiazol-2-yl)-3-(2-cyclopentanecarbonyl-4-methyl-phenyl)-urea (90 mg, 0.25 mmol) and 2-(1,3-Dioxo-1,3-dihydro-isoindol-2-yl)-ethanesulfonyl chloride (70 mg, 0.25 mmol) following the general procedure T. This intermediate was deprotected by heating with excess of hydrazine (0.2

20 mL, 1.0 M. in THF).

LC-MS (m/z): 466 ($M+1$)⁺; ¹H NMR (400 MHz, CD_3OD): δ 1.68 (m, 4H), 1.92 (m, 4H), 2.38 (s, 3H), 3.22 (t, 2H), 3.68 (m, 2H), 3.76 (p, 1H), 4.08 (s, 2H), 4.84 (br, 1H), 5.38 (br, 2H), 6.76 (s, 1H), 7.36 (d, 1H), 7.74 (s, 1H), 8.48 (d, 1H), 10.22 (br, 1H), and 11.36 (br, 1H).

25 EXAMPLE 195

N-{2-[3-(2-Cyclopentanecarbonyl-4-methyl-phenyl)-ureido]-thiazol-4-ylmethyl}-2-methanesulfonylamino-acetamide




N-(2-{3-(2-Cyclopentanecarbonyl-4-methyl-phenyl)-ureido}-thiazol-4-ylmethyl)-2-methanesulfonylamino-acetamide (103 mg, 84%) was prepared from 2-amino-N-(2-{3-(2-cyclopentanecarbonyl-4-methyl-phenyl)-ureido}-thiazol-4-ylmethyl)-acetamide (104 mg, 0.25 mmol) and methanesulfonyl chloride (0.04 mL, 0.5 mmol) following the general procedure L.

LC-MS (m/z): 494 ($M+1$)⁺; ¹H NMR (400 MHz, CDCl₃): δ 1.70 (m, 4H), 1.88 (m, 4H), 2.36 (s, 3H), 3.28 (t, 2H), 3.38 (s, 3H), 3.74 (p, 1H), 4.16 (d, 2H), 6.84 (s, 1H), 6.96 (br, 1H), 7.38 (d, 1H), 7.44 (br, 1H), 7.69 (s, 1H), 8.36 (d, 1H), 10.58 (br, 1H), and 11.28 (br, 1H).

EXAMPLE 196

{[[[2-{3-(2-Cyclopentanecarbonyl-4-methyl-phenyl)-ureido}-thiazol-4-ylmethyl]-carbamoyl]-methyl]-amino}-acetic acid




{[[[2-{3-(2-Cyclopentanecarbonyl-4-methyl-phenyl)-ureido}-thiazol-4-ylmethyl]-carbamoyl]-methyl]-amino}-acetic acid (44 mg, 88%) was prepared from ({2-[3-(2-cyclopentanecarbonyl-4-methyl-phenyl)-ureido]-thiazol-4-ylmethyl}-methoxycarbonylmethyl-amino)-acetic acid methyl ester (50 mg, 0.1 mmol) following the general procedure J.

LC-MS (*m/z*): 474 (*M*+1)⁺; ¹H NMR (400 MHz, CD₃OD): δ 1.58 (m, 4H), 1.84 (m, 4H), 2.31 (s, 3H), 2.79 (d, 2H), 3.58 (d, 2H) 3.70 (m, 1H), 4.18 (s, 2H), 4.96 (br, 1H), 6.82 (s, 1H), 7.03 (br, 1H), 7.30 (d, 1H), 7.72 (s, 1H), 8.44 (d, 1H), 9.66 (br, 1H), 10.08 (br, 1H), and 11.46 (br, 1H).

5

Example 197

{[(2-[3-(2-Cyclopentanecarbonyl-4-methyl-phenyl)-ureido]-thiazol-4-ylmethyl)-carbamoyl]-methyl]-amino}-acetic acid methyl ester




10 {[(2-[3-(2-Cyclopentanecarbonyl-4-methyl-phenyl)-ureido]-thiazol-4-ylmethyl)-carbamoyl]-methyl]-amino}-acetic acid methyl ester (44 mg, 88%) was prepared by treating 2-amino-N-{2-[3-(2-cyclopentanecarbonyl-4-methyl-phenyl)-ureido]-thiazol-4-ylmethyl}-acetamide (104 mg, 0.25 mmol) with methyl bromoacetate (0.023 mL, 0.25 mmol) in THF (5 mL) at 60 °C for 2h followed by column purification [silica, DCM:ethyl acetate (80:20 to
15 20:80)].

LC-MS (*m/z*): 488 (*M*+1)⁺; ¹H NMR (400 MHz, CDCl₃): δ 1.64 (m, 4H), 1.82 (m, 4H), 2.38 (s, 3H), 2.66 (s, 2H), 2.78, (s, 2H), 3.18 (s, 2H), 3.52 (s, 3H), 3.72 (m, 1H), 6.76 (s, 1H), 7.06 (br, 1H), 7.28 (d, 1H), 7.66 (s, 1H), 8.38 (d, 1H), 8.86 (br, 1H), 9.74 (br, 1H), and 11.46 (br, 1H).

20

Example 198

3-({2-[3-(2-Cyclopentanecarbonyl-4-methyl-phenyl)-ureido]-thiazol-4-ylmethyl}-sulfamoyl)-propionic acid




3-({2-[3-(2-Cyclopentanecarbonyl-4-methyl-phenyl)-ureido]-thiazol-4-ylmethyl}-sulfamoyl)-propionic acid (85 mg, 87%) was prepared from 3-({2-[3-(2-cyclopentanecarbonyl-4-methyl-phenyl)-ureido]-thiazol-4-ylmethyl}-sulfamoyl)-propionic acid methyl ester (101 mg, 0.20 mmol) following the general procedure J.

LC-MS (m/z): 495 ($M+1$)⁺; ¹H NMR (400 MHz, CD₃OD): δ 1.74 (m, 4H), 1.88 (m, 4H), 2.33 (s, 3H), 3.12 (t, 2H), 3.48 (t, 2H), 3.72 (p, 1H), 3.88 (d, 2H), 6.72 (s, 1H), 7.36 (d, 1H), 7.68 (s, 1H), 8.42 (d, 1H), 9.54 (br, 1H), 10.88 (br, 1H), 11.12 (br, 1H), and 11.54 (br, 1H).

Example 199

N-{2-[3-(2-Cyclopentanecarbonyl-4-methyl-phenyl)-ureido]-thiazol-4-ylmethyl}-2-methanesulfonyl-acetamide




N-{2-[3-(2-Cyclopentanecarbonyl-4-methyl-phenyl)-ureido]-thiazol-4-ylmethyl}-2-methanesulfonyl-acetamide (103 mg, 84%) was prepared from 1-(4-aminomethyl-thiazol-2-yl)-3-(2-cyclopentanecarbonyl-4-methyl-phenyl)-urea (90 mg, 0.25 mmol) and methanesulfonyl acetic acid (55 mg, 0.4 mmol) following the general procedure T.

LC-MS (m/z): 479 ($M+1$)⁺; ¹H NMR (400 MHz, CD₃OD): δ 1.68 (m, 4H), 1.84 (m, 4H), 2.35 (s, 3H), 2.78 (s, 2H), 3.12 (s, 3H), 3.86 (m, 1H), 4.08 (d, 2H), 6.89 (s, 1H), 7.04 (br, 1H), 7.37 (d, 1H), 7.80 (s, 1H), 8.24 (d, 1H), 8.52 (br, 1H), and 11.36 (br, 1H).

5

Example 200

1-{2-[3-(2-Cyclopentanecarbonyl-4-methyl-phenyl)-ureido]-thiazol-4-ylmethyl}-piperidine-S-3-carboxylic acid ethyl ester




10 1-{2-[3-(2-Cyclopentanecarbonyl-4-methyl-phenyl)-ureido]-thiazol-4-ylmethyl}-piperidine-S-3-carboxylic acid ethyl ester (107 mg, 86%) was prepared from 2-[3-(2-(1-(2-cyclopentanecarbonyl-4-methyl-phenyl)-3-(4-formyl-thiazol-2-yl)-urea (89 mg, 0.25 mmol) and piperidine-S-3-carboxylic acid ethyl ester (40 mg, 0.25 mmol) following the general procedure O.

15 LC-MS (m/z): 499 ($M+1$)⁺; ¹H NMR (400 MHz, CDCl₃): δ 1.22 (t, 3H), 1.63-1.70 (m, 6H), 1.89 (m, 4H), 2.22 (m, 1H), 2.35 (s, 3H), 2.60 (m, 2H), 2.84 (m, 2H), 3.00 (d, 2H), 3.55 (m, 2H), 3.72 (p, 1H), 4.10 (q, 2H), 6.67 (s, 1H), 7.33 (d, 1H), 7.68 (s, 1H), 8.40 (br, 1H), 10.50 (br, 1H), and 11.50 (br, 1H).

Example 201

20 1-{2-[3-(2-Cyclopentanecarbonyl-4-methyl-phenyl)-ureido]-thiazol-4-ylmethyl}-piperidine-S-3-carboxylic acid




1-{2-[3-(2-Cyclopentanecarbonyl-4-methyl-phenyl)-ureido]-thiazol-4-ylmethyl}-piperidine-S-3-carboxylic acid (104 mg, 81%) was prepared from 1-{2-[3-(2-cyclopentanecarbonyl-4-methyl-phenyl)-ureido]-thiazol-4-ylmethyl}-piperidine-S-3-carboxylic acid ethyl ester (135 mg, 0.25 mmol) following the general procedure J.

LC-MS (m/z): 471 ($M+1$)⁺; ¹H NMR (400 MHz, CDCl₃): δ 1.58-1.64 (m, 4H), 1.84-1.96 (m, 6H), 2.16 (m, 1H), 2.28 (s, 3H), 2.80 (m, 1H), 3.05 (m, 1H), 3.34-3.49 (m, 2H), 3.66 (d, 1H), 3.93 (m, 1H), 4.34-4.50 (m, 2H), 4.70 (m, 1H), 6.83 (s, 1H), 7.36 (d, 1H), 7.60 (s, 1H), 8.25 (d, 1H), 9.00 (br, 1H), 10.20 (br, 1H), and 11.36 (br, 1H).

10

Example 202

((2-[3-(2-Cyclopentanecarbonyl-4-methyl-phenyl)-ureido]-thiazol-4-ylmethyl)-amino)-(tetrahydro-pyran-4-yl)-acetic acid methyl ester



15


((2-[3-(2-Cyclopentanecarbonyl-4-methyl-phenyl)-ureido]-thiazol-4-ylmethyl)-amino)-(tetrahydro-pyran-4-yl)-acetic acid methyl ester (113 mg, 88%) was prepared from 2-[3-(2-(2-cyclopentanecarbonyl-4-methyl-phenyl)-3-(4-formyl-thiazol-2-yl)-urea (89 mg, 0.25 mmol)

and amino-(tetrahydro-pyran-4-yl)-acetic acid methyl ester (44 mg, 0.25 mmol) following the general procedure O.

LC-MS (*m/z*): 515 (*M*+1)⁺; ¹H NMR (400 MHz, CDCl₃): δ 1.42 (m, 4H), 1.66-1.73 (m, 4H), 1.80-1.93 (m, 4H), 2.35 (s, 3H), 3.14 (d, 2H), 3.32 (t, 2H), 3.64 (t, 2H), 3.72 (p, 1H), 3.78 (d, 2H), 3.92 (s, 3H), 6.24 (br, 1H), 6.64 (s, 1H), 7.32 (d, 1H), 7.68 (s, 1H), 8.42 (br, 1H), 9.78 (br, 1H), and 11.50 (br, 1H).

Example 203

((2-[3-(2-Cyclopentanecarbonyl-4-methyl-phenyl)-ureido]-thiazol-4-ylmethyl)-amino)-(tetrahydro-pyran-4-yl)-acetic acid




((2-[3-(2-Cyclopentanecarbonyl-4-methyl-phenyl)-ureido]-thiazol-4-ylmethyl)-amino)-(tetrahydro-pyran-4-yl)-acetic acid (87 mg, 87%) was prepared from ((2-[3-(2-cyclopentanecarbonyl-4-methyl-phenyl)-ureido]-thiazol-4-ylmethyl)-amino)-(tetrahydro-pyran-4-yl)-acetic acid methyl ester (128 mg, 0.25 mmol) following the general procedure J.

LC-MS (*m/z*): 501 (*M*+1)⁺; ¹H NMR (400 MHz, CDCl₃): δ 1.48 (m, 4H), 1.68-1.75 (m, 4H), 1.82-1.92 (m, 4H), 2.36 (s, 3H), 3.36 (d, 2H), 3.44 (t, 2H), 3.68 (t, 2H), 3.74 (p, 1H), 3.82 (d, 2H), 5.24 (br, 1H), 6.76 (s, 1H), 7.34 (d, 1H), 7.71 (s, 1H), 8.44 (d, 1H), 9.62 (br, 1H), 10.38 (br, 1H), and 11.20 (br, 1H).

Example 204

1-(2-Cyclopentanecarbonyl-4-methyl-phenyl)-3-{4-[(2-morpholin-4-yl-ethylamino)-methyl]-thiazol-2-yl}-urea




1-(2-Cyclopentanecarbonyl-4-methyl-phenyl)-3-{4-[(2-morpholin-4-yl-ethylamino)-methyl]-thiazol-2-yl}-urea (97 mg, 82%) was prepared from 2-[3-(2-1-(2-cyclopentanecarbonyl-4-methyl-phenyl)-3-(4-formyl-thiazol-2-yl)-urea (89 mg, 0.25 mmol) and 2-morpholin-4-yl-ethylamine (35 mg, 0.25 mmol) following the general procedure O.

LC-MS (m/z): 472 ($M+1$)⁺; ¹H NMR (400 MHz, CDCl₃): δ 1.66 (m, 4H), 1.88 (m, 4H), 2.33 (s, 3H), 2.38 (m, 2H), 2.50 (t, 2H), 2.80 (t, 2H), 3.59-3.64 (m, 6H), 3.70 p, 1H), 3.85 (m, 2H), 5.88 (br, 1H), 6.58 (s, 1H), 7.32 (d, 1H), 7.66 (s, 1H), 8.42 (d, 1H), 10.08 (br, 1H), and 11.28 (br, 1H).

Example 205

1-(2-Cyclopentanecarbonyl-4-methyl-phenyl)-3-(4-morpholin-4-ylmethyl-thiazol-2-yl)-urea




1-(2-Cyclopentanecarbonyl-4-methyl-phenyl)-3-(4-morpholin-4-ylmethyl-thiazol-2-yl)-urea (83 mg, 78%) was prepared from 2-[3-(2-1-(2-cyclopentanecarbonyl-4-methyl-phenyl)-

3-(4-formyl-thiazol-2-yl)-urea (89 mg, 0.25 mmol) and (30 mg, 0.3 mmol) and morpholine (0.25 mL, 0.3 mmol) following the general procedure O.

LC-MS (*m/z*): 429 (*M*+1)⁺; ¹H NMR (400 MHz, CDCl₃): δ 1.62-1.80 (m, 4H), 1.80-1.94 (m, 4H), 2.35 (s, 3H), 2.50 (m, 4H), 3.53 (s, 2H), 3.70 (t, 4H), 3.75 (m, 1H), 6.70 (s, 1H), 7.34 (d, 1H), 7.69 (s, 1H), 8.40 (d, 1H), 9.75 (br, 1H), and 11.58 (br, 1H).

Example 206

1-(2-Cyclopentanecarbonyl-4-methyl-phenyl)-3-[4-(3-oxo-piperazin-1-ylmethyl)-thiazol-2-yl]-urea




1-(2-Cyclopentanecarbonyl-4-methyl-phenyl)-3-[4-(3-oxo-piperazin-1-ylmethyl)-thiazol-2-yl]-urea (79 mg, 82%) was prepared from 2-[3-(2-1-(2-cyclopentanecarbonyl-4-methyl-phenyl)-3-(4-formyl-thiazol-2-yl)-urea (89 mg, 0.25 mmol) and piperazine-2-one (30 mg, 0.30 mmol) following the general procedure O.

LC-MS (*m/z*): 442 (*M*+1)⁺; ¹H NMR (400 MHz, CDCl₃): δ 1.60-1.74 (m, 4H), 1.80-1.94 (m, 4H), 2.35 (s, 3H), 2.67 (t, 2H), 3.30 (s, 2H), 3.38 (m, 2H), 3.58 (s, 2H), 3.73 (p, 1H), 6.66 (s, 1H), 7.32 (d, 1H), 7.67 (s, 1H), 7.78 (br, 1H), 8.38 (d, 1H), 10.44 (br, 1H), and 11.40 (br, 1H).

Example 207

(4-{2-[3-(2-Cyclopentanecarbonyl-4-methyl-phenyl)-ureido]-thiazol-4-ylmethyl}-piperazin-1-yl)-acetic acid ethyl ester




(4-{2-[3-(2-Cyclopentanecarbonyl-4-methyl-phenyl)-ureido]-thiazol-4-ylmethyl}-piperazin-1-yl)-acetic acid ethyl ester (79 mg, 82%) was prepared from 2-[3-(2-(2-cyclopentanecarbonyl-4-methyl-phenyl)-3-(4-formyl-thiazol-2-yl)-urea (89 mg, 0.25 mmol) and piperazin-1-yl)-acetic acid ethyl ester (43 mg, 0.25 mmol) following the general procedure O.

LC-MS (m/z): 514 ($M+1$)⁺; ¹H NMR (400 MHz, CDCl₃): δ 1.28 (t, 3H), 1.46-1.62 (m, 4H), 1.66-1.86 (m, 4H), 2.35 (s, 3H), 3.28 (m, 2H), 3.42 (s, 2H), 3.40-3.64 (m, 4H), 3.83 (m, 1H), 3.92 (s, 2H), 4.12 (m, 2H), 4.48 (s, 2H), 6.84 (s, 1H), 7.29 (d, 1H), 7.67 (s, 1H), 8.28 (d, 1H), 11.14 (br, 1H), and 11.62 (br, 1H).

Example 208

(4-{2-[3-(2-Cyclopentanecarbonyl-4-methyl-phenyl)-ureido]-thiazol-4-ylmethyl}-piperazin-1-yl)-acetic acid




(4-{2-[3-(2-Cyclopentanecarbonyl-4-methyl-phenyl)-ureido]-thiazol-4-ylmethyl}-piperazin-1-yl)-acetic acid (40 mg, 83%) was prepared from (4-{2-[3-(2-cyclopentanecarbonyl-4-methyl-phenyl)-ureido]-thiazol-4-ylmethyl}-piperazin-1-yl)-acetic acid ethyl ester (52 mg, 0.10 mmol) following the general procedure J.

LC-MS (m/z): 486 ($M+1$)⁺; ¹H NMR (400 MHz, CDCl₃): δ 1.44–1.58 (m, 4H), 1.62–1.80 (m, 4H), 2.22 (s, 3H), 2.98 (s, 2H), 3.18 (m, 2H), 3.62 (p, 1H), 4.02–4.18 (m, 8H), 7.00 (s, 1H), 7.22 (d, 1H), 7.58 (s, 1H), 8.18 (d, 1H), 9.54 (br, 1H), 10.78 (br, 1H), and 11.44 (br, 1H).

5

Example 209

3-(4-{2-[3-(2-Cyclopentanecarbonyl-4-methyl-phenyl)-ureido]-thiazol-4-ylmethyl}-piperazin-1-yl)-propionic acid ethyl ester




3-(4-{2-[3-(2-Cyclopentanecarbonyl-4-methyl-phenyl)-ureido]-thiazol-4-ylmethyl}-piperazin-1-yl)-propionic acid ethyl ester (118 mg, 89%) was prepared from 2-[3-(2-(2-cyclopentanecarbonyl-4-methyl-phenyl)-3-(4-formyl-thiazol-2-yl)-urea (89 mg, 0.25 mmol) and 3-piperazin-1-yl-propionic acid ethyl ester (61 mg, 0.30 mmol) following the general procedure O.

LC-MS (m/z): 528 ($M+1$)⁺; ¹H NMR (400 MHz, CDCl₃): δ 1.24 (t, 3H), 1.44–1.64 (m, 4H), 1.64–1.80 (m, 4H), 2.36 (s, 3H), 3.22–3.38 (m, 6H), 3.40–3.64 (m, 4H), 3.83 (m, 1H), 3.96 (m, 2H), 4.08 (m, 2H), 4.58 (s, 2H), 6.97 (s, 1H), 7.14 (d, 1H), 7.64 (s, 1H), 8.16 (d, 1H), 11.04 (br, 1H), and 11.22 (br, 1H).

Example 210

3-(4-{2-[3-(2-Cyclopentanecarbonyl-4-methyl-phenyl)-ureido]-thiazol-4-ylmethyl}-piperazin-1-yl)-propionic acid




3-(4-{2-[3-(2-Cyclopentanecarbonyl-4-methyl-phenyl)-ureido]-thiazol-4-ylmethyl}-piperazin-1-yl)-propionic acid (38 mg, 76%) was prepared from 3-(4-{2-[3-(2-cyclopentanecarbonyl-4-methyl-phenyl)-ureido]-thiazol-4-ylmethyl}-piperazin-1-yl)-propionic acid ethyl ester (53 mg, 0.10 mmol) following the general procedure J.

LC-MS (m/z): 500 ($M+1$)⁺; ¹H NMR (400 MHz, CDCl₃): δ 1.42-1.60 (m, 4H), 1.64-1.78 (m, 4H), 2.26 (s, 3H), 3.38-3.44 (m, 6H), 3.48-3.66 (m, 4H), 3.77 (m, 1H), 4.02 (m, 2H), 4.64 (s, 2H), 6.88 (s, 1H), 7.30 (d, 1H), 7.73 (s, 1H), 8.28 (d, 1H), 9.88 (br, 1H), 10.38 (br, 1H), and 11.34 (br, 1H).

10

Example 211

(3-{2-[3-(2-Cyclopentanecarbonyl-4-methyl-phenyl)-ureido]-thiazol-4-ylmethyl}-ureido)-acetic acid




(3-{2-[3-(2-Cyclopentanecarbonyl-4-methyl-phenyl)-ureido]-thiazol-4-ylmethyl}-ureido)-acetic acid methyl ester (106 mg, 90%) was prepared from 2-[3-(2-cyclopentanecarbonyl-4-methyl-phenyl)-ureido]-thiazol-4-yl acetic acid (97 mg, 0.25 mmol) and glycine methyl ester (48 mg, 0.5 mmol) following the general procedure M.

5 LC-MS (*m/z*): 460 (*M*+1)⁺; ¹H NMR (400 MHz, CDCl₃): δ 1.73 (m, 4H), 1.88 (m, 4H), 2.36 (s, 3H), 3.46 (s, 2H), 3.66 (s, 2H), 3.78 (p, 1H), 6.74 (s, 1H), 7.36 (d, 1H), 7.71 (s, 1H), 8.28 (d, 1H), 8.44 (br, 1H), 8.92 (br, 1H), 9.96 (br, 1H), and 11.68 (br, 1H).

Example 212

10 1-(2-Cyclopentanecarbonyl-4-methyl-phenyl)-3-[4-(2,5-dioxo-imidazolidin-1-ylmethyl)-thiazol-2-yl]-urea




1-(2-Cyclopentanecarbonyl-4-methyl-phenyl)-3-[4-(2,5-dioxo-imidazolidin-1-ylmethyl)-thiazol-2-yl]-urea (78 mg, 69%) was prepared from 1-(4-aminomethyl-thiazol-2-yl)-3-(2-cyclopentanecarbonyl-4-methyl-phenyl)-urea (358 mg, 1.0 mmol) and t-Boc-glycine (90 mg, 0.80 mmol) following the general procedure U.

15 LC-MS (*m/z*): 442 (*M*+1)⁺; ¹H NMR (400 MHz, CDCl₃): δ 1.66 (m, 4H), 1.90 (m, 4H), 2.38 (s, 3H), 3.43 (s, 2H), 3.76 (p, 1H), 4.24 (s, 2H), 5.48 (br, 1H), 6.78 (s, 1H), 7.34 (d, 1H), 7.72 (s, 1H), 8.44 (d, 1H), 9.66 (br, 1H), and 11.64 (br, 1H).

20

Example 213

4-({2-[3-(2-Cyclopentanecarbonyl-4-methyl-phenyl)-ureido]-thiazol-4-ylmethyl}-sulfamoyl)-benzoic acid




4-((2-[3-(2-Cyclopentanecarbonyl-4-methyl-phenyl)-ureido]-thiazol-4-ylmethyl)-sulfamoyl)-benzoic acid (78 mg, 58%) was prepared from 1-(4-aminomethyl-thiazol-2-yl)-3-(2-cyclopentanecarbonyl-4-methyl-phenyl)-urea (90 mg, 0.25 mmol) following the general procedure T.

LC-MS (m/z): 543 ($M+1$)⁺; ¹H NMR (400 MHz, CD₃OD): δ 1.73 (m, 4H), 1.88 (m, 4H), 2.33 (s, 3H), 3.68 (m, 2H), 3.74 (p, 1H), 5.88 (br, 1H), 6.82 (s, 1H), 7.36 (d, 1H), 7.48 (d, 2H), 7.68 (d, 2H), 7.74 (s, 1H), 8.48 (d, 1H), 9.76 (br, 1H), 10.38 (br, 1H), and 11.44 (br, 1H).

Example 214

3-((2-[3-(2-Cyclopentanecarbonyl-4-methyl-phenyl)-ureido]-thiazol-4-ylmethyl)-sulfamoyl)-propionic acid methyl ester




3-((2-[3-(2-Cyclopentanecarbonyl-4-methyl-phenyl)-ureido]-thiazol-4-ylmethyl)-sulfamoyl)-propionic acid methyl ester (116 mg, 91%) was prepared from 1-(4-aminomethyl-thiazol-2-yl)-3-(2-cyclopentanecarbonyl-4-methyl-phenyl)-urea (90 mg, 0.25 mmol) and 3-

chlorosulfonyl-propionic acid methyl ester (47 mg, 0.25 mmol) following the general procedure T.

LC-MS (m/z): 509 ($M+1$)⁺; ¹H NMR (400 MHz, CD₃OD): δ 1.73 (m, 4H), 1.88 (m, 4H), 2.33 (s, 3H), 3.04 (t, 2H), 3.42 (t, 2H), 3.74 (p, 1H), 4.06 (d, 2H), 4.26 (s, 3H), 6.58 (s, 1H), 7.32 (d, 1H), 7.72 (s, 1H), 8.48 (d, 1H), 9.76 (br, 1H), 10.38 (br, 1H), and 11.52 (br, 1H).

Example 215

{2-[3-(2-Cyclopentanecarbonyl-5-methyl-phenyl)-ureido]-thiazol-4-yl}-acetic acid ethyl ester




{2-[3-(2-Cyclopentanecarbonyl-5-methyl-phenyl)-ureido]-thiazol-4-yl}-acetic acid ethyl ester (154 mg, 80 %) was prepared from 2-amino-4 methyl-phenyl-cyclopentyl-methanone (102 mg, 0.5 mmol) and ethyl-2-amino-4-thiazolyl acetate (112 mg, 0.5 mmol) following the general procedure D.

LC-MS (m/z): 416 ($M+1$)⁺; ¹H NMR (400 MHz, CDCl₃): δ 1.26 (t, 3H), 1.69 (m, 4H), 1.88 (m, 4H), 2.41 (s, 3H), 3.70 (s, 2H), 3.72 (m, 1H), 4.18 (m, 2H), 6.71 (s, 1H), 6.88 (d, 1H), 7.7.81 (d, 1H), 8.44 (s, 1H), 9.48 (br, 1H), 11.82 (br, 1H).

Example 216

{2-[3-(2-Cyclopentanecarbonyl-5-methyl-phenyl)-ureido]-thiazol-4-yl}-acetic acid




{2-[3-(2-Cyclopentanecarbonyl-5-methyl-phenyl)-ureido]-thiazol-4-yl}-acetic acid (175 mg, 90%) was prepared from {2-[3-(2-cyclopentanecarbonyl-5-methyl-phenyl)-ureido]-thiazol-4-yl}-acetic acid ethyl ester (208 mg, 0.5 mmol) following the general procedure J.

LC-MS (m/z): 388 ($M+1$)⁺; ¹H NMR (400 MHz, DMSO- d_6): δ 1.74 (m, 4H), 1.87 (m, 4H), 2.35 (s, 3H), 3.56 (s, 2H), 3.84 (m, 1H), 6.85 (s, 1H), 6.96 (d, 1H), 7.95 (d, 1H), 8.17 (s, 1H), 10.96 (br, 1H), 12.22 (br, 1H), 12.40 (br, 1H).

10

Example 217

{2-[3-(2-Cyclopentanecarbonyl-5-fluoro-phenyl)-ureido]-thiazol-4-yl}-acetic acid ethyl ester




{2-[3-(2-Cyclopentanecarbonyl-4-fluorol-phenyl)-ureido]-thiazol-4-yl}-acetic acid ethyl ester (168 mg, 80%) was prepared from 2-amino-4-fluoro methyl-phenyl-cyclopentyl-methanone (103 mg, 0.5 mmol) and ethyl-2-amino-4-thiazolyl acetate (112 mg, 0.6 mmol) following the general procedure D.

LC-MS (m/z): 420 ($M+1$)⁺; ¹H NMR (400 MHz, CDCl₃): δ 1.26 (t, 3H), 1.69 (m, 4H), 1.88 (m, 4H), 3.67 (m, 1H), 3.72 (s, 2H), 4.18 (m, 2H), 6.68 (d, 1H), 6.75 (m, 1H), 7.93 (t, 1H), 8.44 (d, 1H), 10.20 (br, 1H), 11.90 (br, 1H).

20

Example 218

{2-[3-(2-Cyclopentanecarbonyl-5-fluoro-phenyl)-ureido]-thiazol-4-yl}-acetic acid




- 5 {2-[3-(2-Cyclopentanecarbonyl-5-methyl-phenyl)-ureido]-thiazol-4-yl}-acetic acid
(180 mg, 92%) was prepared from {2-[3-(2-cyclopentanecarbonyl-5-methyl-phenyl)-
ureido]-thiazol-4-yl}-acetic acid ethyl ester (209 mg, 0.5 mmol) following the general
procedure J.

LC-MS (*m/z*): 392 (*M*+1)⁺; ¹H NMR (400 MHz, DMSO-*d*₆): δ 1.61 (m, 4H), 1.74 (m,
10 2H), 1.87 (m, 2H), 3.57 (s 2H), 3.84 (m, 1H), 6.88 (s, 1H), 6.98 (t, 1 H), 8.21 (m, 2 H), 11.23
(br, 1H), 12.20 (br, 1H).

Example 219

- 15 {2-[3-(4-Bromo-2-cyclopentanecarbonyl-phenyl)-ureido]-thiazol-4-yl}-acetic acid
ethyl ester




- 20 {2-[3-(4-Bromo-2-cyclopentanecarbonyl-phenyl)-ureido]-thiazol-4-yl}-acetic acid
ethyl ester (192 mg, 80 %) was prepared from 2-amino-5-bromo-phenyl-cyclopentyl-
methanone (134 mg, 0.5 mmol) and ethyl-2-amino-4-thiazolyl acetate (112 mg, 0.6 mmol)
following the general procedure D.

LC-MS (m/z): 481 ($M+1$)⁺; ¹H NMR (400 MHz, CDCl₃): δ 1.27 (t, 3H), 1.70 (m, 4H), 1.87 (m, 4H), 3.67 (m, 1H), 3.72 (s, 2H), 4.19 (m, 2H), 6.70 (s, 1H), 7.59 (dd, 1H), 7.99 (s, 1H), 8.53 (br, 1H), 9.59 (br, 1H), 11.56 (br, 1H).

5

Example 220

{2-[3-(4-Bromo-2-cyclopentanecarbonyl-phenyl)-ureido]-thiazol-4-yl}-acetic acid



{2-[3-(4-Bromo-2-cyclopentanecarbonyl-phenyl)-ureido]-thiazol-4-yl}-acetic acid (204 mg, 90 %) was prepared from {2-[3-(4-bromo-2-cyclopentanecarbonyl-phenyl)-ureido]-thiazol-4-yl}-acetic acid ethyl ester (240 mg, 0.5 mmol) following the general procedure J.


10

LC-MS (m/z): 453 ($M+1$)⁺; ¹H NMR (400 MHz, DMSO- d_6): δ 1.72 (m, 4H), 1.82 (m, 4H), 3.54 (s, 2H), 3.78 (m, 1H), 6.86 (s, 1H), 7.72 (br, 1H), 8.10 (br, 1H), 8.22 (s, 1H), 10.65 (br, 1H), 11.70 (br, 1H), 12.40 (br, 1H).

15

Example 221

{4-[3-(5-Chloro-thiazol-2-yl)-ureido]-3-cyclopentanecarbonyl-phenyl}-acetic acid ethyl ester




{4-[3-(5-Chloro-thiazol-2-yl)-ureido]-3-cyclopentanecarbonyl-phenyl}-acetic acid ethyl ester (157 mg, 72%) was prepared from ethyl (4-methylcarboxy)-2-cyclopentanoylaniline (138 mg, 0.5 mmol) and 5-chloro-thiazol-2-ylamine (81mg, 0.6 mmol) following the general procedure D.

20

LC-MS (*m/z*): 436 (*M*+1)⁺; ¹H NMR (400 MHz, DMSO-*d*₆): δ 0.90 (t, 3H), 1.62 (m, 2H), 1.46 (m, 2H), 1.62 (m, 4H), 3.73 (s, 2 H), 3.85 (m, 1H), 4.15 (q, 2H), 7.42 (s, 1H), 7.48 (d, 1H), 7.98 (s, 1H), 8.20 (d, 1 H), 10.79 (br, 1 H), 12.10 (br 1H).

5 **Example 222**

{4-[3-(5-Chloro-thiazol-2-yl)-ureido]-3-cyclopentanecarbonyl-phenyl}-acetic acid




{4-[3-(5-Chloro-thiazol-2-yl)-ureido]-3-cyclopentanecarbonyl-phenyl}-acetic acid (180 mg, 88 %) was prepared from {4-[3-(5-chloro-thiazol-2-yl)-ureido]-3-cyclopentanecarbonyl-phenyl}-acetic acid ethyl ester (218 mg, 0.5 mmol) following the general procedure J.

LC-MS (*m/z*): 408 (*M*+1)⁺; ¹H NMR (400 MHz, DMSO-*d*₆): δ 1.73 (m, 4H), 1.89 (m, 4H), 3.35 (s, 2H), 3.85 (m, 1H), 7.42 (s, 1H), 7.47 (d, 1H), 7.96 (s, 1H), 8.20 (d, 1 H), 10.79 (s, 1 H), 12.20 (br, 1H),

15 **Example 223**

2-{4-[3-(5-Chloro-thiazol-2-yl)-ureido]-3-cyclopentanecarbonyl-phenyl}-N-(2-methanesulfonyl-ethyl)-acetamide




2-{4-[3-(5-Chloro-thiazol-2-yl)-ureido]-3-cyclopentanecarbonyl-phenyl}-N-(2-methanesulfonyl-ethyl)-acetamide (180 mg, 70 %) was prepared from {4-[3-(5-chloro-thiazol-2-yl)-ureido]-3-cyclopentanecarbonyl-phenyl}-acetic acid (204 mg, 0.5 mmol) and 2-methanesulfonyl-ethylamine (62 mg, 0.5 mmol) following the general procedure K.

LC-MS (m/z): 514 ($M+1$)⁺; ¹H NMR (400 MHz, DMSO- d_6): δ 1.75 (m, 4H), 1.91 (m, 4H), 2.97 (s, 3H), 3.07 (q, 2 H), 3.23 (t, 2H), 3.42 (s, 2H), 3.84 (m, 1H), 7.42 (s, 1H), 7.44 (d, 1H), 7.94 (s, 1H), 8.18 (d, 1 H), 8.35 (t, 1H), 10.77 (s, 1 H), 12.01 (br, 1H),

5

Example 224

[2-(3-(2-Cyclopentanecarbonyl-4-[(2-methanesulfonyl-ethylcarbamoyl)-methyl]-phenyl)-ureido)-thiazol-4-yl]-acetic acid ethyl ester




10 [2-(3-(2-Cyclopentanecarbonyl-4-[(2-methanesulfonyl-ethylcarbamoyl)-methyl]-phenyl)-ureido)-thiazol-4-yl]-acetic acid ethyl ester (174 mg, 65%) was prepared from 2-(4-amino-3-cyclopentanecarbonyl-phenyl)-N-(2-methanesulfonyl-ethyl)-acetamide (176 mg, 0.5 mmol) and (2-amino-thiazol-4-yl)-acetic acid ethyl ester (112 mg, 0.6 mmol) following the general procedure D.

15 LC-MS (m/z): 565 ($M+1$)⁺; ¹H NMR (400 MHz, CDCl₃): δ 1.28 (t, 3H), 1.65 (m, 4H), 1.79 (m, 2H), 1.88 (m, 2H), 3.01 (s, 3H), 3.31 (t, 2H), 3.53 (s, 2H), 3.68 (s, 2H), 3.83 (m, 1H), 4.20 (m, 3H), 7.42 (s, 1H), 7.44 (d, 1H), 7.94 (s, 1H), 8.18 (d, 1 H), 8.35 (t, 1H), 10.77 (s, 1 H), 12.01 (br, 1H),

Example 225

2-(3-(2-Cyclopentanecarbonyl-4-[2-methanesulfonyl-ethyl carbamoyl]-methyl]-phenyl)-ureido)-thiazol-4-yl-acetic acid




2-(3-(2-Cyclopentanecarbonyl-4-[2-methanesulfonyl-ethyl carbamoyl]-methyl]-phenyl)-ureido)-thiazol-4-yl-acetic acid (214 mg, 80%) was prepared from 2-(3-(2-cyclopentanecarbonyl-4-[2-methanesulfonyl-ethyl carbamoyl]-methyl]-phenyl)-ureido)-thiazol-4-yl-acetic acid ethyl ester (282 mg, 0.5 mmol) following the general procedure J.

LC-MS (m/z): 537 ($M+1$)⁺; ¹H NMR (400 MHz, CDCl₃): δ 1.65 (m, 4H), 1.79 (m, 2H), 1.88 (m, 2H), 2.97 (s, 3H), 3.22 (t, 2H), 3.45 (m, 2H), 3.47 (s, 2H), 3.56 (s, 2H), 3.82 (m, 1H), 6.84 (s, 1H), 7.42 (d, 1H), 7.92 (s, 1H), 8.18 (d, 1H), 8.33 (t, 1H), 10.70 (s, 1H), 11.84 (br, 1H),

Example 226

3-Cyclopentanecarbonyl-4-(3-[1,3,4]thiadiazol-2-yl-ureido)-phenyl]-acetic acid ethyl ester




3-Cyclopentanecarbonyl-4-(3-[1,3,4]thiadiazol-2-yl-ureido)-phenyl]-acetic acid ethyl ester (147 mg, 73 %) was prepared from (4-amino-3-cyclopentanecarbonyl-phenyl)-acetic acid ethyl ester (138 mg, 0.5 mmol) and [1,3,4]thiadiazol-2-ylamine (60 mg, 0.6 mmol) following the general procedure D.

LC-MS (m/z): 403 ($M+1$)⁺; ¹H NMR (400 MHz, CDCl₃): δ 1.27 (t, 6H), 1.67 (m, 4H), 1.91 (m, 4H), 3.64 (s, 2H), 3.77 (m, 1H), 4.17 (m, 4H), 7.46 (d, 1H), 7.89 (s, 2H), 8.47 (s, 1H), 8.77 (s, 1H), 10.40 (br, 1H), 11.89 (br, 1H).

5

Example 227

1-[2-Cyclopentanecarbonyl-4-(2-morpholin-4-yl-2-oxo-ethyl)-phenyl]-3-thiazol-2-yl-urea




10 1-[2-Cyclopentanecarbonyl-4-(2-morpholin-4-yl-2-oxo-ethyl)-phenyl]-3-thiazol-2-yl-urea (151 mg, 68 %) was prepared from [3-cyclopentanecarbonyl-4-(3-thiazol-2-yl-ureido)-phenyl]-acetic acid (186 mg, 0.5 mmol) and morpholine (44 mg, 0.5 mmol) following the general procedure K.

LC-MS (m/z): 443 ($M+1$)⁺; ¹H NMR (400 MHz, CDCl₃/DMSO-*d*₆): δ 1.67 (m, 4H), 1.88 (m, 4H), 3.51 (t, 4H), 3.69 (t, 4H), 3.74 (m, 3H), 6.91 (d, 1H), 7.37 (dd, 1H), 7.64 (d, 1H),
15 7.84 (s, 1H), 8.52 (br, 2H), 11.32 (br, 1H).

Example 228

1-[2-Cyclopentanecarbonyl-4-[2-(4-methyl-piperazin-1-yl)-2-oxo-ethyl]-phenyl]-3-thiazol-2-yl-urea



20


1-[2-Cyclopentanecarbonyl-4-[2-(4-methyl-piperazin-1-yl)-2-oxo-ethyl]-phenyl]-3-thiazol-2-yl-urea (137 mg, 60 %) was prepared from [3-cyclopentanecarbonyl-4-(3-thiazol-2-

yl-ureido)-phenyl]-acetic acid (186 mg, 0.5 mmol) and N-methyl piperazine (50 mg, 0.5 mmol) following the general procedure K.

LC-MS (*m/z*): 456 (M+1)⁺; ¹H NMR (400 MHz, CDCl₃/DMSO-*d*₆): δ 1.67 (m, 4H), 1.89 (m, 4H), 2.29 (s, 3H), 3.51 (t, 4H), 3.61 (s, 2H), 3.69 (m, 5H), 6.92 (d, 1H), 7.38 (d, 1 H),
 5 7.58(s, 1H), 7.84 (s, 1H), 8.53 (d, 1H), 11.56 (br, 1H), 11.67 (br, 1H).

Example 229

{2-[3-Cyclopentanecarbonyl-4-(3-thiazol-2-yl-ureido)-phenyl]-acetylamino}-acetic acid



10


{2-[3-Cyclopentanecarbonyl-4-(3-thiazol-2-yl-ureido)-phenyl]-acetylamino}-acetic acid tert.butyl ester (158 mg, 65 %) was prepared from [3-cyclopentanecarbonyl-4-(3-thiazol-2-yl-ureido)-phenyl]-acetic acid (186 mg, 0.5 mmol) and glycine ter.butyl ester (60 mg, 0.5 mmol) following the general procedure K. The ester intermediate (122 mg, 0.25 mmol) upon
 15 hydrolysis with TFA furnished {2-[3-cyclopentanecarbonyl-4-(3-thiazol-2-yl-ureido)-phenyl]-acetylamino}-acetic acid (86 mg, 80%).

20

LC-MS (*m/z*): 431 (M+1)⁺; ¹H NMR (400 MHz, CDCl₃/DMSO-*d*₆): δ 1.67 (m, 4H), 1.94 (m, 4H), 3.69 (dd, 2H), 3.82 (m, 3H), 3.95 (t, 1H), 6.99 (m, 1H), 7.32 (dd, 1 H), 7.39 (d, 1H), 7.75 (d, 1H), 8.45 (d, 1H), 12.16 (br, 1H).

Example 230

{2-[3-(2-Cyclopentanecarbonyl-4-methylcarbamoylmethyl-phenyl)-ureido]-thiazol-4-yl}-acetic acid




{2-[3-(2-Cyclopentanecarbonyl-4-methylcarbamoylmethyl-phenyl)-ureido]-thiazol-4-yl}-acetic acid (153 mg, 65 %) was prepared from 2-(4-amino-3-cyclopentane-carbonyl-phenyl)-N-methyl-acetamide (130 mg, 0.5 mmol) and (2-amino-thiazol-4-yl)-acetic acid ethyl ester (112 mg, 0.6 mmol) following the general procedure D. The ester intermediate (236 mg, 0.5 mmol) upon hydrolysis following the general procedure J furnished {2-[3-(2-cyclopentanecarbonyl-4-methylcarbamoylmethyl-phenyl)-ureido]-thiazol-4-yl}-acetic acid (167 mg, 75%).

LC-MS (m/z): 445 ($M+1$)⁺; ¹H NMR (400 MHz, DMSO- d_6): δ 1.62 (m, 4H), 1.74 (m, 2H), 1.90 (m, 2H), 2.57 (d, 3H), 3.43 (s, 2H), 3.56 (s, 2H), 3.81 (m, 1H), 6.85 (d, 2H), 7.41 (d, 1H), 7.92 (s, 1H), 7.96 (d, 1H), 8.18 (d, 1H), 10.69 (s, 1H), 11.96 (br, 1H).

Example 231

{3-Cyclopentanecarbonyl-4-[3-(4-methylcarbamoylmethyl-thiazol-2-yl)-ureido]-phenyl}-acetic acid ethyl ester




{3-Cyclopentanecarbonyl-4-[3-(4-methylcarbamoylmethyl-thiazol-2-yl)-ureido]-phenyl}-acetic acid ethyl ester (153 mg, 65 %) was prepared from (4-amino-3-cyclopentane-carbonyl-phenyl)-acetic acid ethyl ester (138 mg, 0.5 mmol) and 2-(2-amino-thiazol-4-yl)-N-methyl-acetamide (103 mg, 0.6 mmol) following the general procedure D.

LC-MS (m/z): 473 ($M+1$)⁺; ¹H NMR (400 MHz, CDCl₃): δ 1.26 (t, 3H), 1.70 (m, 4H), 1.86 (m, 2H), 1.92 (m, 2H), 2.81 (d, 3H), 3.62 (s, 2H), 3.65 (s, 2H), 3.77 (m, 1H), 4.16 (q, 2H), 6.65 (s, 1H), 6.71 (br, 1H), 7.44 (dd, 1H), 7.88 (s, 1H), 8.52 (d, 1H), 9.39 (br, 1H), 11.71 (br, 1H).

5

Example 232

2-{3-Cyclopentanecarbonyl-4-[3-(4-methylcarbamoylmethyl-thiazol-2-yl)-ureido]-phenyl}-N-methyl-acetamide



10 {3-Cyclopentanecarbonyl-4-[3-(4-methylcarbamoylmethyl-thiazol-2-yl)-ureido]-phenyl}-acetic acid (155 mg, 70 %) was prepared from {3-cyclopentanecarbonyl-4-[3-(4-methylcarbamoylmethyl-thiazol-2-yl)-ureido]-phenyl}-acetic acid ethyl ester (236 mg, 0.5 mmol) following the general procedure J. The acid (222 mg, 0.5 mmol) was coupled with 1 N methyl amine following the general procedure K furnished 2-{3-cyclopentanecarbonyl-4-[3-(4-methylcarbamoylmethyl-thiazol-2-yl)-ureido]-phenyl}-N-methyl-acetamide (137 mg, 60%).


15

LC-MS (m/z): 458 ($M+1$)⁺; ¹H NMR (400 MHz, CDCl₃/DMSO- d_6): δ 1.69 (m, 4H), 1.82 (m, 4H), 2.65 (d, 3H), 3.68 (d, 3H), 3.43 (s, 2H), 3.48 (s, 2H), 3.67 (m, 1H), 6.51 (d, 1H), 7.14 (br, 1H), 7.31 (m, 2H), 7.54 (d, 1H), 7.75 (dd, 1H), 8.31 (d, 1H), 11.13 (br, 1H).

20

Example 233

1-(2-Cyclopentanecarbonyl-4-methyl-phenyl)-3-{4-[2-(pyridin-2-yloxy)-ethyl]-thiazol-2-yl}-urea




To a solution of {2-[3-(2-cyclopentanecarbonyl-4-methyl-phenyl)-ureido]-thiazol-4-yl}-acetic acid (0.38g, 1.0 mmol) in THF (5mL) at 0°C was added 1.0 M solution of BH₃-THF (3.0 mL, 3.0mmol) dropwise. After stirring at 0°C for 1h, the excess BH₃-THF complex was quenched with MeOH. The solution was concentrated under vacuum and purified on silica gel with 1:1 EtOAc/hexanes to obtain the alcohol as a white powder in 80% yield.

1-(2-Cyclopentanecarbonyl-4-methyl-phenyl)-3-{4-[2-(pyridin-2-yloxy)-ethyl]-thiazol-2-yl}-urea (30mg, 50%) was prepared from 1-[2-(cyclopentanecarbonyl-4-methyl-phenyl)]-3-[4-(2-hydroxy-ethyl)-thiazol-2-yl]-urea (0.05g, 0.134mmol), 2-hydroxypyridine (0.013g, 0.143mmol) following the general procedure P.

LC/MS (m/z): 451 (M+1)⁺; ¹H NMR (400 MHz, CDCl₃) δ 1.67-1.72 (m, 4H), 1.89-1.85 (m, 4H), 2.36 (s, 3H), 3.13 (t, 2H), 3.64-3.80 (m, 1H), 4.60 (t, 2H), 6.60 (s, 1H), 6.72 (d, 1H), 6.86 (m, 1H), 7.34 (d, 1H), 7.56 (m, 1H), 7.70 (s, 1H), 8.15 (d, 1H), 8.45 (br, 1H), 11.85 (br, 1H).

15 Example 234

1-(2-Cyclopentanecarbonyl-4-methyl-phenyl)-3-{4-[2-(pyridin-4-yloxy)-ethyl]-thiazol-2-yl}-urea




1-(2-Cyclopentanecarbonyl-4-methyl-phenyl)-3-{4-[2-(pyridin-4-yloxy)-ethyl]-thiazol-2-yl}-urea (50mg, 42%) was prepared from 1-[2-(cyclopentanecarbonyl-4-methyl-phenyl)]-3-[4-(2-hydroxy-ethyl)-thiazol-2-yl]-urea (0.10g, 0.27mmol) and 4-hydroxypyridine (0.05g, 0.54mmol) following the general procedure P.

LC/MS (m/z): 451 (M+1)⁺; ¹H NMR (400 MHz, CDCl₃) δ 1.62-1.73 (m, 4H), 1.80-1.85 (m, 4H), 2.37 (s, 3H), 2.85 (t, 2H), 3.87 (t, 2H), 3.75 (br, 1H), 6.52 (s, 1H), 7.46 (d, 2H), 7.48 (d, 2H), 7.52 (d, 1H), 7.70 (s, 1H), 8.44 (d, 1H),

EXAMPLE 235

5-(2-{2-[3-(2-Cyclopentanecarbonyl-4-methyl-phenyl)-ureido]-thiazol-4-yl} ethoxy) nicotinic acid methyl ester




5 5-(2-{2-[3-(2-Cyclopentanecarbonyl-4-methyl-phenyl)-ureido]-thiazol-4-yl} ethoxy) nicotinic acid methyl ester (30mg, 44%) was prepared from 1-[2-(cyclopentanecarbonyl-4-methyl-phenyl)-3-[4-(2-hydroxy-ethyl)-thiazol-2-yl]-urea (0.05g, 0.134mmol) and 5-hydroxynicotinic acid methyl ester (0.04g, 0.27mmol) following the general procedure P.

LC/MS (m/z): 509 (M+1)⁺; ¹H NMR (400 MHz, CD₃OD) δ 1.65-1.74 (br, 4H), 1.83-1.89 (br, 4H), 2.35 (s, 3H), 3.15 (t, 2H), 3.74 (br, 1H), 3.92 (s, 3H), 4.42 (t, 2H), 6.75 (s, 1H), 7.72 (s, 1H), 7.85 (d, 1H), 8.27 (s, 1H), 8.42 (d, 1H), 8.57 (s, 1H), 8.67 (s, 1H).

EXAMPLE 236

5-(2-{2-[3-(2-Cyclopentanecarbonyl-4-methyl-phenyl)-ureido]-thiazol-4-yl} ethoxy) nicotinic acid




15 5-(2-{2-[3-(2-Cyclopentanecarbonyl-4-methyl-phenyl)-ureido]-thiazol-4-yl} ethoxy) nicotinic acid (23mg, 95%) was prepared from 5-(2-{2-[3-(2-cyclopentanecarbonyl-4-methyl-phenyl)-ureido]-thiazol-4-yl} ethoxy) nicotinic acid methyl ester (0.025g, 0.05mmol) and 2.5M LiOH (20ul) following the general procedure J.

20 LC/MS (m/z): 495 (M+1)⁺; ¹H NMR (400 MHz, CD₃OD) δ 1.65-1.75 (br, 4H), 1.83-1.90 (br, 4H), 2.31 (s, 3H), 3.05 (t, 2H), 3.72-3.80 (m, 1H), 4.42 (t, 2H), 6.85 (s, 1H), 7.57 (s, 1H), 7.74 (d, 1H), 8.30 (s, 1H), 8.48 (d, 1H), 8.52 (s, 1H), 8.65 (s, 1H), 11.43 (br, 1H).

EXAMPLE 237

4-(2-{2-[3-(2-Cyclopentanecarbonyl-4-methyl-phenyl)-ureido]-thiazol-4-yl} ethoxy) benzoic acid methyl ester




5 4-(2-{2-[3-(2-Cyclopentanecarbonyl-4-methyl-phenyl)-ureido]-thiazol-4-yl} ethoxy) benzoic acid methyl ester (50mg, 37%) was prepared from 1-[2-(cyclopentanecarbonyl-4-methyl-phenyl]-3-[4-(2-hydroxy-ethyl)-thiazol-2-yl]-urea (0.10g, 0.27mmol) and 4-hydroxybenzoic acid methyl ester (0.08g, 0.54mmol) following the general procedure P.

LC/MS (m/z): 508 (M+1)⁺ ; ¹H NMR (400 MHz, CDCl₃) δ 1.64-1.78 (m, 4H), 1.82-1.90 (m, 4H), 2.32 (s, 3H), 3.04 (t, 2H), 3.60-3.85 (m, 1H), 3.92 (s, 3H), 4.32 (t, 2H), 6.81 (s, 1H), 6.98 (d, 2H), 7.33 (d, 1H), 7.80 (s, 1H), 7.86 (d, 2H), 8.15 (br, 1H), 10.70 (br, 1H), 11.88 (br, 1H).

EXAMPLE 238

15 4-(2-{2-[3-(2-Cyclopentanecarbonyl-4-methyl-phenyl)-ureido]-thiazol-4-yl} ethoxy) benzoic acid




20 4-(2-{2-[3-(2-cyclopentanecarbonyl-4-methyl-phenyl)-ureido]-thiazol-4-yl} ethoxy) benzoic acid (19mg, 96%) was prepared from 4-(2-{2-[3-(2-cyclopentanecarbonyl-4-methyl-phenyl)-ureido]-thiazol-4-yl} ethoxy) benzoic acid methyl ester (0.020g, 0.04mmol) and 2.5M LiOH (20uL) following the general procedure J.

LC/MS (m/z): 494 (M+1)⁺ ; ¹H NMR (400 MHz, DMSO) δ 1.65-1.78 (m, 4H), 1.80-1.92 (m, 4H), 2.32 (s, 3H), 3.04 (t, 2H), 3.60-3.85 (m, 1H), 4.34 (t, 2H), 6.82 (s, 1H), 7.01 (d, 2H), 7.36 (d, 2H), 7.81 (s, 1H), 7.86 (d, 2H), 8.13 (br, 1H), 8.88 (s, 1H), 10.66 (br, 1H).

EXAMPLE 239

[4-(2-{2-[3-(2-Cyclopentanecarbonyl-4-methyl-phenyl)-ureido]-thiazol-4-yl} ethoxy)-phenyl] acetic acid methyl ester



5

[4-(2-{2-[3-(2-Cyclopentanecarbonyl-4-methyl-phenyl)-ureido]-thiazol-4-yl} ethoxy)-phenyl] acetic acid methyl ester (56mg, 40%) was prepared from 1-[2-(cyclopentanecarbonyl-4-methyl-phenyl)-3-[4-(2-hydroxy-ethyl)-thiazol-2-yl]-urea (0.10g, 0.27mmol) and 4-hydroxyphenylacetic acid methyl ester (0.09g, 0.54mmol) following the general procedure P.


10

LC/MS (M/Z): 522 (M+1)⁺.

Example 240

[4-(2-{2-[3-(2-Cyclopentanecarbonyl-4-methyl-phenyl)-ureido]-thiazol-4-yl} ethoxy)-phenyl] acetic acid

15



[4-(2-{2-[3-(2-Cyclopentanecarbonyl-4-methyl-phenyl)-ureido]-thiazol-4-yl} ethoxy)-phenyl] acetic acid (25mg, 86%) was prepared from [4-(2-{2-[3-(2-Cyclopentanecarbonyl-4-methyl-phenyl)-ureido]-thiazol-4-yl} ethoxy)-phenyl] acetic acid methyl ester (0.03g, 0.06mmol) and 2.5M LiOH (20uL) following the general procedure J.


20

LC/MS (m/z): 508 (M+1)⁺; ¹H NMR (400 MHz, CDCl₃) δ 1.65-1.76 (m, 4H), 1.83-1.95 (m, 4H), 2.36 (s, 3H), 3.10 (t, 2H), 3.70-3.82 (m, 1H), 4.24 (t, 2H), 4.96-5.02 (m, 2H), 6.50 (s, 1H), 6.88 (d, 2H), 7.20 (d, 2H), 7.30 (d, 1H), 7.68 (s, 1H), 8.45 (br, 1H), 11.45 (br, 1H).

25

Example 241

1-(2-Cyclopentanecarbonyl-4-methyl-phenyl)-3-[4-(2-morpholin-4-yl-ethyl)-thiazol-2-yl]-urea




To 1-(2-cyclopentanecarbonyl-4-methyl-phenyl)-3-[4-(2-hydroxy-ethyl)-thiazol-2-yl]-urea (2.68 mmol) in CH_2Cl_2 (20mL) at 0 °C was added Et_3N (1.64 mL, 11.70 mmol), dimethyl sulfoxide (10 mL) followed by sulfur trioxide-pyridine (1.46 g, 9.21 mmol). The reaction mixture was stirred for 1h and then poured into water (30 mL). The mixture was extracted with CH_2Cl_2 and the organic layers was washed with 1.0 N ammonium chloride (2 x 20 mL), water (2 x 20 mL), brine (1 x20 mL), dried (Na_2SO_4) and concentrated to give a solid. The solid was purified by column chromatography (silica, Hexanes/ EtOAc, 20-50%) to obtain the aldehyde as a white solid in 50% yield.

1-(2-Cyclopentanecarbonyl-4-methyl-phenyl)-3-[4-(2-morpholin-4-yl-ethyl)-thiazol-2-yl]-urea (15mg, 31%) was prepared from 1-(2-Cyclopentanecarbonyl-4-methyl-phenyl)-3-[4-(2-oxo-ethyl)-thiazol-2-yl]-urea (0.04g, 0.11mmol) and morpholine (0.009g, 0.11mmol) following the general procedure O.

LC/MS (m/z): 443 ($M+1$)⁺; ^1H NMR (400 MHz, CDCl_3) δ 1.68-1.72 (m, 4H), 1.88-1.91 (m, 4H), 2.36 (s, 3H), 2.53 (br, 4H), 2.72 (t, 2H), 2.85 (t, 2H), 3.72-3.74 (m, 5H), 6.52 (s, 1H), 7.35 (d, 1H), 7.70 (s, 1H), 8.45 (br, 1H), 11.62 (br, 1H).

Example 242

1-(2-Cyclopentanecarbonyl-4-methyl-phenyl)-3-[4-[2-(2-morpholin-4-yl-ethylamino)-ethyl]-thiazol-2-yl]-urea




1-(2-Cyclopentanecarbonyl-4-methyl-phenyl)-3-{4-[2-(2-morpholin-4-yl-ethylamino)-ethyl]-thiazol-2-yl}-urea (23mg, 44%) was prepared from (0.04g, 0.11mmol) and 4-ethylamino morpholine (0.014g, 0.11mmol) following the general procedure O.

5 LC/MS (m/z): 486 (M+1)⁺; ¹H NMR (400 MHz, CDCl₃) δ 1.69-1.73 (m, 4H), 1.88-1.92 (m, 4H), 2.35 (s, 3H), 2.50 (t, 2H), 2.53 (br, 4H), 2.72 (t, 2H), 2.80 (t, 2H), 2.85 (t, 2H), 3.72-3.74 (m, 5H), 6.53 (s, 1H), 7.35 (d, 1H), 7.70 (s, 1H), 8.45 (br, 1H), 11.60 (br, 1H).

10 Example 243

1-(2-{2-[3-(2-Cyclopentanecarbonyl-4-methyl-phenyl)-ureido]-thiazol-4-yl}-ethyl)-piperidin-3-carboxylic acid ethyl ester




15 1-(2-{2-[3-(2-cyclopentanecarbonyl-4-methyl-phenyl)-ureido]-thiazol-4-yl}-ethyl)-piperidin-3-carboxylic acid ethyl ester (28mg, 50%) was prepared from (0.04g, 0.11mmol) and (s)-(+)-nipecotic acid (0.017g, 0.11mmol) following the general procedure O.

LC/MS (m/z): 528 (M+1)⁺.

Example 244

1-(2-{2-[3-(2-Cyclopentanecarbonyl-4-methyl-phenyl)-ureido]-thiazol-4-yl}-ethyl)-piperidin-3-carboxylic acid




5 1-(2-{2-[3-(2-Cyclopentanecarbonyl-4-methyl-phenyl)-ureido]-thiazol-4-yl}-ethyl)-piperidin-3-carboxylic acid (18mg, 95%) was prepared from 1-(2-{2-[3-(2-cyclopentanecarbonyl-4-methyl-phenyl)-ureido]-thiazol-4-yl}-ethyl)-piperidin-3-carboxylic acid ethyl ester (0.02g, 0.04mmol) and 2.5M LiOH (20uL) following the general procedure J.

10 LC/MS (m/z): 513 (M+1)⁺; ¹H NMR (400 MHz, CDCl₃) δ 1.60-1.69 (m, 4H), 1.80-1.98 (m, 4H), 2.34 (s, 3H), 2.90-3.05 (m, 5H), 3.10-3.20 (m, 4H), 3.65-3.75 (m, 1H), 6.49 (s, 1H), 7.34 (d, 1H), 7.63 (s, 1H), 8.25 (d, 1H), 11.40 (br, 1H).

Example 245

15 1-(2-Cyclopentanecarbonyl-4-methyl-phenyl)-3-[4-(2-piperazin-1-yl-ethyl)-thiazol-2-yl]-urea




20 1-(2-Cyclopentanecarbonyl-4-methyl-phenyl)-3-[4-(2-piperazin-1-yl-ethyl)-thiazol-2-yl]-urea (35mg, 70%) was prepared from methanesulfonic acid 2-{2-[3-(2-cyclopentanecarbonyl-4-methyl-phenyl)-ureido]-thiazol-4-yl}-ethyl ester (0.06g, 0.13mmol) and piperazine (0.50g, 5.82mmol) following the general procedure Z.

LC/MS (m/z): 442 (M+1)⁺; ¹H NMR (400 MHz, CDCl₃) δ 1.66-1.72 (m, 4H), 1.86-1.90 (m, 4H), 2.36 (s, 3H), 2.45-2.60 (br, 4H), 2.71 (t, 2H), 2.84 (t, 2H), 2.89-2.98 (m, 4H), 3.70-3.80 (m, 1H), 6.47 (s, 1H), 7.34 (d, 1H), 7.69 (s, 1H), 8.45 (br, 1H), 11.45 (br, 1H).

5 **Example 246**

1-(2-Cyclopentanecarbonyl-4-methyl-phenyl)-3-{4-[2-(4-methanesulfonyl-piperazin-1-yl-ethyl)-thiazol-2-yl]-urea




10 1-(2-Cyclopentanecarbonyl-4-methyl-phenyl)-3-{4-[2-(4-methanesulfonyl-piperazin-1-yl-ethyl)-thiazol-2-yl]-urea (14mg, 40%) was prepared from 1-(2-cyclopentanecarbonyl-4-methyl-phenyl)-3-[4-(2-piperazin-1-yl-ethyl)-thiazol-2-yl]-urea (0.03g, 0.07mmol) and sulfonyl chloride (0.008g, 0.068 mmol) following the general procedure T.

15 LC/MS (m/z): 520 (M+1)⁺; ¹H NMR (400 MHz, CDCl₃) δ 1.65-1.75 (m, 4H), 1.85-1.95 (m, 4H), 2.37 (s, 3H), 2.60-2.64 (br, 4H), 2.76-2.95 (m, 4H), 2.78 (s, 3H), 3.25 (br, 4H), 3.70-3.80 (m, 1H), 6.50 (s, 1H), 7.35 (d, 1H), 7.70 (s, 1H), 8.45 (br, 1H), 11.65 (br, 1H).

Example 247

20 1-[4-(2-Amino-ethyl)-thiazol-2-yl]-3-(2-cyclopentanecarbonyl-4-methyl-phenyl)-urea




1-[4-(2-Amino-ethyl)-thiazol-2-yl]-3-(2-cyclopentanecarbonyl-4-methyl-phenyl)-urea (0.15g, 94%) was prepared from 1-[4-(2-azido-ethyl)-thiazol-2-yl]-3-(2-cyclopentanecarbonyl-4-methyl-phenyl)-urea (0.17g, 0.43mmol) following the general procedure T.

5 LC/MS (m/z): 373 (M+1)⁺ ; ¹H NMR (400 MHz, CDCl₃) δ 1.68-1.74 (m, 4H), 1.88-1.93 (m, 4H), 2.36 (s, 3H), 2.84 (t, 2H), 3.28 (t, 2H), 3.72-3.80 (m, 1H), 4.5 (br, 2H), 6.47 (s, 1H), 7.34 (d, 1H), 7.69 (s, 1H), 8.55 (d, 1H), 11.34 (br, 1H).

Example 248

10 3-(2-{2-[3-(2-Cyclopentanecarbonyl-4-methyl-phenyl)-ureido]-thiazol-4-yl}-ethyl amino)-propionic acid




15 1-[4-(amino-ethyl)-thiazol-2-yl]-3-(2-cyclopentanecarbonyl-4-methyl-phenyl)-urea (0.03g, 0.08mmol) and succinic anhydride (0.02g, 0.16mmol) in 10mL DCM were refluxed for 2h. Reaction was concentrated and purified on silica gel using 5% MeOH/EtOAc to yield the pure product (20mg, 53%).

LC/MS (m/z): 473 (M+1)⁺ ; ¹H NMR (400 MHz, DMSO) δ 1.60-1.75 (m, 4H), 1.85-1.92 (m, 4H), 2.38 (s, 3H), 2.67 (t, 2H), 3.28-3.32 (m, 4H), 3.80-3.92 (br, 1H), 4.02 (t, 2H), 6.70 (s, 1H), 7.38 (d, 1H), 7.83 (s, 1H), 8.15 (d, 1H), 10.65 (br, 1H).

20

Example 249

(2-{2-[3-(2-Cyclopentanecarbonyl-4-methyl-phenyl)-ureido]-thiazol-4-yl}-ethyl amino)-acetic acid methyl ester




To 1-[4-(amino-ethyl)-thiazol-2-yl]-3-(2-cyclopentanecarbonyl-4-methyl-phenyl)-urea (0.03g, 0.08mmol) in 10mL DCM was added methyl bromoacetate (0.01g, 0.08mmol) followed by Et₃N (0.008g, 0.08mmol). After stirring at room temperature overnight, reaction was concentrated and purified by flash chromatography on silica gel with 5%MeOH/EtOAc to obtain the product in 50% yield.

LC/MS (m/z): 445 (M+1)⁺

Example 250

(2-{2-[3-(2-Cyclopentanecarbonyl-4-methyl-phenyl)-ureido]-thiazol-4-yl}-ethyl amino)-acetic acid




(2-{2-[3-(2-cyclopentanecarbonyl-4-methyl-phenyl)-ureido]-thiazol-4-yl}-ethyl amino)-acetic acid (18mg, 95%) was prepared from (2-{2-[3-(2-cyclopentanecarbonyl-4-methyl-phenyl)-ureido]-thiazol-4-yl}-ethyl amino)-acetic acid ethyl ester (0.02g, 0.05mmol) and 2.5M LiOH (20uL) following the general procedure J.

LC/MS (m/z): 431 (M+1)⁺; ¹H NMR (400 MHz, CDCl₃) δ 1.63-1.74 (m, 4H), 1.86-1.92 (m, 4H), 2.36 (s, 3H), 2.81 (t, 2H), 3.07 (t, 2H), 3.59 (s, 2H), 3.69-3.76 (m, 1H), 6.54 (s, 1H), 7.33 (d, 1H), 7.70 (s, 1H), 8.46 (br, 1H), 11.60 (br, 1H).

Example 251

N-(2-{2-[3-(2-Cyclopentanecarbonyl-4-methyl-phenyl)-ureido]-thiazol-4-yl}-ethyl)-acetamide




To 1-[4-(amino-ethyl)-thiazol-2-yl]-3-(2-cyclopentanecarbonyl-4-methyl-phenyl)-urea
5 (0.05g, 0.13mmol) in 10mL DCM at 0°C was added acetyl chloride (0.02g, 0.27mmol)
followed by pyridine (0.02g, 0.21mmol). After stirring for 3h, reaction was concentrated
and purified by flash chromatography on silica gel with 50%EtOAc/hexane to obtain the
product in 64% yield.

LC/MS (m/z): 415 (M+1)⁺; ¹H NMR (400 MHz, CDCl₃) δ 1.65-1.75 (m, 4H), 1.84-
10 1.90 (m, 4H), 1.97 (s, 3H), 2.37 (s, 3H), 2.84 (t, 2H), 3.58 (t, 2H), 3.70-3.80 (m, 1H), 6.54 (s,
1H), 7.34(d, 1H), 7.70 (s, 1H), 8.45 (br, 1H), 11.65 (br, 1H).

Example 252

N-(2-{2-[3-(2-Cyclopentanecarbonyl-4-methyl-phenyl)-ureido]-thiazol-4-yl}-ethyl)-
15 methanesulfonamide




To 1-[4-(amino-ethyl)-thiazol-2-yl]-3-(2-cyclopentanecarbonyl-4-methyl-phenyl)-urea
(0.05g, 0.13mmol) in 10mL DCM at 0°C was added methanesulfonyl chloride (0.02g,
0.13mmol) followed by pyridine (0.02g, 0.21mmol). The reaction was stirred at ambient
20 temperature overnight, concentrated and purified by flash chromatography on silica gel
with 20-50%EtOAc/hexane to obtain the product as white solid in 75% yield.

LC/MS (m/z): 451 (M+1)⁺; ¹H NMR (400 MHz, DMSO) δ 1.62-1.76 (m, 4H), 1.82-1.92 (m, 4H), 2.32 (s, 3H), 2.75 (t, 2H), 2.84 (s, 3H), 3.80-3.90 (m, 1H), 4.10 (t, 2H), 6.76 (s, 1H), 7.37 (d, 1H), 7.83 (s, 1H), 8.16 (d, 1H), 10.64 (br, 1H), 11.80 (br, 1H).

5 Example 253

N-(2-{2-[3-(2-Cyclopentanecarbonyl-4-methyl-phenyl)-ureido]-thiazol-4-yl}-ethyl)-2-dimethylamino acetamide




N-(2-{2-[3-(2-Cyclopentanecarbonyl-4-methyl-phenyl)-ureido]-thiazol-4-yl}-ethyl)-2-dimethylamino acetamide (37mg, 61%) was prepared from 1-[4-(amino-ethyl)-thiazol-2-yl]-3-(2-cyclopentanecarbonyl-4-methyl-phenyl)-urea (0.05g, 0.13mmol), dimethylamino acetic acid (0.02g, 0.13mmol) following the general procedure K.

LC/MS (m/z): 458 (M+1)⁺; ¹H NMR (400 MHz, CD₃OD) δ 1.63-1.70 (m, 4H), 1.80-1.95 (m, 4H), 2.36 (s, 3H), 2.70 (s, 6H), 2.85 (t, 2H), 3.60 (t, 2H), 3.62 (s, 2H), 3.86 (br, 1H), 6.65 (s, 1H), 7.38-7.28 (m, 2H), 7.65-7.74 (m, 1H), 7.80 (br, 1H), 8.20 (br, 1H),

Example 254

N-(2-{2-[3-(2-Cyclopentanecarbonyl-4-methyl-phenyl)-ureido]-thiazol-4-yl}-ethyl)-succinamic acid




1-[4-(amino-methyl)-thiazol-2-yl]-3-(2-cyclopentanecarbonyl-4-methyl-phenyl)-urea (0.10g, 0.28mmol) and succinic anhydride (0.08g, 0.78mmol) in 10mL DCM were

refluxed for 2h. Reaction was concentrated and purified on silica gel using 5% MeOH/EtOAc to yield the desired product (77mg, 60%).

LC/MS (m/z): 459 (M+1)⁺ ; ¹H NMR (400 MHz, CD₃OD) δ 1.70-1.77 (m, 4H), 1.85-1.95 (m, 4H), 2.39 (s, 3H), 2.62-2.72 (m, 4H), 3.71 (s, 2H), 3.80-3.90 (m, 1H), 6.08 (s, 1H),
5 7.40- (d, 1H), 7.75 (s, 1H), 8.70 (d, 1H), 11.30 (s, 1H),

Example 255


3-{2-[3-(2-Cyclopentanecarbonyl-4-methyl-phenyl)-ureido]-thiazol-4-yl}-acrylic acid ethyl ester



10

3-{2-[3-(2-Cyclopentanecarbonyl-4-methyl-phenyl)-ureido]-thiazol-4-yl}-acrylic acid ethyl ester (70mg, 58%) was prepared from 1-(2-Cyclopentanecarbonyl-4-methyl-phenyl)-3-[4-(1-oxo-methyl)-thiazol-2-yl]-urea (0.10g, 0.28mmol) and (carbethoxymethylene)-triphenylphosphorane (0.12g, 0.34mmol) following the general
15 procedure X.

LC/MS (m/z): 428 (M+1)⁺ ; ¹H NMR (400 MHz, CDCl₃) δ 1.29 (t, 3H), 1.67-1.75 (m, 4H), 1.80-1.95 (m, 4H), 2.37 (s, 3H), 3.70-3.80 (m, 1H), 4.19-4.22 (q, 2H), 6.56 (d, 1H), 7.07 (s, 1H), 7.36 (d, 1H), 7.50 (d, 1H), 7.72 (s, 1H), 8.45 (d, 1H),


EXAMPLE 256**3-{2-[3-(2-CYCLOPENTANECARBONYL-4-METHYL-PHENYL)-UREIDO]-THIAZOL-4-YL}-PROPIONIC ACID ETHYL ESTER**

5 3-{2-[3-(2-Cyclopentanecarbonyl-4-methyl-phenyl)-ureido]-thiazol-4-yl}-propionic acid ethyl ester (48mg, 96%) was prepared by hydrogenation of 3-{2-[3-(2-Cyclopentanecarbonyl-4-methyl-phenyl)-ureido]-thiazol-4-yl}-acrylic acid ethyl ester (0.05g, 0.12mmol) with Pd/C.

LC/MS (m/z): 430 (M+1)⁺; ¹H NMR (400 MHz, CDCl₃) δ 1.22 (t, 3H), 1.64-1.70 (m, 4H), 1.80-1.95 (m, 4H), 2.35 (s, 3H), 2.69 (t, 2H), 2.98 (t, 2H), 3.67-3.80 (m, 1H), 4.01-4.15 (q, 2H), 6.50 (s, 1H), 7.32 (d, 1H), 7.68 (s, 1H), 8.45 (br, 1H), 11.60 (br, 1H).

EXAMPLE 257

3-{2-[3-(2-Cyclopentanecarbonyl-4-methyl-phenyl)-ureido]-thiazol-4-yl}-propionic acid




15

3-{2-[3-(2-Cyclopentanecarbonyl-4-methyl-phenyl)-ureido]-thiazol-4-yl}-propionic acid (15mg, 88%) was prepared from 3-{2-[3-(2-Cyclopentanecarbonyl-4-methyl-phenyl)-ureido]-thiazol-4-yl}-propionic acid ethyl ester (0.03g, 0.04mmol) and 2.5M LiOH (20 μL) following the general procedure J.

LC/MS (m/z): 402 (M+1)⁺ ; ¹H NMR (400 MHz, CD₃OD) δ 1.70-1.80 (m, 4H), 1.82-1.95 (m, 4H), 2.34 (s, 3H), 2.58 (t, 2H), 2.80 (t, 2H), 3.82-3.94 (m, 1H), 6.69 (s, 1H), 7.38 (d, 1H), 7.85 (s, 1H), 8.19 (br, 1H), 10.65 (br, 1H).

EXAMPLE 258

- 5 4-{2-[3-(2-Cyclopentanecarbonyl-4-methyl-phenyl)-ureido]-thiazol-4-yl}-but-2-enoic acid ethyl ester




- 10 4-{2-[3-(2-Cyclopentanecarbonyl-4-methyl-phenyl)-ureido]-thiazol-4-yl}-but-2-enoic acid ethyl ester (50mg, 42%) was prepared from 1-(2-Cyclopentanecarbonyl-4-methyl-phenyl)-3-[4-(2-oxo-ethyl)-thiazol-2-yl]-urea (0.10g, 0.27mmol) and (carbethoxymethylene)-triphenylphosphorane (0.09g, 0.27mmol) following the general procedure X.

- 15 LC/MS (m/z): 442 (M+1)⁺ ; ¹H NMR (400 MHz, CDCl₃) δ 1.28 (t, 3H), 1.67-1.73 (m, 4H), 2.37 (s, 3H), 3.56 (d, 1H), 3.70-3.80 (m, 1H), 4.15-4.21 (q, 2H), 5.88 (d, 1H), 6.55 (s, 1H), 7.37 (d, 1H), 7.65 (s, 1H), 8.43 (br, 1H), 11.65 (br, 1H).

Example 259

- 20 1-(2-Cyclopentanecarbonyl-4-methyl-phenyl)-3-[4-(1H-imidazol-2-ylsulfanylmethyl)-thiazol-2-yl]-urea




1-(2-Cyclopentanecarbonyl-4-methyl-phenyl)-3-[4-(1H-imidazol-2-ylsulfanylmethyl)-thiazol-2-yl]-urea (76 mg, 69%) was prepared from 1-(4-Chloromethyl-thiazol-2-yl)-3-(2-cyclopentanecarbonyl-4-methyl-phenyl)-urea (94 mg, 0.25 mmol) following the general
5 procedure Q.

LC-MS (m/z): 442 ($M+1$)⁺; ¹H NMR (400 MHz, CDCl₃): δ 1.76 (m, 4H), 1.92 (m, 4H), 2.32 (s, 3H), 3.88 (p, 1H), 4.43 (s, 2H), 6.94 (s, 1H), 7.11 (d, 1H), 7.36 (d, 1H), 7.88 (d, 1H), 7.71 (s, 1H), 8.52 (d, 1H), 10.34 (br, 2H), and 11.42 (br, 1H).

10 Example 260

1-(2-Cyclopentanecarbonyl-4-methyl-phenyl)-3-[4-(1-methyl-1H-tetrazol-5-ylsulfanylmethyl)-thiazol-2-yl]-urea




1-(2-Cyclopentanecarbonyl-4-methyl-phenyl)-3-[4-(1-methyl-1H-tetrazol-5-ylsulfanylmethyl)-thiazol-2-yl]-urea (93 mg, 79%) was prepared from 1-(4-chloromethyl-
15

thiazol-2-yl)-3-(2-cyclopentanecarbonyl-4-methyl-phenyl)-urea (94 mg, 0.25 mmol) following the general procedure Q.

LC-MS (*m/z*): 458 (*M*+1)⁺; ¹H NMR (400 MHz, CDCl₃): δ 1.73 (m, 4H), 1.88 (m, 4H), 2.37 (s, 3H), 3.76 (p, 1H), 3.86 (s, 3H), 4.55 (s, 2H), 6.88 (s, 1H), 7.34 (d, 1H), 7.70 (s, 1H), 8.40 (d, 1H), 8.52 (br, 1H), and 11.78 (br, 1H).

Example 261

1-(2-Cyclopentanecarbonyl-4-methyl-phenyl)-3-[4-(9H-purin-6-ylsulfanylmethyl)-thiazol-2-yl]-urea



10


1-(2-Cyclopentanecarbonyl-4-methyl-phenyl)-3-[4-(9H-purin-6-ylsulfanylmethyl)-thiazol-2-yl]-urea (106 mg, 86%) was prepared from 1-(4-chloromethyl-thiazol-2-yl)-3-(2-cyclopentanecarbonyl-4-methyl-phenyl)-urea (94 mg, 0.25 mmol) following the general procedure Q.

LC-MS (*m/z*): 494 (*M*+1)⁺; ¹H NMR (400 MHz, CDCl₃): δ 1.66 (m, 4H), 1.84 (m, 4H), 2.21 (s, 3H), 3.72 (p, 1H), 4.29 (s, 2H), 4.63 (br, 1H), 6.78 (s, 1H), 7.11 (s, 1H), 7.28 (d, 1H), 7.75 (s, 1H), 8.13 (s, 1H), 8.42 (d, 1H), 9.24 (br, 1H), and 11.40 (br, 1H).

Example 262

1-(2-Cyclopentanecarbonyl-4-methyl-phenyl)-3-[4-(9H-purine-6-sulfonylmethyl)-thiazol-2-yl]-urea

20




1-(2-Cyclopentanecarbonyl-4-methyl-phenyl)-3-[4-(9H-purine-6-sulfonylmethyl)-thiazol-2-yl]-urea (78 mg, 74%) was prepared from 1-(2-cyclopentanecarbonyl-4-methyl-phenyl)-3-[4-(9H-purin-6-ylsulfanylmethyl)-thiazol-2-yl]-urea (99 mg, 0.20 mmol) and

5 following the general procedure R to give title compound.

LC-MS (m/z): 526 ($M+1$)⁺; ¹H NMR (400 MHz, CDCl₃): δ 1.58 (m, 4H), 1.66 (m, 2H), 1.74 (m, 2H), 2.32 (s, 3H), 3.74 (p, 1H), 4.58 (s, 2H), 6.71 (d, 1H), 7.06 (d, 1H), 7.16 (s, 1H), 7.42 (d, 1H), 7.72 (br, 1H), 8.18 (d, 1H), 8.98 (s, 1H), 9.88 (br, 1H), and 10.26 (br, 1H).

Example 263

10 1-(2-Cyclopentanecarbonyl-4-methyl-phenyl)-3-[4-(pyridin-4-ylsulfanylmethyl)-thiazol-2-yl]-urea




1-(2-Cyclopentanecarbonyl-4-methyl-phenyl)-3-[4-(pyridin-4-ylsulfanylmethyl)-thiazol-2-yl]-urea (79 mg, 70%) was prepared from 1-(4-chloromethyl-thiazol-2-yl)-3-(2-

15 cyclopentanecarbonyl-4-methyl-phenyl)-urea (94 mg, 0.25 mmol) following the general procedure Q.

LC-MS (m/z): 453 ($M+1$)⁺; ¹H NMR (400 MHz, CDCl₃): δ 1.68 (m, 4H), 1.86 (m, 4H), 2.36 (s, 3H), 3.76 (p, 1H), 4.21 (s, 2H), 6.78 (s, 1H), 7.16-7.20 (dd, 2H), 7.34 (d, 1H), 7.70 (s, 1H), 8.13 (s, 1H), 8.38 (dd, 1H), 8.42 (d, 1H), 9.46 (br, 1H), and 11.70 (br, 1H).

Example 264

5 1-(2-Cyclopentanecarbonyl-4-methyl-phenyl)-3-[4-(pyridine-4-sulfonylmethyl)-thiazol-2-yl]-urea




1-(2-Cyclopentanecarbonyl-4-methyl-phenyl)-3-[4-(pyridine-4-sulfonylmethyl)-thiazol-2-yl]-urea (73 mg, 75%) was prepared from 1-(2-cyclopentanecarbonyl-4-methyl-phenyl)-3-[4-(pyridin-4-ylsulfonylmethyl)-thiazol-2-yl]-urea (90 mg, 0.20 mmol) and following the general procedure R.

LC-MS (m/z): 485 ($M+1$)⁺; ¹H NMR (400 MHz, CDCl₃): δ 1.66 (m, 4H), 1.86 (m, 4H), 2.37 (s, 3H), 3.78 (p, 1H), 4.49 (s, 2H), 6.88 (s, 1H), 7.36 (d, 1H), 7.62 (dd, 2H), 7.72 (s, 1H), 8.42 (d, 1H), 8.86 (d, 2H), 9.40 (br, 1H), and 11.52 (br, 1H).

Example 265

15 (5-{2-[3-(2-Cyclopentanecarbonyl-4-methyl-phenyl)-ureido]-thiazol-4-ylmethylsulfanyl}-tetrazol-1-yl)-acetic acid




(5-{2-[3-(2-Cyclopentanecarbonyl-4-methyl-phenyl)-ureido]-thiazol-4-ylmethylsulfanyl}-tetrazol-1-yl)-acetic acid (100 mg, 80%) was prepared from 1-(4-chloromethyl-thiazol-2-yl)-3-(2-cyclopentanecarbonyl-4-methyl-phenyl)-urea (94 mg, 0.25 mmol) following the general procedure Q.

- 5 LC-MS (*m/z*): 501 (*M*+1)⁺; ¹H NMR (400 MHz, CDCl₃): δ 1.74 (m, 4H), 1.90 (m, 4H), 2.33 (s, 3H), 3.72 (p, 1H), 4.22 (s, 2H), 5.00 (s, 2H), 6.92 (s, 1H), 7.52 (d, 1H), 7.72 (s, 1H), 8.38 (d, 1H), 9.82 (br, 1H), 10.32 (br, 1H), and 11.70 (br, 1H).

Example 266

- 10 (5-{2-[3-(2-Cyclopentanecarbonyl-4-methyl-phenyl)-ureido]-thiazol-4-ylmethylsulfanyl}-4-cyclopropyl-4H-[1,2,4]triazol-3-ylmethyl)-carbamic acid tert-butyl ester




- 15 (5-{2-[3-(2-Cyclopentanecarbonyl-4-methyl-phenyl)-ureido]-thiazol-4-ylmethylsulfanyl}-4-cyclopropyl-4H-[1,2,4]triazol-3-ylmethyl)-carbamic acid tert-butyl ester (136 mg, 89%) was prepared from 1-(4-chloromethyl-thiazol-2-yl)-3-(2-cyclopentanecarbonyl-4-methyl-phenyl)-urea (94 mg, 0.25 mmol) following the general procedure Q.

LC-MS (*m/z*): 612 (*M*+1)⁺; ¹H NMR (400 MHz, CDCl₃): δ 1.04 (m, 2H), 1.14 (m, 2H), 1.40 (s, 9H), 1.66 (m, 4H), 1.84 (m, 4H), 2.35 (s, 3H), 2.88 (p, 1H), 3.70 (p, 1H), 4.52 (s, 2H), 4.54 (s, 2H), 5.78 (br, 1H), 6.80 (s, 1H), 7.34 (d, 1H), 7.66 (s, 1H), 8.32 (d, 1H), 10.24 (br, 1H), and 11.36 (br, 1H).

- 20 **Example 267**

1-[4-(5-Aminomethyl-4-cyclopropyl-4H-[1,2,4]triazol-3-ylsulfanylmethyl)-thiazol-2-yl]-
3-(2-cyclopentanecarbonyl-4-methyl-phenyl)-urea hydrochloride salt




1-[4-(5-Aminomethyl-4-cyclopropyl-4H-[1,2,4]triazol-3-ylsulfanylmethyl)-thiazol-2-yl]-
3-(2-cyclopentanecarbonyl-4-methyl-phenyl)-urea (51 mg, 99%) was prepared from (5-{2-[3-(2-Cyclopentanecarbonyl-4-methyl-phenyl)-ureido]-thiazol-4-ylmethyl sulfanyl}-4-cyclopropyl-4H-[1,2,4]triazol-3-ylmethyl)-carbamic acid tert-butyl ester (61 mg, 0.10 mmol) and hydrochloric acid (4 mL, 4.0 M. in dioxane) following the general deprotection procedure.

LC-MS (*m/z*): 512 (*M*+1)⁺; ¹H NMR (400 MHz, CDCl₃): δ 1.24 (m, 2H), 1.38 (m, 2H),
1.72 (m, 4H), 1.94 (m, 4H), 2.38 (s, 3H), 2.92 (p, 1H), 3.65 (s, 2H), 3.77 (s, 2H), 4.08 (m, 1H),
5.24 (br, 2H), 6.90 (s, 1H), 7.52 (d, 1H), 7.72 (s, 1H), 8.42 (d, 1H), 10.24 (br, 1H), and 11.36 (br, 1H).

Example 268

2-{2-[3-(2-Cyclopentanecarbonyl-4-methyl-phenyl)-ureido]-thiazol-4-ylmethylsulfanyl}-1H-imidazole-4-carboxylic acid ethyl ester




2-{2-[3-(2-Cyclopentanecarbonyl-4-methyl-phenyl)-ureido]-thiazol-4-ylmethylsulfanyl}-
1H-imidazole-4-carboxylic acid ethyl ester (112 mg, 88%) was prepared from 1-(4-
chloromethyl-thiazol-2-yl)-3-(2-cyclopentanecarbonyl-4-methyl-phenyl)-urea (94 mg, 0.25
5 mmol) following the general procedure Q.

LC-MS (m/z): 486 ($M+1$)⁺; ¹H NMR (400 MHz, CDCl₃): δ 1.36 (t, 3H), 1.74 (m, 4H),
1.90 (m, 4H), 2.36 (s, 3H), 3.78 (p, 1H), 4.08 (q, 2H), 4.34 (s, 2H), 6.84 (s, 1H), 7.34 (d, 1H),
7.72 (d, 1H), 7.88 (d, 1H), 8.18 (d, 1H), 8.38 (d, 1H), 10.24 (br, 1H), and 11.88 (br, 1H).

10 **Example 269**

2-{2-[3-(2-Cyclopentanecarbonyl-4-methyl-phenyl)-ureido]-thiazol-4-
ylmethylsulfanyl}-1H-imidazole-4-carboxylic acid




2-{2-[3-(2-Cyclopentanecarbonyl-4-methyl-phenyl)-ureido]-thiazol-4-ylmethylsulfanyl}-
1H-imidazole-4-carboxylic acid (91 mg, 94%) was prepared from 2-{2-[3-(2-
5 cyclopentanecarbonyl-4-methyl-phenyl)-ureido]-thiazol-4-ylmethylsulfanyl}-1H-imidazole-4-
carboxylic acid ethyl ester (102 mg, 0.20 mmol) following the general procedure J.

LC-MS (m/z): 474 ($M+1$)⁺; ¹H NMR (400 MHz, CDCl₃): δ 1.74 (m, 4H), 1.92 (m, 4H),
2.34 (s, 3H), 3.87 (p, 1H), 4.37 (s, 2H), 6.89 (s, 1H), 7.39 (d, 1H), 7.68 (d, 1H), 7.84 (d, 1H),
8.15 (d, 1H), 8.26 (d, 1H), 9.28 (br, 1H), 10.66 (br, 1H), and 11.90 (br, 1H).

10

Example 270

2-{2-[3-(2-Cyclopentanecarbonyl-4-methyl-phenyl)-ureido]-thiazol-4-
ylmethanesulfonyl}-1H-imidazole-4-carboxylic acid ethyl ester



15


2-{2-[3-(2-Cyclopentanecarbonyl-4-methyl-phenyl)-ureido]-thiazol-4-
ylmethanesulfonyl}-1H-imidazole-4-carboxylic acid ethyl ester (35 mg, 64%) was prepared

from 2-{2-[3-(2-cyclopentanecarbonyl-4-methyl-phenyl)-ureido]-thiazol-4-ylmethylsulfanyl}-1H-imidazole-4-carboxylic acid ethyl ester (51 mg, 0.10 mmol) following the general procedure R.

LC-MS (*m/z*): 546 (*M*+1)⁺; ¹H NMR (400 MHz, CDCl₃): δ 1.42 (t, 3H), 1.63 (m, 4H), 1.76 (m, 2H), 1.88 (m, 2H), 2.34 (s, 3H), 3.88 (p, 1H), 4.12 (q, 2H), 4.82 (s, 2H), 6.94 (s, 1H), 7.38 (d, 1H), 7.55 (d, 1H), 7.72 (d, 1H), 7.91 (d, 1H), 6 (d, 1H), 9.78 (br, 1H), and 11.90 (br, 1H).

Example 271

2-{2-[3-(2-Cyclopentanecarbonyl-4-methyl-phenyl)-ureido]-thiazol-4-ylmethanesulfonyl}-1H-imidazole-4-carboxylic acid




2-{2-[3-(2-Cyclopentanecarbonyl-4-methyl-phenyl)-ureido]-thiazol-4-ylmethanesulfonyl}-1H-imidazole-4-carboxylic acid (43 mg, 83%) was prepared from 2-{2-[3-(2-cyclopentanecarbonyl-4-methyl-phenyl)-ureido]-thiazol-4-ylmethanesulfonyl}-1H-imidazole-4-carboxylic acid ethyl ester (52 mg, 0.10 mmol) following the general procedure J.

LC-MS (*m/z*): 518 (*M*+1)⁺; ¹H NMR (400 MHz, DMSO-*d*₆): δ 1.63 (m, 4H), 1.74 (m, 2H), 1.86 (m, 2H), 2.34 (s, 3H), 3.85 (p, 1H), 4.82 (s, 2H), 6.92 (s, 1H), 7.38 (d, 1H), 7.54 (d, 1H), 7.72 (d, 1H), 7.90 (s, 1H), 7.96 (br, 1H), 8.22 (d, 1H), 10.58 (br, 1H), and 11.88 (br, 1H).

Example 272

4-Amino-2-{2-[3-(2-cyclopentanecarbonyl-4-methyl-phenyl)-ureido]-thiazol-4-ylmethylsulfanyl}-pyrimidine-5-carboxylic acid ethyl ester




4-Amino-2-{2-[3-(2-cyclopentanecarbonyl-4-methyl-phenyl)-ureido]-thiazol-4-ylmethylsulfanyl}-pyrimidine-5-carboxylic acid ethyl ester (113 mg, 84%) was prepared from 1-(4-chloromethyl-thiazol-2-yl)-3-(2-cyclopentanecarbonyl-4-methyl-phenyl)-urea (94 mg, 0.25 mmol) following the general procedure Q.

LC-MS (m/z): 541 ($M+1$)⁺; ¹H NMR (400 MHz, CDCl₃): δ 1.34 (t, 3H), 1.72 (m, 4H), 1.88 (m, 4H), 2.34 (s, 3H), 3.62 (br, 2H), 3.74 (p, 1H), 4.10 (q, 2H), 4.36 (s, 2H), 6.90 (s, 1H), 7.36 (d, 1H), 7.74 (d, 1H), 8.18 (dd, 1H), 8.56 (s, 1H), 10.12 (br, 1H), and 11.78 (br, 1H).

Example 273

4-Amino-2-{2-[3-(2-cyclopentanecarbonyl-4-methyl-phenyl)-ureido]-thiazol-4-ylmethylsulfanyl}-pyrimidine-5-carboxylic acid




4-Amino-2-{2-[3-(2-cyclopentanecarbonyl-4-methyl-phenyl)-ureido]-thiazol-4-ylmethylsulfanylmethyl}-pyrimidine-5-carboxylic acid (94 mg, 91%) was prepared from 4-amino-2-{2-[3-(2-cyclopentanecarbonyl-4-methyl-phenyl)-ureido]-thiazol-4-ylmethylsulfanylmethyl}-pyrimidine-5-carboxylic acid ethyl ester (108 mg, 0.20 mmol) following the general procedure J.

LC-MS (m/z): 513 ($M+1$)⁺; ¹H NMR (400 MHz, DMSO- d_6): δ 1.72 (m, 4H), 1.88 (m, 4H), 2.34 (s, 3H), 3.66 (br, 2H), 3.86 (p, 1H), 4.36 (s, 2H), 7.07 (s, 1H), 7.32 (d, 1H), 7.72 (d, 1H), 8.20 (dd, 1H), 8.56 (s, 1H), 9.76 (br, 1H), 10.64 (br, 1H), and 11.74 (br, 1H).

10

Example 274

2-{2-[3-(2-Cyclopentanecarbonyl-4-methyl-phenyl)-ureido]-thiazol-4-ylmethylsulfanylmethyl}-4-hydroxy-pyrimidine-5-carboxylic acid




2-{2-[3-(2-Cyclopentanecarbonyl-4-methyl-phenyl)-ureido]-thiazol-4-ylmethylsulfanyl}-4-hydroxy-pyrimidine-5-carboxylic acid (104 mg, 81%) was prepared from 2-{2-[3-(2-cyclopentanecarbonyl-4-methyl-phenyl)-ureido]-thiazol-4-ylmethylsulfanyl}-4-hydroxy-pyrimidine-5-carboxylic acid ethyl ester (135 mg, 0.25 mmol) following the general procedure J.

LC-MS (m/z): 514 ($M+1$)⁺; ¹H NMR (400 MHz, CD₃OD): δ 1.74 (m, 4H), 1.90 (m, 4H), 2.36 (s, 3H), 3.84 (p, 1H), 4.28 (s, 2H), 6.98 (s, 1H), 7.36 (d, 1H), 7.70 (d, 1H), 8.36 (dd, 1H), 8.56 (s, 1H), 9.04 (br, 1H), 9.76 (br, 1H), 10.64 (br, 1H), and 11.74 (br, 1H).

10

Example 275

1-[4-(5-Amino-thiazol-2-ylsulfanylmethyl)-thiazol-2-yl]-3-(2-cyclopentanecarbonyl-4-methyl-phenyl)-urea




1-[4-(5-Amino-thiazol-2-ylsulfanylmethyl)-thiazol-2-yl]-3-(2-cyclopentanecarbonyl-4-methyl-phenyl)-urea (96 mg, 81%) was prepared from 1-(4-chloromethyl-thiazol-2-yl)-3-(2-cyclopentanecarbonyl-4-methyl-phenyl)-urea (94 mg, 0.25 mmol) following the general procedure Q.

5 LC-MS (m/z): 474 ($M+1$)⁺; ¹H NMR (400 MHz, CDCl₃): δ 1.68 (m, 4H), 1.84 (m, 4H), 2.37 (s, 3H), 3.72 (p, 1H), 4.34 (s, 2H), 5.18 (br, 2H), 6.58 (s, 1H), 7.16 (s, 1H), 7.36 (d, 1H), 7.64 (s, 1H), 8.26 (d, 1H), 10.56 (br, 1H), and 11.60 (br, 1H).

Example 276

10 {2-[3-(2-Cyclopentanecarbonyl-5-methanesulfonylamino-4-methyl-phenyl)-ureido]-thiazol-4-yl}-acetic acid ethyl ester




{2-[3-(2-Cyclopentanecarbonyl-5-methanesulfonylamino-4-methyl-phenyl)-ureido]-thiazol-4-yl}-acetic acid ethyl ester (102 mg, 80 %) was prepared from N-(5-amino-4-cyclopentanecarbonyl-2-methyl-phenyl)-methanesulfonamide (198 mg, 0.5 mmol) and ethyl-2-amino-4-thiazolyl acetate (93 mg, 0.5 mmol) following the general procedure D.

15 LC-MS (m/z): 509 ($M+1$)⁺; ¹H NMR (400 MHz, CDCl₃): δ 1.27 (t, 3H), 1.70 (m, 4H), 1.88 (m, 4H), 2.29 (s, 3H), 3.30 (s, 2H), 3.68 (s, 3H), 4.19 (q, 2H), 4.24 (p, 1H), 6.48 (s, 1H), 6.72 (s, 1H), 7.74 (s, 1H), 8.64 (br, 1H), 9.62 (br, 1H), and 11.82 (br, 1H).

20

Example 277

{2-[3-(2-Cyclopentanecarbonyl-5-methanesulfonylamino-4-methyl-phenyl)-ureido]-thiazol-4-yl}-acetic acid




{2-[3-(2-Cyclopentanecarbonyl-5-methanesulfonylamino-4-methyl-phenyl)-ureido]-thiazol-4-yl}-acetic acid ethyl ester (116 mg, 97%) was prepared from {2-[3-(2-cyclopentanecarbonyl-5-methanesulfonylamino-4-methyl-phenyl)-ureido]-thiazol-4-yl}-acetic acid ethyl ester (127 mg, 0.25 mmol) following the general procedure J.

LC-MS (m/z): 481 ($M+1$)⁺; ¹H NMR (400 MHz, DMSO- d_6): δ 1.68 (m, 4H), 1.84 (m, 4H), 2.38 (s, 3H), 3.03 (s, 3H), 3.64 (s, 2H), 4.24 (p, 1H), 5.36 (br, 1H), 6.50 (s, 1H), 6.78 (s, 1H), 7.68 (s, 1H), 9.62 (br, 1H), 11.24 (br, 1H), and 12.54 (br, 1H).

10 Example 278

1-(2-Cyclopentanecarbonyl-4-methyl-phenyl)-3-{4-[2-(1H-imidazol-2-ylsulfanyl)-ethyl]-thiazol-2-yl}-urea




1-(2-Cyclopentanecarbonyl-4-methyl-phenyl)-3-{4-[2-(1H-imidazol-2-ylsulfanyl)-ethyl]-thiazol-2-yl}-urea (25mg, 50%) was prepared from methanesulfonic acid 2-{2-[3-(2-cyclopentanecarbonyl-4-methyl-phenyl)-ureido]-thiazol-4-yl}-ethyl ester (0.05g, 0.11mmol) and 1H-imidazol-2-thiol (0.009g, 0.11mmol) following the general procedure Z.

LC/MS (m/z): 456 ($M+1$)⁺; ¹H NMR (400 MHz, CDCl₃) δ 1.63-1.68 (m, 4H), 1.83-1.88 (m, 4H), 2.35 (s, 3H), 2.87-2.93 (m, 2H), 3.25-3.28 (m, 2H), 3.60-3.75 (m, 1H), 6.74 (s, 1H), 6.85-7.10 (m, 2H), 7.33 (d, 1H), 7.65 (s, 1H), 8.4 (d, 1H), 11.35 (br, 1H).

Example 279

1-(2-Cyclopentanecarbonyl-4-methyl-phenyl)-3-{4-[2-(1-methyl-1H-imidazol-2-ylsulfanyl)-ethyl]-thiazol-2-yl}-urea




1-(2-Cyclopentanecarbonyl-4-methyl-phenyl)-3-{4-[2-(1-methyl-1H-imidazol-2-ylsulfanyl)-ethyl]-thiazol-2-yl}-urea (45mg, 43%) was prepared from methanesulfonic acid 2-{2-[3-(2-cyclopentanecarbonyl-4-methyl-phenyl)-ureido]-thiazol-4-yl}-ethyl ester (0.10g, 0.22mmol) and 1-methyl-1H-imidazole-2-thiol (0.04g, 0.33mmol) following the general procedure Z.

LC/MS (m/z): 470 (M+1)⁺; ¹H NMR (400 MHz, CDCl₃) δ 1.73-1.76 (m, 4H), 1.90-1.93 (m, 4H), 2.36 (s, 3H), 3.04 (t, 2H), 3.35 (t, 2H), 3.63 (s, 3H), 3.65-3.80 (m, 1H), 6.51 (s, 1H), 6.71 (d, 2H), 7.33 (d, 1H), 7.68 (s, 1H), 8.40 (br, 1H), 11.43 (br, 1H).

Example 280

1-(2-Cyclopentanecarbonyl-4-methyl-phenyl)-3-{4-[2-(pyridin-2-ylsulfanyl)-ethyl]-thiazol-2-yl}-urea




1-(2-cyclopentanecarbonyl-4-methyl-phenyl)-3-{4-[2-(pyridin-2-ylsulfanyl)-ethyl]-thiazol-2-yl}-urea (24mg, 467%) was prepared from methanesulfonic acid 2-{2-[3-(2-cyclopentanecarbonyl-4-methyl-phenyl)-ureido]-thiazol-4-yl}-ethyl ester (0.05g, 0.11mmol) and 2-mercapto pyridine (0.012g, 0.11mmol) following the general procedure Z.

LC/MS (m/z): 467 (M+1)⁺; ¹H NMR (400 MHz, CDCl₃) δ 1.62-1.74 (m, 4H), 1.85-1.93 (m, 4H), 2.36 (s, 3H), 3.05 (t, 2H), 3.50 (t, 2H), 3.72-3.80 (m, 1H), 6.54 (s, 1H), 7.13-7.10 (m, 2H), 7.60-7.65 (m, 3H), 8.46-8.49 (m, 2H), 11.72 (br, 1H).

Example 281

1-(2-Cyclopentanecarbonyl-4-methyl-phenyl)-3-{4-[2-(pyridin-4-ylsulfanyl)-ethyl]-thiazol-2-yl}-urea




5

1-(2-Cyclopentanecarbonyl-4-methyl-phenyl)-3-{4-[2-(pyridin-4-ylsulfanyl)-ethyl]-thiazol-2-yl}-urea (74mg, 60%) was prepared from methanesulfonic acid 2-{2-[3-(2-cyclopentanecarbonyl-4-methyl-phenyl)-ureido]-thiazol-4-yl}-ethyl ester (0.12g, 0.27mmol) and 2-mercaptopyridine (0.04g, 0.39mmol) following the general procedure Z.

10 LC/MS (m/z): 467 (M+1)⁺; ¹H NMR (400 MHz, CDCl₃) δ 1.69-1.73 (m, 4H), 1.87-1.92 (m, 4H), 2.37 (s, 3H), 3.03 (t, 2H), 3.34 (t, 2H), 3.75-3.79 (m, 1H), 6.58 (s, 1H), 7.16 (d, 2H), 7.25 (d, 2H), 7.36 (d, 1H), 7.71 (s, 1H), 8.46 (d, 1H), 11.74 (s, 1H).

Example 282

15 3-{2-[3-(2-Acetyl-4-methyl-phenyl)-ureido]-thiazol-4-ylmethylsulfanyl}-propionic acid



3-(2-Amino-thiazol-4-ylmethylsulfanyl)-propionic acid methyl ester (0.75g, 60%) was prepared from 4-chloromethyl-thiazol-2-ylamine (1.00g, 5.41mmol) and 3-mercapto-propionic acid methyl ester (0.97g, 8.11mmol) according to procedure Z.

20 3-{2-[3-(2-Acetyl-4-methyl-phenyl)-ureido]-thiazol-4-ylmethylsulfanyl}-propionic acid methyl ester (0.28g, 61%) was prepared from 3-(2-amino-thiazol-4-ylmethylsulfanyl)-


propionic acid methyl ester (0.23g, 0.99mmol) and (2-amino-5-methyl-phenyl)-cyclopentyl-methanone (0.24g, 1.49mmol) following the procedure D.

3-{2-[3-(2-Acetyl-4-methyl-phenyl)-ureido]-thiazol-4-ylmethylsulfanyl}-propionic acid (40mg, 83%) was prepared from [4-(2-{2-[3-(2-cyclopentanecarbonyl-4-methyl-phenyl)-ureido]-thiazol-4-yl} ethoxy)-phenyl] acetic acid methyl ester (0.05g, 0.11mmol) and 2.5M LiOH (20uL) following the general procedure J.

LC/MS (*m/z*): 448 (*M*+1)⁺ ; ¹H NMR (400 MHz, DMSO) δ 1.70-1.80 (m, 4H), 1.85-1.97 (m, 4H), 2.28 (s, 3H), 2.67 (t, 2H), 2.83 (t, 2H), 3.70 (s, 2H), 3.85-3.90 (m, 1H), 6.89 (s, 1H), 7.20 (d, 2H), 7.85 (s, 1H), 8.20 (d, 1H), 10.64 (br, 1H), 10.73 (br, 1H), 12.14 (br, 1H).

Example 283

1-(2-Cyclopentanecarbonyl-4-methyl-phenyl)-3-(5-methylsulfanyl-thiazol-2-yl)-urea




1-(2-Cyclopentanecarbonyl-4-methyl-phenyl)-3-(5-methylsulfanyl-thiazol-2-yl)-urea (94 mg, 25%) was prepared from 1-(5-bromo-thiazol-2-yl)-3-(2-cyclopentanecarbonyl-4-methyl-phenyl)-urea (408 mg, 1 mmol) and sodium methanethiolate (140 mg, 2 mmol) following the general procedure Q.

LC-MS (*m/z*): 376 (*M*+1)⁺; ¹H NMR (400 MHz, CDCl₃): δ 1.74 (m, 4H), 1.93 (m, 4H) 2.38 (s, 3H), 2.43 (s, 3H), 3.79 (m, 1H), 7.37 (d, 1H), 7.73 (s, 2H), 8.45 (d, 1H), 11.65 (br, 1H), 12.20 (br, 1H).

Example 284

{2-[3-(2-Cyclopentanecarbonyl-4-methyl-phenyl)-ureido]-thiazol-5-ylsulfanyl}-acetic acid methyl ester




{2-[3-(2-Cyclopentanecarbonyl-4-methyl-phenyl)-ureido]-thiazol-5-ylsulfanyl}-acetic acid methyl ester (130 mg, 30%) was prepared from 1-(5-bromo-thiazol-2-yl)-3-(2-cyclopentanecarbonyl-4-methyl-phenyl)-urea (408 mg, 1 mmol), and mercapto-acetic acid methyl ester (212 mg, 2 mmol) following the general procedure Q.

LC-MS (m/z): 434 ($M+1$)⁺; ¹H NMR (400 MHz, CDCl₃): δ 1.71 (m, 4H), 1.91 (m, 4H), 2.38 (s, 3H), 3.47 (s, 3H), 3.72 (s, 3H), 3.79 (m, 1H), 7.39 (d, 1H), 7.75 (d, 2H), 8.43 (d, 1H), 11.40 (br, 1H), 11.66 (br, 1H).

10

Example 285

{2-[3-(2-Cyclopentanecarbonyl-4-methyl-phenyl)-ureido]-thiazol-5-ylsulfanyl}-acetic acid




{2-[3-(2-Cyclopentanecarbonyl-4-methyl-phenyl)-ureido]-thiazol-5-ylsulfanyl}-acetic acid (161 mg, 75 %) was prepared from {2-[3-(2-cyclopentanecarbonyl-4-methyl-phenyl)-ureido]-thiazol-5-ylsulfanyl}-acetic acid methyl ester (229 mg, 0.5 mmol) following the general procedure J.

LC-MS (m/z): 420 ($M+1$)⁺; ¹H NMR (400 MHz, DMSO-*d*₆): δ 1.61 (m, 4H), 1.74 (m, 2H), 1.89 (m, 2H), 2.33 (s, 3H), 3.49 (s, 3H), 3.88 (m, 1H), 7.41 (d, 2H), 7.92 (d, 1H), 8.33 (d, 1H), 10.80 (br, 1H), 10.98 (br, 1H), 12.12 (br, 1H).

20

Example 286

{2-[3-(2-Cyclopentanecarbonyl-4-methyl-phenyl)-ureido]-thiazole-5-sulfonyl}-acetic acid methyl ester




{2-[3-(2-Cyclopentanecarbonyl-4-methyl-phenyl)-ureido]-thiazole-5-sulfonyl}-acetic acid methyl ester (175 mg, 75%) was prepared from {2-[3-(2-cyclopentanecarbonyl-4-methyl-phenyl)-ureido]-thiazole-5-sulfonyl}-acetic acid methyl ester (217 mg, 0.5 mmol) by oxidation with *m*-cpba following the general procedure R.

LC-MS (*m/z*): 466 (*M*+1)⁺; ¹H NMR (400 MHz, CDCl₃): δ 1.71 (m, 4H), 1.91 (m, 4H), 2.38 (s, 3H), 3.78 (m, 4H), 4.24 (s, 2H), 7.41 (d, 1H), 7.77 (s, 1H), 8.30 (s, 1H), 8.42 (d, 1H), 11.89 (br, 1H).

10

Example 287

{2-[3-(2-Cyclopentanecarbonyl-4-methyl-phenyl)-ureido]-thiazole-5-sulfonyl}-acetic acid



15


{2-[3-(2-Cyclopentanecarbonyl-4-methyl-phenyl)-ureido]-thiazol-5-ylsulfonyl}-acetic acid (158 mg, 70 %) was prepared from {2-[3-(2-cyclopentanecarbonyl-4-methyl-phenyl)-ureido]-thiazol-5-ylsulfonyl}-acetic acid methyl ester (233 mg, 0.5 mmol) following the general procedure J.

20

LC-MS (*m/z*): 452 (*M*+1)⁺; ¹H NMR (400 MHz, acetone-*d*₆): δ 1.67 (m, 4H), 1.71 (m, 4H), 1.91 (m, 4H), 2.38 (s, 3H), 3.95 (m, 1H), 4.44 (s, 2H), 7.44 (d, 1H), 7.97 (s, 2H), 8.42 (d, 1H), 11.15 (br, 1H), 11.35 (br, 1H).

Example 288

3-{2-[3-(2-Cyclopentanecarbonyl-4-methyl-phenyl)-ureido]-thiazol-5-ylsulfonyl}-propionic acid methyl ester




{2-[3-(2-Cyclopentanecarbonyl-4-methyl-phenyl)-ureido]-thiazol-5-ylsulfanyl}-propionic acid methyl ester (125 mg, 28 %) was prepared from 1-(5-bromo-thiazol-2-yl)-3-(2-cyclopentanecarbonyl-4-methyl-phenyl)-urea (408 mg, 1 mmol), and mercapto-propionic acid methyl ester (240 mg, 2 mmol) following the general procedure Q.

LC-MS (m/z): 448 ($M+1$)⁺; ¹H NMR (400 MHz, CDCl₃): δ 1.71 (m, 4H), 1.91 (m, 4H), 2.38 (s, 3H), 2.65 (s, 2H), 3.51 (s, 2H), 3.66 (s, 3H), 3.78 (m, 1H), 6.90 (d, 1H), 7.38 (d, 1H), 7.73 (s, 1H), 8.42 (d, 1H) 11.62 (br, 1H), 11.65 (br, 1H).

10 Example 289

3-{2-[3-(2-Cyclopentanecarbonyl-4-methyl-phenyl)-ureido]-thiazol-5-ylsulfanyl}-propionic acid




{2-[3-(2-Cyclopentanecarbonyl-4-methyl-phenyl)-ureido]-thiazol-5-ylsulfanyl}-propionic acid (138 mg, 72 %) was prepared from {2-[3-(2-cyclopentanecarbonyl-4-methyl-phenyl)-ureido]-thiazol-5-ylsulfanyl}-propionic acid methyl ester (217 mg, 0.5 mmol) following the general procedure J.

LC-MS (m/z): 433 ($M+1$)⁺; ¹H NMR (400 MHz, CDCl₃): δ 1.67 (m, 4H), 1.89 (m, 4H), 2.36 (s, 3H), 2.51 (s, 2H), 3.59 (s, 2H), 3.78 (m, 1H), 6.90 (d, 1H), 7.41 (d, 1H), 7.93 (s, 1H), 8.45 (d, 1H), 9.56 (br, 1H), 11.16 (br, 1H), 11.25 (br, 1H).

Example 290

3-{2-[3-(2-Cyclopentanecarbonyl-4-methyl-phenyl)-ureido]-thiazole-5-sulfonyl}-propionic acid methyl ester




5 {2-[3-(2-Cyclopentanecarbonyl-4-methyl-phenyl)-ureido]-thiazole-5-sulfonyl}-propionic acid methyl ester (163 mg, 68%) was prepared from {2-[3-(2-cyclopentanecarbonyl-4-methyl-phenyl)-ureido]-thiazole-5-sulfonyl}-propionic acid methyl ester (224 mg, 0.5 mmol) by oxidation with *m*-cpba following the general procedure R.

LC-MS (*m/z*): 480 (*M*+1)⁺; ¹H NMR (400 MHz, CDCl₃): δ 1.71 (m, 6H), 1.91 (m, 2H),
 10 2.40 (s, 3H), 2.85 (m, 2H), 3.56 (m, 2H), 3.67 (s, 3H), 3.81 (m, 1H), 7.41 (dd, 1H), 7.78 (s, 2H), 8.29 (s, 1H), 8.42 (d, 1H), 11.40 (br, 1H), 11.91 (br, 1H).

Example 291

15 3-{2-[3-(2-Cyclopentanecarbonyl-4-methyl-phenyl)-ureido]-thiazole-5-sulfonyl}-propionic acid




20 {2-[3-(2-Cyclopentanecarbonyl-4-methyl-phenyl)-ureido]-thiazol-5-ylsulfonyl}-propionic acid (168 mg, 72 %) was prepared from {2-[3-(2-cyclopentanecarbonyl-4-methyl-phenyl)-ureido]-thiazol-5-ylsulfonyl}-propionic acid methyl ester (240 mg, 0.5 mmol) following the general procedure J.

LC-MS (m/z): 466 ($M+1$)⁺; ¹H NMR (400 MHz, CDCl₃): δ 1.71 (m, 4H), 1.91 (m, 4H), 2.38 (s, 3H), 2.89 (s, 2H), 3.63 (s, 2H), 3.81 (m, 1H), 7.39 (d, 1H), 7.75 (d, 2H), 8.43 (d, 1H), 11.40 (br, 1H), 11.66 (br, 1H).

5 EXAMPLE 292

N-(2-{2-[3-(2-Cyclopentanecarbonyl-4-methyl-phenyl)-ureido]-thiazol-5-yl-sulfanyl}-ethyl)-acetamide




N-(2-{2-[3-(2-Cyclopentanecarbonyl-4-methyl-phenyl)-ureido]-thiazol-5-yl-sulfanyl}-ethyl)-acetamide (156 mg, 35 %) was prepared from 1-(5-bromo-thiazol-2-yl)-3-(2-cyclopentanecarbonyl-4-methyl-phenyl)-urea (408 mg, 1 mmol), and N-(2-mercapto-ethyl)-acetamide (240 mg, 2 mmol) following the general procedure Q.

LC-MS (m/z): 447 ($M+1$)⁺; ¹H NMR (400 MHz, CDCl₃): δ 1.71 (m, 4H), 1.91 (m, 4H), 1.99 (s, 3H), 2.38 (s, 3H), 2.85 (t, 3H), 3.46 (q, 2H), 3.79 (m, 1H), 5.95 (br, 1H), 7.37 (d, 1H), 7.69 (d, 2H), 8.42 (d, 1H), 11.35 (br, 1H), 11.57 (br, 1H).

Example 293

N-(2-{2-[3-(2-Cyclopentanecarbonyl-4-methyl-phenyl)-ureido]-thiazol-5-yl-sulfonyl}-ethyl)-acetamide




N-(2-{2-[3-(2-Cyclopentanecarbonyl-4-methyl-phenyl)-ureido]-thiazol-5-yl-sulfonyl}-ethyl)-acetamide (167 mg, 70%) was prepared from N-(2-{2-[3-(2-cyclopentanecarbonyl-4-methyl-phenyl)-ureido]-thiazol-5-yl-sulfonyl}-ethyl)-acetamide (223 mg, 0.5 mmol) by oxidation with *m*-cpba following the general procedure R.

- 5 LC-MS (*m/z*): 479 (M+1)⁺; ¹H NMR (400 MHz, CDCl₃): δ 1.70 (m, 4H), 1.92 (m, 4H), 2.01 (s, 3H), 2.39 (s, 3H), 3.42 (m, 2H), 3.76 (m, 3H), 6.32 (m, 1H), 7.39 (d, 1H), 7.76 (s, 2H), 8.13 (s, 1H), 8.40 (d, 1H), 11.66 (br, 1H).

Example 294

- 10 2-Acetylamino-3-{2-[3-(2-cyclopentanecarbonyl-4-methyl-phenyl)-ureido]-thiazol-5-ylsulfanyl}-propionic acid methyl ester




- 15 2-Acetylamino-3-{2-[3-(2-cyclopentanecarbonyl-4-methyl-phenyl)-ureido]-thiazol-5-ylsulfanyl}-propionic acid methyl ester (156 mg, 35 %) was prepared from 1-(5-bromo-thiazol-2-yl)-3-(2-cyclopentanecarbonyl-4-methyl-phenyl)-urea (408 mg, 1 mmol) and 2-acetylamino-3-mercapto-propionic acid methyl ester (354 mg, 2 mmol) following the general procedure Q.

- 20 LC-MS (*m/z*): 505 (M+1)⁺; ¹H NMR (400 MHz, CDCl₃): δ 1.71 (m, 6H), 1.91 (m, 2H), 2.03 (s, 3H), 2.38 (s, 3H), 3.24 (m, 2H), 3.66 (s, 3H), 3.79 (m, 1H), 4.84 (m, 1H), 6.42 (d, 1H), 7.39, d, 1H), 7.60 (s, 1H), 7.73 (s, 1H), 8.43 (d, 1H), 10.88 (br, 1H), 11.64 (br, 1H).

Example 295

2-Acetylamino-3-{2-[3-(2-cyclopentanecarbonyl-4-methyl-phenyl)-ureido]-thiazol-5-ylsulfanyl}-propionic acid




- 2-Acetylamino-3-{2-[3-(2-cyclopentanecarbonyl-4-methyl-phenyl)-ureido]-thiazol-5-ylsulfanyl}-propionic acid (187 mg, 76 %) was prepared from 2-acetylamino-3-{2-[3-(2-cyclopentanecarbonyl-4-methyl-phenyl)-ureido]-thiazol-5-ylsulfanyl}-propionic acid methyl ester (245 mg, 0.5 mmol) following the general procedure J.

LC-MS (m/z): 491 ($M+1$)⁺; ¹H NMR (400 MHz, Acetone- d_6): δ 1.68 (m, 4H), 1.84 (m, 4H), 1.97 (s, 3H), 2.37 (s, 3H), 3.10 (m, 1H), 3.26 (m, 1H), 3.94 (m, 1H), 4.64 (m, 1H), 5.8 (br, 1H), 7.45 (d, 1H), 7.54 (d, 1H), 8.01 (s, 1H), 8.43 (d, 1H), 11.20 (br, 1H), 11.44 (br, 1H).

10 Example 296

2-tert-Butoxycarbonylamino-3-{2-[3-(2-cyclopentanecarbonyl-4-methyl-phenyl)-ureido]thiazol-5-ylsulfanyl}-propionic acid methyl ester




- 2-tert-Butoxycarbonylamino-3-{2-[3-(2-cyclopentanecarbonyl-4-methyl-phenyl)-ureido]thiazol-5-ylsulfanyl}-propionic acid methyl ester (196 mg, 35 %) was prepared from 1-(5-bromo-thiazol-2-yl)-3-(2-cyclopentanecarbonyl-4-methyl-phenyl)-urea (408 mg, 1 mmol) and 2-tert-butoxycarbonylamino-3-mercapto-propionic acid methyl ester (470 mg, 2 mmol) following the general procedure Q.

LC-MS (m/z): 563 ($M+1$)⁺; ¹H NMR (400 MHz, CDCl₃): δ 1.55 (s, 9H), 1.71 (m, 4H), 1.91 (m, 4H), 2.37 (s, 3H), 3.18 (m, 2H), 3.76 (m, 4H), 4.53 (m, 1H), 5.45 (m, 1H), 7.35 (d, 1H), 7.65 (s, 1H), 7.72 (s, 1H), 8.43 (d, 1H), 11.60 (br, 1H), 11.66 (br, 1H).

5

Example 297

(2-{2-[3-(2-Cyclopentanecarbonyl-4-methyl-phenyl)-ureido]-thiazol-5-ylsulfanyl}-ethyl)-carbamic acid tert-butyl ester




(2-{2-[3-(2-Cyclopentanecarbonyl-4-methyl-phenyl)-ureido]-thiazol-5-ylsulfanyl}-ethyl)-carbamic acid tert-butyl ester (202 mg, 40 %) was prepared from 1-(5-bromo-thiazol-2-yl)-3-(2-cyclopentanecarbonyl-4-methyl-phenyl)-urea (408 mg, 1 mmol) and (2-mercapto-ethyl)-carbamic acid tert-butyl ester (354 mg, 2 mmol) following the general procedure Q.

LC-MS (m/z): 505 ($M+1$)⁺; ¹H NMR (400 MHz, CDCl₃): δ 1.45(s, 9H), 1.71 (m, 4H), 1.92 (m, 4H), 2.38 (s, 3H), 2.83 (m, 2H), 3.32 (m, 2H), 3.78 (m, 1H), 4.95 (br, 1H), 7.36 (d, 1H), 7.73 (s, 2H), 8.43 (d, 1H), 11.48 (br, 1H), 11.62 (br, 1H).

15

Example 298

2-{2-[3-(2-Cyclopentanecarbonyl-4-methyl-phenyl)-ureido]-thiazol-5-ylsulfanyl}-ethyl-ammonium; chloride




20

2-{2-[3-(2-Cyclopentanecarbonyl-4-methyl-phenyl)-ureido]-thiazol-5-ylsulfanyl}-ethyl-ammonium; chloride (183 mg, 90%) was prepared by reacting (2-{2-[3-(2-cyclopentane-carbonyl-4-methyl-phenyl)-ureido]-thiazol-5-ylsulfanyl}-ethyl)-carbamic acid tert-butyl ester (253 mg, 0.5 mmol) with 4 N HCl in dioxane

5 LC-MS (m/z): 406 ($M+1$)⁺; ¹H NMR (400 MHz, DMSO- d_6): δ 1.63 (m, 4 H), 1.73 (m, 2H), 1.91 (m, 2H), 2.33 (s, 3H), 2.91 (m, 2 H), 3.18 (m, 2H), 3.87 (m, 1H), 7.39 (d, 1H), 7.56 (s, 2H), 7.85 (s, 1H), 8.16 (d, 1H), 10.71 (br, 1H), 12.20 (br, 1H).

Example 299

10 (2-{2-[3-(2-Cyclopentanecarbonyl-4-methyl-phenyl)-ureido]-thiazole-5-sulfonyl}-ethyl) carbamic acid tert-butyl ester




15 (2-{2-[3-(2-Cyclopentanecarbonyl-4-methyl-phenyl)-ureido]-thiazole-5-sulfonyl}-ethyl)-carbamic acid tert-butyl ester (195 mg, 73%) was prepared from (2-{2-[3-(2-cyclopentane-carbonyl-4-methyl-phenyl)-ureido]-thiazole-5-sulfanyl}-ethyl) carbamic acid tert-butyl ester (252 mg, 0.5 mmol) by oxidation with *m*-cpba following the general procedure R.

LC-MS (m/z): 537 ($M+1$)⁺; ¹H NMR (400 MHz, CDCl₃): δ 1.44 (s, 9H), 1.71 (m, 4H), 1.91 (m, 4H), 2.38 (s, 3H), 3.46 (t, 2H), 3.78 (m, 3H), 3.81 (m, 1H), 7.39 (d, 1H), 7.75 (d, 2H), 8.43 (d, 1H), 11.40 (br, 1H), 11.66 (br, 1H).

20

Example 300

2-{2-[3-(2-Cyclopentanecarbonyl-4-methyl-phenyl)-ureido]-thiazole-5-sulfonyl}-ethyl ammonium; chloride




2-{2-[3-(2-Cyclopentanecarbonyl-4-methyl-phenyl)-ureido]-thiazol-5-ylsulfonyl}-ethyl-ammonium; chloride (185 mg, 85%) was prepared by reacting 2-{2-[3-(2-cyclopentanecarbonyl-4-methyl-phenyl)-ureido]-thiazol-5-ylsulfonyl}-ethyl-carbamic acid tert-butyl ester (253 mg, 0,5 mmol) with 4 N HCl in dioxane

LC-MS (*m/z*): 438 (*M*+1)⁺; ¹H NMR (400 MHz, DMSO-*d*₆): δ 1.61(m, 4H), 1.72 (m, 2H), 1.88 (m, 2H), 2.34 (s, 3H), 3.01(br, 2H), 3.70 (t, 2H), 3.88 (m, 1H), 7.41 (d, 1H), 7.87 (s, 1H), 8.13 (d, 1H), 8.15 (br, 1H), 10.75 (br, 1H), 11.62 (br, 1H).

10

Example 301

1-(2-Cyclopentanecarbonyl-4-methyl-phenyl)-3-[5-(2-dimethylamino-ethylsulfanyl)-thiazol-2-yl]-urea




1-(2-Cyclopentanecarbonyl-4-methyl-phenyl)-3-[5-(2-dimethylamino-ethylsulfanyl)-thiazol-2-yl]-urea (130 mg, 30 %) was prepared from 1-(5-bromo-thiazol-2-yl)-3-(2-cyclopentanecarbonyl-4-methyl-phenyl)-urea (408 mg, 1 mmol) and 2-dimethylamino-ethanethiol (210 mg, 2 mmol) following the general procedure Q.

LC-MS (*m/z*): 433 (*M*+1)⁺; ¹H NMR (400 MHz, CDCl₃): δ 1.71 (m, 4H), 1.92 (m, 4H), 2.24 (s, 6H), 2.38 (s, 3H), 2.55(m, 2H), 2.83 (m, 2H), 3.78 (m, 1H), 7.36 (d, 1H), 7.69 (s, 1H), 7.74(s, 1H), 8.44 (d, 1H), 11.65 (br, 1H).

20

Example 302

1-(2-Cyclopentanecarbonyl-4-methyl-phenyl)-3-[5-(2-hydroxy-ethylsulfanyl)-thiazol-2-yl]-urea




1-(2-Cyclopentanecarbonyl-4-methyl-phenyl)-3-[5-(2-hydroxy-ethylsulfanyl)-thiazol-2-yl]-urea (114 mg, 28 %) was prepared from 1-(5-bromo-thiazol-2-yl)-3-(2-cyclopentanecarbonyl-4-methyl-phenyl)-urea (408 mg, 1 mmol) and 2-hydroxy-ethanethiol (158 mg, 2 mmol) following the general procedure Q.

LC-MS (m/z): 406 ($M+1$)⁺; ¹H NMR (400 MHz, CDCl₃/acetone- d_6): δ 1.52 (m, 4H), 1.68 (m, 2H), 1.75 (m, 2H), 2.33 (s, 3H), 2.71 (t, 3H), 3.58 (m, 2H), 3.90 (m, 1H), 7.17 (d, 1H), 7.31 (s, 1H), 7.56 (s, 1H), 8.27 (d, 1H), 11.20 (br, 1H).

Example 303

1-(2-Cyclopentanecarbonyl-4-methyl-phenyl)-3-[5-(2-hydroxy-ethylsulfonyl)-thiazol-2-yl]-urea




1-(2-Cyclopentanecarbonyl-4-methyl-phenyl)-3-[5-(2-hydroxy-ethylsulfonyl)-thiazol-2-yl]-urea (153, 70%) was prepared from 1-(2-cyclopentanecarbonyl-4-methyl-phenyl)-3-[5-(2-hydroxy-ethylsulfanyl)-thiazol-2-yl]-urea (204 mg, 0.5 mmol) by oxidation with *m*-cpba following the general procedure R.

LC-MS (m/z): 438 ($M+1$)⁺; ¹H NMR (400 MHz, CDCl₃/acetone- d_6): δ 1.66 (m, 4H), 1.89 (m, 4H), 2.37 (s, 3H), 3.48 (m, 2H), 3.72 (m, 2H), 4.07 (m, 2H), 7.37 (d, 1H), 7.54 (d, 1H), 7.66 (d, 1H), 8.01 (d, 1H), 11.88 (br, 1H).

Example 304

1-(2-Cyclopentanecarbonyl-4-methyl-phenyl)-3-[5-(pyridin-2-ylsulfanyl)-thiazol-2-yl]-urea




5 1-(2-Cyclopentanecarbonyl-4-methyl-phenyl)-3-[5-(pyridin-2-ylsulfanyl)-thiazol-2-yl]-urea (140 mg, 32 %) was prepared from 1-(5-bromo-thiazol-2-yl)-3-(2-cyclopentanecarbonyl-4-methyl-phenyl)-urea (408 mg, 1 mmol) and 2-mercaptopyridine (222 mg, 2 mmol) following the general procedure Q.

LC-MS (*m/z*): 439 (*M*+1)⁺; ¹H NMR (400 MHz, CDCl₃): δ 1.63 (m, 4H), 1.86 (m, 4H),
 10 2.37 (s, 3H), 3.76 (m, 1H), 7.04 (m, 2H), 7.37 (d, 1H), 7.51 (m, 1H), 7.73 (s, 1H), 7.92 (s, 1H),
 8.42 (d, 1H), 8.48 (d, 1H), 11.16 (br, 1H), 11.81 (br, 1H).

Example 305

15 1-(2-Cyclopentanecarbonyl-4-methyl-phenyl)-3-[5-(pyridine-2-sulfonyl)-thiazol-2-yl]-urea




1-(2-Cyclopentanecarbonyl-4-methyl-phenyl)-3-[5-(pyridine-2-sulfonyl)-thiazol-2-yl]-urea (176 mg, 75%) was prepared from 1-(2-cyclopentanecarbonyl-4-methyl-phenyl)-3-[5-(pyridine-2-sulfanyl)-thiazol-2-yl]-urea (219 mg, 0.5 mmol) by oxidation with *m*-cpba following the general procedure R.

20 LC-MS (*m/z*): 471 (*M*+1)⁺; ¹H NMR (400 MHz, CDCl₃): δ 1.68 (m, 4H), 1.93 (m, 4H),
 2.39 (s, 3H), 3.79 (m, 1H), 7.39 (d, 1H), 7.48 (m, 1H), 7.76 (s, 1H), 7.93 (dd, 1H), 8.15 (d,
 1H), 8.33 (br, 1H), 8.42 (d, 1H), 8.70 (d, 1H), 10.8 (br, 1H), 11.93 (br, 1H).

Example 306

1-(2-Cyclopentanecarbonyl-4-methyl-phenyl)-3-[5-(pyrimidin-2-ylsulfanyl)-thiazol-2-yl]-urea



5

1-(2-Cyclopentanecarbonyl-4-methyl-phenyl)-3-[5-(pyrimidin-2-ylsulfanyl)-thiazol-2-yl]-urea (142 mg, 32 %) was prepared from 1-(5-bromo-thiazol-2-yl)-3-(2-cyclopentanecarbonyl-4-methyl-phenyl)-urea (408 mg, 1 mmol) and 2-mercaptopyrimidine (222 mg, 2 mmol) following the general procedure Q.


10

LC-MS (m/z): 440 ($M+1$)⁺; ¹H NMR (400 MHz, CDCl₃): δ 1.64 (m, 4H), 1.89 (m, 4H), 2.36 (s, 3H), 3.81 (m, 1H), 7.01 (t, 1H), 7.34 (d, 2H), 7.72 (s, 1H), 7.87 (s, 1H), 8.47 (d, 1H), 8.58 (d, 1H), 8.67 (d, 1H), 11.70 (br, 1H), 11.78 (br, 1H).

Example 307

15

1-(2-Cyclopentanecarbonyl-4-methyl-phenyl)-3-[5-(pyrimidin-2-ylsulfonyl)-thiazol-2-yl]-urea



20


1-(2-Cyclopentanecarbonyl-4-methyl-phenyl)-3-[5-(pyrimidine-2-sulfonyl)-thiazol-2-yl]-urea (169 mg, 72%) was prepared from 1-(2-cyclopentanecarbonyl-4-methyl-phenyl)-3-[5-(pyrimidine-2-sulfanyl)-thiazol-2-yl]-urea (220 mg, 0.5 mmol) by oxidation with *m*-cpba following the general procedure R.

LC-MS (m/z): 472 ($M+1$)⁺; ¹H NMR (400 MHz, CDCl₃): δ 1.71 (m, 4H), 1.91 (m, 4H), 2.38 (s, 3H), 3.78 (m, 1H), 7.39 (d, 1H), 7.49 (t, 1H), 7.76 (s, 1H), 8.31 (s, 1H), 8.43 (d, 1H), 8.93 (d, 2H), 11.80 (br, 1H), 11.94 (br, 1H).

5

Example 308

1-(2-Cyclopentanecarbonyl-4-methyl-phenyl)-3-[5-(pyridin-4-ylsulfanyl)-thiazol-2-yl]-
urea




10 1-(2-Cyclopentanecarbonyl-4-methyl-phenyl)-3-[5-(pyridin-4-ylsulfanyl)-thiazol-2-yl]-
urea (110 mg, 25 %) was prepared from 1-(5-bromo-thiazol-2-yl)-3-(2-cyclopentanecarbonyl-
4-methyl-phenyl)-urea (408 mg, 1 mmol) and 4-mercaptopyridine (222 mg, 2 mmol) following
the general procedure Q.

LC-MS (m/z): 439 ($M+1$)⁺; ¹H NMR (400 MHz, CDCl₃): δ 1.62 (m, 4H), 1.84 (m, 4H),
2.35 (s, 3H), 3.73 (m, 1H), 7.10 (d, 2H), 7.34 (d, 2H), 7.71 (s, 1H), 7.95 (s, 1H), 8.38 (d, 1H),
15 8.42 (d, 1H), 11.65 (br, 1H), 11.75 (br, 1H).

Example 309

1-(2-Cyclopentanecarbonyl-4-methyl-phenyl)-3-[5-(1-methyl-1H-imidazol-2-
ylsulfanyl)-thiazol-2-yl]-urea



20


1-(2-Cyclopentanecarbonyl-4-methyl-phenyl)-3-[5-(1-methyl-1H-imidazol-2-
ylsulfanyl)-thiazol-2-yl]-urea (141 mg, 32 %) was prepared from 1-(5-bromo-thiazol-2-yl)-3-(2-
cyclopentanecarbonyl-4-methyl-phenyl)-urea (408 mg, 1 mmol) and 1-methyl-1H-imidazole-
2-thiol (228 mg, 2 mmol) following the general procedure Q.

LC-MS (*m/z*): 442 (*M*+1)⁺; ¹H NMR (400 MHz, CDCl₃): δ 1.66 (m, 4H), 1.88 (m, 4H), 2.35 (s, 3H), 3.74 (m, 1H), 3.77 (s, 3H), 6.93 (s, 1H), 7.06 (s, 1H), 7.32 (d, 1H), 7.69 (s, 1H), 7.82 (s, 1H), 8.40 (d, 1H), 11.51 (br, 1H), 12.20 (br, 1H).

5

Example 310

1-(2-Cyclopentanecarbonyl-4-methyl-phenyl)-3-[5-(1H-imidazol-2-ylsulfanyl)-thiazol-2-yl]-urea




10 1-(2-Cyclopentanecarbonyl-4-methyl-phenyl)-3-[5-(1H-imidazol-2-ylsulfanyl)-thiazol-2-yl]-urea (149 mg, 35 %) was prepared from 1-(5-bromo-thiazol-2-yl)-3-(2-cyclopentanecarbonyl-4-methyl-phenyl)-urea (408 mg, 1 mmol) and 1H-imidazole-2-thiol (200 mg, 2 mmol) following the general procedure Q.

LC-MS (*m/z*): 428 (*M*+1)⁺; ¹H NMR (400 MHz, CDCl₃): δ 1.70 (m, 4H), 1.91 (m, 4H), 2.37 (s, 3H), 3.83 (m, 1H), 7.10 (s, 1H), 7.34 (d, 1H), 7.57 (s, 2H), 7.66 (s, 1H), 7.78 (s, 1H), 8.42 (d, 1H), 11.22 (br, 1H).

15

Example 311

1-(2-Cyclopentanecarbonyl-4-methyl-phenyl)-3-[5-(7H-purin-6-ylsulfanyl)-thiazol-2-yl]-urea



20


1-(2-Cyclopentanecarbonyl-4-methyl-phenyl)-3-[5-(7H-purin-6-ylsulfanyl)-thiazol-2-yl]-urea (124 mg, 28%) was prepared from 1-(5-bromo-thiazol-2-yl)-3-(2-cyclopentanecarbonyl-4-methyl-phenyl)-urea (408 mg, 1 mmol) and 7H-purine-6-thiol (304 mg, 2 mmol) following the general procedure Q.

LC-MS (m/z): 480 ($M+1$)⁺; ¹H NMR (400 MHz, CDCl₃): δ 1.61 (m, 4H), 1.81 (m, 4H), 2.26 (s, 3H), 3.64 (m, 1H), 7.21 (d, 2H), 7.54 (d, 2H), 8.02 (s, 1H), 8.26 (d, 1H), 8.56 (s, 1H), 11.12 (br, 1H), 11.62 (br, 1H).

5

Example 312

2-{2-[3-(2-Cyclopentanecarbonyl-4-methyl-phenyl)-ureido]-thiazol-5-ylsulfanyl}-
nicotinic acid methyl ester




2-{2-[3-(2-Cyclopentanecarbonyl-4-methyl-phenyl)-ureido]-thiazol-5-ylsulfanyl}-
10 nicotinic acid methyl ester (124 mg, 25%) was prepared from 1-(5-bromo-thiazol-2-yl)-3-(2-
cyclopentanecarbonyl-4-methyl-phenyl)-urea (408 mg, 1 mmol) and 2-mercapto-nicotinic
acid methyl ester (340 mg, 2 mmol) following the general procedure Q.

LC-MS (m/z): 497 ($M+1$)⁺; ¹H NMR (400 MHz, CDCl₃): δ 1.64 (m, 4H), 1.85 (m, 4H),
2.36 (s, 3H), 3.74 (m, 1H), 3.98 (s, 3H), 7.08 (dd, 1H), 7.34 (d, 2H), 7.71 (s, 1H), 7.79 (s 1H),
15 8.22 (dd, 1H), 8.45 (d, 1H), 8.49 (dd, 1H), 11.45 (br, 1H), 11.75 (br, 1H).

Example 313

2-{2-[3-(2-Cyclopentanecarbonyl-4-methyl-phenyl)-ureido]-thiazol-5-ylsulfanyl}-
nicotinic acid



20


2-{2-[3-(2-Cyclopentanecarbonyl-4-methyl-phenyl)-ureido]-thiazol-5-ylsulfanyl}-
nicotinic acid (186 mg, 75 %) was prepared from 2-{2-[3-(2-cyclopentanecarbonyl-4-methyl-

phenyl)-ureido]-thiazol-5-ylsulfanyl]-nicotinic acid methyl ester (248 mg, 0.5 mmol) following the general procedure J.

LC-MS (m/z): 483 ($M+1$)⁺; ¹H NMR (400 MHz, DMSO- d_6): δ 1.62 (m, 4H), 1.75 (m, 2H), 1.89 (m, 2H), 2.33 (s, 3H), 3.88 (m, 1H), 7.23 (dd, 1H), 7.39 (d, 1H), 7.47 (s, 1H), 7.86 (s, 1H), 8.19 (m, 2H), 8.24 (dd, 1H), 10.70 (br, 1H), 12.23 (br, 1H)

Example 314

2-{2-[3-(2-Cyclopentanecarbonyl-4-methyl-phenyl)-ureido]-thiazol-5-ylsulfonyl]-nicotinic acid methyl ester



10

2-{2-[3-(2-Cyclopentanecarbonyl-4-methyl-phenyl)-ureido]-thiazol-5-ylsulfonyl]-nicotinic acid methyl ester (185 mg, 70%) was prepared from 2-{2-[3-(2-cyclopentanecarbonyl-4-methyl-phenyl)-ureido]-thiazol-5-ylsulfanyl]-nicotinic acid methyl ester (248 mg, 0.5 mmol) by oxidation with *m*-cpba following the general procedure R.


LC-MS (m/z): 529 ($M+1$)⁺, ¹H NMR (400 MHz, CDCl₃): δ 1.68 (m, 4H), 1.85 (m, 2H), 1.95 (m, 2H), 2.38 (s, 3H), 3.95 (m, 1H), 3.99 (s, 3H), 7.42 (d, 1H), 7.79 (dd, 2H), 7.97 (s, 1H), 8.14 (s, 1H), 8.15 (d, 1H), 8.82 (d, 1H), 8.79 (d, 1H), 11.35 (br, 1H), 11.68 (br, 1H).

15

Example 315

2-{2-[3-(2-Cyclopentanecarbonyl-4-methyl-phenyl)-ureido]-thiazol-5-ylsulfonyl]-nicotinic acid

20




2-{2-[3-(2-Cyclopentanecarbonyl-4-methyl-phenyl)-ureido]-thiazol-5-ylsulfonyl}-nicotinic acid (200 mg, 78 %) was prepared from 2-{2-[3-(2-cyclopentanecarbonyl-4-methyl-phenyl)-ureido]-thiazol-5-ylsulfonyl}-nicotinic acid methyl ester (264 mg, 0.5 mmol) following the general procedure J.

5 LC-MS (*m/z*): 515 (*M*+1)⁺; ¹H NMR (400 MHz, acetone-*d*₆): δ 1.67 (m, 4H), 1.84 (m, 2H), 1.95 (m, 2H), 2.38 (s, 3H), 3.95 (m, 1H), 7.44 (d, 1H), 7.76 (dd, 1H), 7.96 (s, 1H), 8.05 (s, 1H), 8.18 (d, 1H), 8.42 (d, 1H), 8.77 (d, 1H), 11.23 (br, 1H), 11.36 (br, 1H).

Example 316

10 6-{2-[3-(2-Cyclopentanecarbonyl-4-methyl-phenyl)-ureido]-thiazol-5-ylsulfanyl}-nicotinic acid methyl ester




15 6-{2-[3-(2-Cyclopentanecarbonyl-4-methyl-phenyl)-ureido]-thiazol-5-ylsulfanyl}-nicotinic acid methyl ester (100 mg, 20%) was prepared from 1-(5-bromo-thiazol-2-yl)-3-(2-cyclopentanecarbonyl-4-methyl-phenyl)-urea (408 mg, 1 mmol) and 6-mercapto-nicotinic acid methyl ester (340 mg, 2 mmol) following the general procedure Q.

LC-MS (*m/z*): 497 (*M*+1)⁺; ¹H NMR (400 MHz, CDCl₃): δ 1.64 (m, 4H), 1.85 (m, 4H), 2.33 (s, 3H), 3.79 (m, 1H), 3.99 (s, 3H), 7.10 (dd, 1H), 7.34 (dd, 1H), 7.71 (s, 1H), 7.79 (s, 1H), 8.22 (dd, 1H), 8.45 (d, 1H), 8.50 (dd, 1H), 8.79 (s, 1H), 11.50 (br, 1H), 11.75 (br, 1H).

20 Example 317

6-{2-[3-(2-Cyclopentanecarbonyl-4-methyl-phenyl)-ureido]-thiazol-5-ylsulfanyl}-nicotinic acid




6-{2-[3-(2-Cyclopentanecarbonyl-4-methyl-phenyl)-ureido]-thiazol-5-ylsulfanyl}-
 nicotinic acid (169 mg, 70 %) was prepared from 6-{2-[3-(2-cyclopentanecarbonyl-4-methyl-
 phenyl)-ureido]-thiazol-5-ylsulfanyl}-nicotinic acid methyl ester (248 mg, 0.5 mmol) following
 5 the general procedure J.

LC-MS (m/z): 483 ($M+1$)⁺; ¹H NMR (400 MHz, DMSO- d_6): δ 1.62 (m, 4H), 1.75 (m, 2H), 1.89 (m, 2H), 2.33 (s, 3H), 3.89 (m, 1H), 7.12 (d, 1H), 7.38 (d, 1H), 7.75 (s, 1H), 7.87 (s, 1H), 8.12 (dd, 2H), 8.24 (dd, 1H), 8.85 (s, 1H), 10.76 (br, 1H), 12.60 (br, 1H).

10 Example 318

6-{2-[3-(2-Cyclopentanecarbonyl-4-methyl-phenyl)-ureido]-thiazol-5-ylsulfonyl}-
 nicotinic acid methyl ester




6-{2-[3-(2-Cyclopentanecarbonyl-4-methyl-phenyl)-ureido]-thiazol-5-ylsulfonyl}-
 15 nicotinic acid methyl ester (172 mg, 65%) was prepared from 6-{2-[3-(2-cyclopentane-
 carbonyl-4-methyl-phenyl)-ureido]-thiazol-5-ylsulfanyl}-nicotinic acid methyl ester (248 mg,
 0.5 mmol) by oxidation with *m*-cpba following the general procedure R.

LC-MS (m/z): 529 ($M+1$)⁺; ¹H NMR (400 MHz, CDCl₃): δ 1.74 (m, 4H), 1.97 (m, 4H),
 2.38 (s, 3H), 3.81 (m, 1H), 3.98 (s, 3H), 7.44 (t, 1H), 7.58 (d, 1H), 8.02 (dd, 1H), 8.11 (s, 1
 20 H), 8.23 (d, 1H), 8.52 (dd, 1H), 9.26 (d, 1H), 10.56 (br, 1H), 12.11 (br, 1H).

Example 319

3-{2-[3-(2-Cyclopentanecarbonyl-4-methyl-phenyl)-ureido]-thiazol-5-ylsulfanyl}-pyridine-2-carboxylic acid methyl ester



5


3-{2-[3-(2-Cyclopentanecarbonyl-4-methyl-phenyl)-ureido]-thiazol-5-ylsulfanyl}-pyridine-2-carboxylic acid methyl ester (149 mg, 30%) was prepared from 1-(5-bromothiiazol-2-yl)-3-(2-cyclopentanecarbonyl-4-methyl-phenyl)-urea (408 mg, 1 mmol) and 3-mercapto-pyridine-2-carboxylic acid methyl ester (340 mg, 2 mmol) following the general procedure Q.

10

LC-MS (m/z): 497 ($M+1$)⁺; ¹H NMR (400 MHz, CDCl₃): δ 1.59 (m, 4H), 1.80 (m, 2H), 1.91 (m, 2H), 2.37 (s, 3H), 3.73 (m, 1H), 4.01 (s, 3H), 7.29 (dd, 1H), 7.36 (d, 1H), 7.52 (dd, 1H), 7.72 (s, 1H), 7.97 (s, 1H), 8.11 (dd, 1H), 8.46 (m, 1H), 11.83 (br, 1H).

15 EXAMPLE 320

3-{2-[3-(2-Cyclopentanecarbonyl-4-methyl-phenyl)-ureido]-thiazol-5-ylsulfanyl}-pyridine-2-carboxylic acid



20


3-{2-[3-(2-Cyclopentanecarbonyl-4-methyl-phenyl)-ureido]-thiazol-5-ylsulfanyl}-pyridine-2-carboxylic acid (181 mg, 75 %) was prepared from 3-{2-[3-(2-Cyclopentanecarbonyl-4-methyl-phenyl)-ureido]-thiazol-5-ylsulfanyl}-pyridine-2-carboxylic acid methyl ester (248 mg, 0.5 mmol) following the general procedure J.

LC-MS (m/z): 483 ($M+1$)⁺; ¹H NMR (400 MHz, DMSO- d_6): δ 1.62 (m, 4H), 1.75 (m, 2H), 1.91 (m, 2H), 2.32 (s, 3H), 3.88 (m, 1H), 7.38 (d, 2H), 7.51 (dd, 1H), 7.72 (s, 1H), 7.87 (s, 1H), 8.13 (d, 1H), 8.44 (d, 1H), 10.74 (br, 1H), 12.25 (br, 1H).

5

Example 321

2-[2-[3-(2-Cyclopentanecarbonyl-4-methyl-phenyl)-ureido]-thiazol-5-ylsulfanyl]-1H-imidazole-4-carboxylic acid ethyl ester




2-[2-[3-(2-Cyclopentanecarbonyl-4-methyl-phenyl)-ureido]-thiazol-5-ylsulfanyl]-1H-imidazole-4-carboxylic acid ethyl ester (190 mg, 38%) was prepared from 1-(5-bromo-thiazol-2-yl)-3-(2-cyclopentanecarbonyl-4-methyl-phenyl)-urea (408 mg, 1 mmol) and 2-mercapto-1H-imidazole-4-carboxylic acid ethyl ester (340 mg, 2 mmol) following the general procedure Q.

LC-MS (m/z): 500 ($M+1$)⁺; ¹H NMR (400 MHz, CDCl₃): δ 1.27 (t, 3H), 1.62 (m, 4H), 1.72 (m, 2H), 1.89 (m, 2H), 2.34 (s, 3H), 3.87 (m, 1H), 4.16 (q, 2H), 6.60 (br, 1H), 7.39 (d, 1H), 7.70 (s, 1H), 7.82 (s, 2H), 8.15 (d, 1H), 10.22 (br, 1H), 11.90 (br, 1H).

Example 322

2-[2-[3-(2-Cyclopentanecarbonyl-4-methyl-phenyl)-ureido]-thiazol-5-ylsulfanyl]-1H-imidazole-4-carboxylic acid

20




2-{2-[3-(2-Cyclopentanecarbonyl-4-methyl-phenyl)-ureido]-thiazol-5-ylsulfany]-1H-imidazole-4-carboxylic acid (167 mg, 71 %) was prepared from 2-{2-[3-(2-cyclopentanecarbonyl-4-methyl-phenyl)-ureido]-thiazol-5-ylsulfany]-1H-imidazole-4-carboxylic acid ethyl ester (250 mg, 0.5 mmol) following the general procedure J.

LC-MS (m/z): 472 ($M+1$)⁺; ¹H NMR (400 MHz, DMSO- d_6): δ 1.61 (m, 4H), 1.73 (m, 2H), 1.89 (m, 2H), 2.33 (s, 3H), 3.87 (m, 1H), 6.53 (br, 1H), 7.37 (d, 1H), 7.64 (s, 1H), 7.83 (s, 2H), 8.14 (d, 1H), 10.70 (br, 1H), 12.20 (br, 1H).

10 Example 323

2-{2-[3-(2-Cyclopentanecarbonyl-4-methyl-phenyl)-ureido]-thiazol-5-ylsulfonyl]-1H-imidazole-4-carboxylic acid ethyl ester




2-{2-[3-(2-Cyclopentanecarbonyl-4-methyl-phenyl)-ureido]-thiazol-5-ylsulfonyl]-1H-imidazole-4-carboxylic acid ethyl ester (181 mg, 68%) was prepared from 2-{2-[3-(2-cyclopentanecarbonyl-4-methyl-phenyl)-ureido]-thiazol-5-ylsulfonyl]-1H-imidazole-4-carboxylic acid ethyl ester (250 mg, 0.5 mmol) by oxidation with *m*-cpba following the general procedure R.

LC-MS (m/z): 532 ($M+1$)⁺; ¹H NMR (400 MHz, CDCl₃): δ 1.27 (t, 3H), 1.71 (m, 4H), 1.91 (m, 4H), 2.36 (s, 3H), 3.73 (m, 1H), 4.38 (q, 2H), 6.98 (s, 1H), 7.34 (d, 1H), 7.71 (s, 1H), 7.83 (s, 1H), 8.22 (br, 1H), 8.38 (d, 1H), 11.80 (br, 1H).

5

Example 324

2-{2-[3-(2-Cyclopentanecarbonyl-4-methyl-phenyl)-ureido]-thiazol-5-ylsulfonyl}-1H-imidazole-4-carboxylic acid




2-{2-[3-(2-Cyclopentanecarbonyl-4-methyl-phenyl)-ureido]-thiazol-5-ylsulfonyl}-1H-imidazole-4-carboxylic acid (171 mg, 68 %) was prepared from 2-{2-[3-(2-cyclopentanecarbonyl-4-methyl-phenyl)-ureido]-thiazol-5-ylsulfonyl}-1H-imidazole-4-carboxylic acid ethyl ester (266 mg, 0.5 mmol) following the general procedure J.

LC-MS (m/z): 504 ($M+1$)⁺; ¹H NMR (400 MHz, DMSO- d_6): δ 1.61 (m, 4H), 1.75 (m, 2H), 1.89 (m, 2H), 2.33 (s, 3H), 3.88 (m, 1H), 7.41 (d, 1H), 7.53 (dd, 2H), 7.69 (dd, 1H), 7.88 (m, 1H), 8.01 (br, 1H), 8.14 (d, 1H), 10.86 (br, 1H).

15

Example 325

(3-Cyclopentanecarbonyl-4-{3-[5-(pyridin-2-ylsulfanyl)-thiazol-2-yl]-ureido}-phenyl)-acetic acid ethyl ester




20

(3-Cyclopentanecarbonyl-4-{3-[5-(pyridin-2-ylsulfanyl)-thiazol-2-yl]-ureido}-phenyl)-acetic acid ethyl ester (153 mg, 30%) was prepared from {4-[3-(5-bromo-thiazol-2-yl)-ureido]-3-cyclopentanecarbonyl-phenyl}-acetic acid ethyl ester (480 mg, 1 mmol) and 2-mercapto-pyridine (222 mg, 2 mmol) following the general procedure Q.

5 LC-MS (m/z): 511 ($M+1$)⁺; ¹H NMR (400 MHz, CDCl₃): δ 1.26 (t, 3H), 1.61 (m, 4H), 1.85 (m, 4H), 3.62 (s, 2H), 3.76 (m, 1H), 4.16 (q, 2H), 7.01 (dd, 1H), 7.06 (d, 1H), 7.45 (d, 1H), 7.51 (d, 1H), 7.89 (s, 1H), 7.93 (s, 1H), 8.41 (dd, 1H), 8.55 (d, 1H), 11.70 (br, 1H), 11.89 (br, 1H).

10 Example 326

(3-Cyclopentanecarbonyl-4-{3-[5-(pyridin-2-ylsulfanyl)-thiazol-2-yl]-ureido}-phenyl)-acetic acid




15 (3-Cyclopentanecarbonyl-4-{3-[5-(pyridin-2-ylsulfanyl)-thiazol-2-yl]-ureido}-phenyl)-acetic acid (173 mg, 72 %) was prepared from (3-cyclopentanecarbonyl-4-{3-[5-(pyridin-2-ylsulfanyl)-thiazol-2-yl]-ureido}-phenyl)-acetic acid ethyl ester (255 mg, 0.5 mmol) following the general procedure J.

20 LC-MS (m/z): 483 ($M+1$)⁺; ¹H NMR (400 MHz, Acetone d₆): δ 1.64 (m, 4H), 1.86 (m, 2H), 1.98 (m, 4H), 3.71 (s, 2H), 3.94 (m, 1H), 7.08 (d, 1H), 7.15 (dd, 1H), 7.53 (d, 1H), 7.61 (s, 1H), 7.68 (m, 1H), 8.11 (s, 1H), 8.39 (m, 1H), 8.52 (d, 1H), 11.30 (br, 1H), 11.35 (br, 1H).

Example 327

25 (3-Cyclopentanecarbonyl-4-{3-[5-(1H-imidazol-2-ylsulfanyl)-thiazol-2-yl]-ureido}-phenyl)-acetic acid ethyl ester




(3-Cyclopentanecarbonyl-4-{3-[5-(1H-imidazol-2-ylsulfanyl)-thiazol-2-yl]-ureido}-phenyl)acetic acid ethyl ester (120 mg, 24%) was prepared from {4-[3-(5-bromo-thiazol-2-yl)-ureido]-3-cyclopentanecarbonyl-phenyl}-acetic acid ethyl ester (480 mg, 1 mmol) and 1H-imidazole-2-thiol (200 mg, 2 mmol) following the general procedure Q.

LC-MS (m/z): 500 ($M+1$)⁺; ¹H NMR (400 MHz, CDCl₃): δ 1.21 (t, 3H), 1.58 (m, 4H), 1.83 (m, 4H), 3.61 (m, 3H), 4.17 (q, 1H), 6.73 (s, 1H), 7.04 (m, 2H), 7.38 (d, 1H), 7.61 (br, 1H), 7.81 (s, 1H), 8.41 (d, 1H), 11.23 (br, 1H), 11.49 (br, 1H).

10 Example 328

(3-Cyclopentanecarbonyl-4-{3-[5-(1H-imidazol-2-ylsulfanyl)-thiazol-2-yl]-ureido}-phenyl)-acetic acid




(3-Cyclopentanecarbonyl-4-{3-[5-(1H-imidazol-2-ylsulfanyl)-thiazol-2-yl]-ureido}-phenyl)-acetic acid (177 mg, 72 %) was prepared from (3-cyclopentanecarbonyl-4-{3-[5-(1H-imidazol-2-ylsulfanyl)-thiazol-2-yl]-ureido}-phenyl)-acetic acid ethyl ester (250 mg, 0.5 mmol) following the general procedure J.

LC-MS (m/z): 472 ($M+1$)⁺; ¹H NMR (400 MHz, acetone d_6): δ 1.61 (m, 4H), 1.91 (m, 4H), 3.58 (s, 2H), 3.72 (m, 1H), 6.77 (s, 1H), 7.04 (d, 1H), 7.33 m, 2H), 7.64 (br, 1H), 7.80 (s, 1H), 8.43 (d, 1H), 11.44 (br, 1H), 11.59 (br, 1H).

Example 329

1-(2-Cyclopentanecarbonyl-4-methyl-phenyl)-3-[5-(1-methyl-1H-tetrazol-5-ylsulfanyl)-thiazol-2-yl]-urea




1-(2-Cyclopentanecarbonyl-4-methyl-phenyl)-3-[5-(1-methyl-1H-tetrazol-5-ylsulfanyl)-thiazol-2-yl]-urea (133 mg, 30 %) was prepared from 1-(5-bromo-thiazol-2-yl)-3-(2-cyclopentanecarbonyl-4-methyl-phenyl)-urea (408 mg, 1 mmol) and 1-methyl-1H-tetrazole-5-thiol (202 mg, 2 mmol) following the general procedure Y

LC-MS (*m/z*): 444 (*M*+1)⁺, ¹H NMR (400 MHz, CDCl₃): δ 1.70 (m, 4H), 1.91 (m, 4H), 2.37 (s, 3H), 3.78 (m, 1H), 4.06 (s, 3H), 7.37 (d, 1H), 7.73 (s, 1H), 7.93 (s, 1H), 8.41 (d, 1H), 9.30 (br, 1H), 11.74 (br, 1H).

10 Example 330

(5-{2-[3-(2-Cyclopentanecarbonyl-4-methyl-phenyl)-ureido]-thiazol-5-ylsulfanyl}-tetrazol-1-yl)-acetic acid




(5-{2-[3-(2-Cyclopentanecarbonyl-4-methyl-phenyl)-ureido]-thiazol-5-ylsulfanyl}-tetrazol-1-yl)-acetic acid (136 mg, 28 %) was prepared from 1-(5-bromo-thiazol-2-yl)-3-(2-cyclopentanecarbonyl-4-methyl-phenyl)-urea (408 mg, 1 mmol) and (5-mercapto-tetrazol-1-yl)-acetic acid (320 mg, 2 mmol) following the general procedure Y.

LC-MS (*m/z*): 488 (*M*+1)⁺, ¹H NMR (400 MHz, DMSO-*d*₆): δ 1.70 (m, 4H), 1.91 (m, 4H), 2.34 (s, 3H), 3.89 (m, 1H), 5.48 (s, 2H), 7.42 (d, 1H), 7.81 (s, 1H), 7.88 (s, 1H), 8.17 (d, 1H), 10.76 (br, 1H), 12.25 (br, 2H).

Example 331

1-(2-Cyclopentanecarbonyl-4-methyl-phenyl)-3-{5-[1-(2-dimethylamino-ethyl)-1H-tetrazol-5-ylsulfanyl]-thiazol-2-yl}-urea




1-(2-Cyclopentanecarbonyl-4-methyl-phenyl)-3-{5-[1-(2-dimethylamino-ethyl)-1H-tetrazol-5-ylsulfanyl]-thiazol-2-yl}-urea (125 mg, 25 %) was prepared from 1-(5-bromothiazol-2-yl)-3-(2-cyclopentanecarbonyl-4-methyl-phenyl)-urea (408 mg, 1 mmol) and 1-(2-dimethylamino-ethyl)-1H-tetrazole-5-thiol (346 mg, 2 mmol) following the general procedure Y

LC-MS (m/z): 501 ($M+1$)⁺, ¹H NMR (400 MHz, DMSO- d_6): δ 1.70 (m, 4H), 1.91 (m, 4H), 2.28 (s, 6H), 2.34 (s, 3H), 2.76 (t, 2H), 3.74 (m, 1H), 4.42 (t, 2H), 7.32 (d, 1H), 7.68 (s, 1H), 7.83 (s, 1H), 8.36 (d, 1H), 11.25 (br, 1H), 11.40 (br, 1H).

Example 332

1-(2-Cyclopentanecarbonyl-4-methyl-phenyl)-3-[5-(4H-[1,2,4]triazol-3-ylsulfanyl)-thiazol-2-yl]-urea




1-(2-Cyclopentanecarbonyl-4-methyl-phenyl)-3-[5-(4H-[1,2,4]triazol-3-ylsulfanyl)-thiazol-2-yl]-urea (129 mg, 30 %) was prepared from 1-(5-bromothiazol-2-yl)-3-(2-cyclopentanecarbonyl-4-methyl-phenyl)-urea (408 mg, 1 mmol) and 2H-[1,2,4]triazole-3-thiol (202 mg, 2 mmol) following the general procedure Y

LC-MS (m/z): 429 ($M+1$)⁺, ¹H NMR (400 MHz, DMSO- d_6): δ 1.63 (m, 4H), 1.70 (m, 2H), 1.91 (m, 2H), 2.34 (s, 3H), 3.89 (m, 1H), 7.41 (d, 1H), 7.63 (s, 1H), 7.87 (s, 1H), 8.17 (d, 1H), 10.72 (br, 1H), 12.18 (br, 1H).

Example 333

1-(2-Cyclopentanecarbonyl-4-methyl-phenyl)-3-[5-(4-methyl-4H-[1,2,4]triazol-3-ylsulfanyl)-thiazol-2-yl]-urea




5 1-(2-Cyclopentanecarbonyl-4-methyl-phenyl)-3-[5-(4-methyl-4H-[1,2,4]triazol-3-ylsulfanyl)-thiazol-2-yl]-urea (125 mg, 25 %) was prepared from 1-(5-bromo-thiazol-2-yl)-3-(2-cyclopentanecarbonyl-4-methyl-phenyl)-urea (408 mg, 1 mmol) and 4-methyl-4H-[1,2,4]triazole-3-thiol (230mg, 2 mmol) following the general procedure Y

LC-MS (m/z): 443 ($M+1$)⁺, ¹H NMR (400 MHz, DMSO- d_6): δ 1.63 (m, 4H), 1.75 (m, 2H), 1.91 (m, 1H), 2.34 (s, 3H), 3.70 (s, 3H), 3.89 (m, 1H), 7.41 (d, 1H), 7.73 (s, 1H), 7.87 (s, 1H), 8.16 (d, 1H), 8.62 (s, 1H), 10.72 (br, 1H), 12.15 (br, 1H).

Example 334

1-(2-Cyclopentanecarbonyl-4-methyl-phenyl)-3-[5-(4-methyl-4H-[1,2,4]triazole-3-sulfonyl)-thiazol-2-yl]-urea




15 1-(2-Cyclopentanecarbonyl-4-methyl-phenyl)-3-[5-(4-methyl-4H-[1,2,4]triazole-3-sulfonyl)-thiazol-2-yl]-urea (171 mg, 72%) was prepared from 1-(2-cyclopentanecarbonyl-4-methyl-phenyl)-3-[5-(4-methyl-4H-[1,2,4]triazole-3-sulfanyl)-thiazol-2-yl]-urea (221 mg, 0.5 mmol) by oxidation with *m*-cpba following the general procedure R

20 LC-MS (m/z): 475 ($M+1$)⁺, ¹H NMR (400 MHz, DMSO- d_6): δ 1.73 (m, 4H), 1.85 (m, 2H), 1.91 (m, 2H), 2.34 (s, 3H), 3.70 (s, 3H), 3.89 (m, 1H), 7.51 (d, 1H), 7.73 (s, 1H), 7.87 (s, 1H), 8.16 (d, 1H), 8.62 (s, 1H), 10.72 (br, 1H), 12.15 (br, 1H).

Example 335

(4-{2-[3-(2-Cyclopentanecarbonyl-4-methyl-phenyl)-ureido]-thiazol-5-yl sulfanyl}-phenyl)-acetic acid



5

(4-{2-[3-(2-Cyclopentanecarbonyl-4-methyl-phenyl)-ureido]-thiazol-5-yl sulfanyl}-phenyl)-acetic acid (140 mg, 30 %) was prepared from 1-(5-bromo-thiazol-2-yl)-3-(2-cyclopentanecarbonyl-4-methyl-phenyl)-urea (408 mg, 1 mmol) and (4-mercapto-phenyl)-acetic acid (336 mg, 2 mmol) following the general procedure Y.


10

(LC-MS (m/z): 496 ($M+1$)⁺; ¹H NMR (400 MHz, DMSO- d_6): δ 1.63 (m, 4H), 1.75 (m, 2H), 1.91 (m, 2H), 2.33 (s, 3H), 3.59 (s, 2H), 3.89 (m, 1H), 7.20 (m, 3H), 7.40 (d, 2H), 7.68 (s, 1H), 7.87 (s, 1H), 8.16 (d, 1H), 12.36 (br, 3H).

Example 336

15

1-(5-Acetyl-4-methyl-thiazol-2-yl)-3-(2-cyclopentanecarbonyl-4-methyl-phenyl)-urea




20

1-(5-Acetyl-4-methyl-thiazol-2-yl)-3-(2-cyclopentanecarbonyl-4-methyl-phenyl)-urea (116 mg, 60 %) was prepared from (2-amino-5-methyl-phenyl)-cyclopentyl-methanone (102 mg, 0.5 mmol) and 1-(2-amino-4-methyl-thiazol-5-yl)-ethanone (94 mg, 0.6 mmol) following the general procedure D.

LC-MS (m/z): 386 ($M+1$)⁺, ¹H NMR (400 MHz, DMSO- d_6): δ 1.63 (m, 4H), 1.76 (m, 2H), 1.91 (m, 2H), 2.35 (s, 3H), 2.45 (s, 3H), 2.55 (s, 3H), 3.89 (m, 1H), 7.42 (d, 1H), 7.88 (s, 1H), 8.20 (s, 1H), 10.80 (br, 1H), 12.25 (br, 1H).

Example 337

1-(2-Cyclopentanecarbonyl-4-methyl-phenyl)-3-(5-nitro-thiazol-2-yl)-urea




5 1-(2-Cyclopentanecarbonyl-4-methyl-phenyl)-3-(5-nitro-thiazol-2-yl)-urea (103 mg, 55 %) was prepared from (2-amino-5-methyl-phenyl)-cyclopentyl-methanone (102 mg, 0.5 mmol) and 5-nitro-thiazol-2-ylamine (87 mg, 0.6 mmol) following the general procedure D.

LC-MS (*m/z*): 375 (*M*+1)⁺, ¹H NMR (400 MHz, DMSO-*d*₆): δ 1.64 (m, 4H), 1.76 (m, 2H), 1.91 (m, 2H), 2.36 (s, 3H), 3.91 (m, 1H), 7.44 (d, 1H), 7.91 (d, 1H), 8.21 (d, 1H), 8.58 (s, 1H), 10.97 (br, 1H), 12.25 (br, 1H).

10

Example 338

1-(4-tert-Butyl-thiazol-2-yl)-3-(2-cyclopentanecarbonyl-4-methyl-phenyl)-urea




15 1-(4-tert-Butyl-thiazol-2-yl)-3-(2-cyclopentanecarbonyl-4-methyl-phenyl)-urea (127 mg, 66 %) was prepared from (2-amino-5-methyl-phenyl)-cyclopentyl-methanone (102 mg, 0.5 mmol) and 4-tert-butyl-thiazol-2-ylamine (94 mg, 0.6 mmol) following the general procedure D.

LC-MS (*m/z*): 386 (*M*+1)⁺, ¹H NMR (400 MHz, DMSO-*d*₆): δ 1.24 (s, 9H), 1.62 (m, 4H), 1.76 (m, 2H), 1.91 (m, 2H), 2.32 (s, 3H), 3.87 (m, 1H), 6.63 (s, 1H), 7.37 (d, 1H), 7.83 (d, 1H), 8.13 (d, 1H), 10.53 (br, 1H), 11.82 (br, 1H).

20

Example 339

1-(2-Cyclopentanecarbonyl-4-methyl-phenyl)-3-pyridin-2-yl-urea




1-(2-Cyclopentanecarbonyl-4-methyl-phenyl)-3-pyridin-2-yl-urea (94 mg, 58 %) was prepared from (2-amino-5-methyl-phenyl)-cyclopentyl-methanone (102 mg, 0.5 mmol) and 2-aminopyridine (57 mg, 0.6 mmol) following the general procedure D.

LC-MS (m/z): 324($M+1$)⁺, ¹H NMR (400 MHz, DMSO- d_6): δ 1.63 (m, 4H), 1.76 (m, 2H), 1.91 (m, 2H), 2.35 (s, 3H), 3.90 (m, 1H), 6.03 (d, 1H), 7.42 (d, 1H), 7.80 (m, 4H), 8.21 (s, 1H), 10.77 (br, 1H), 12.24 (br, 1H).

10 **Example 340**

1-(2-Cyclopentanecarbonyl-4-methyl-phenyl)-3-(5-formyl-thiazol-2-yl)-urea




1-(2-Cyclopentanecarbonyl-4-methyl-phenyl)-3-(5-formyl-thiazol-2-yl)-urea (116 mg, 65 %) was prepared from (2-amino-5-methyl-phenyl)-cyclopentyl-methanone (102 mg, 0.5 mmol) and 2-amino-thiazol-5-carbaldehyde (77 mg, 0.6 mmol) following the general procedure D.

LC-MS (m/z): 358 ($M+1$)⁺; ¹H NMR (400 MHz, DMSO- d_6): δ 1.65 (m, 4H), 1.76 (m, 2H), 1.90 (m, 2H), 2.35 (s, 3H), 3.89 (m 1H), 7.43 (d, 1H), 7.89 (d, 1H), 8.19 (d, 1H), 8.36 (s, 1H), 9.89 (s, 1H), 10.85 (br, 1H), 12.5 (br, 1H).

20

Example 341

1-(2-Cyclopentanecarbonyl-4-methyl-phenyl)-3-[5-(hydroxyimino-methyl)-thiazol-2-yl]-urea




To a mixture of 2M aq NaHCO₃ (5 mL) and 1-(2-cyclopentanecarbonyl-4-methyl-phenyl)-3-(5-formyl-thiazol-2-yl)-urea (72 mg, 0.2 mmol) in THF (5 mL) was added hydroxylamine hydrochloride (200 mg) at room temperature with stirring. The resulting mixture was heated at 60 °C for 12 h. The reaction mixture was cooled and extracted with ethyl acetate. The organic layer was washed (brine), dried (Na₂SO₄) and concentrated to obtain 1-(2-cyclopentanecarbonyl-4-methyl-phenyl)-3-[5-(hydroxyimino-methyl)-thiazol-2-yl]-urea (56 mg, 76%) as a mixture of *syn* and *anti*-isomers.

LC-MS (*m/z*): 373 (M+1)⁺.

10

Example 342

{2-[3-(2-Cyclopentanecarbonyl-4-methyl-phenyl)-ureido]-thiazol-5-ylmethyleneaminoxy}-acetic acid




To a mixture of 2M aq NaHCO₃ (5 mL) and 1-(2-cyclopentanecarbonyl-4-methyl-phenyl)-3-(5-formyl-thiazol-2-yl)-urea (72 mg, 0.2 mmol) in THF (5 mL) was added hydroxylamine hydrochloride (200 mg) at room temperature with stirring. The resulting mixture was heated at 60 °C for 12 h. The reaction mixture was cooled, acidified to pH 6.5 (citric acid) and extracted with ethyl acetate. The organic layer was washed (brine), dried (Na₂SO₄) and concentrated to obtain {2-[3-(2-cyclopentanecarbonyl-4-methyl-phenyl)-ureido]-thiazol-5-ylmethyleneaminoxy}-acetic acid (56 mg, 76%) as a mixture of *syn* and *anti*-isomers.

LC-MS (*m/z*): 431 (M+1)⁺.

25

Example 343

((2-[3-(2-Cyclopentanecarbonyl-4-methyl-phenyl)-ureido]-thiazol-5-ylmethyl)-amino)-acetic acid methyl ester




((2-[3-(2-Cyclopentanecarbonyl-4-methyl-phenyl)-ureido]-thiazol-5-ylmethyl)-amino)-acetic acid methyl ester (38 mg, 45%) was prepared from 1-(2-cyclopentanecarbonyl-4-methyl-phenyl)-3-(5-formyl-thiazol-2-yl)-urea (0.072 g, 0.2 mmol) and glycine methyl ester hydrochloride following the general procedure O.

LC-MS (m/z): 431 ($M+1$)⁺; ¹H NMR (400 MHz, DMSO- d_6): δ 1.64 (m, 4H), 1.76 (m, 2H), 1.91 (m, 2H), 2.33 (s, 3H), 3.33 (s, 2H), 3.35 (br, 1H), 3.63 (s, 3H), 3.83 (s, 2H), 3.91 (m, 1H), 7.14 (s, 1H), 7.38 (d, 1H), 7.83 (s, 1H), 8.17 (s, 1H), 10.63 (br, 1H), 11.56 (br, 1H).

Example 344

3-{2-[3-(2-Cyclopentanecarbonyl-4-methyl-phenyl)-ureido]-thiazol-5-yl}-acrylic acid ethyl ester




3-{2-[3-(2-Cyclopentanecarbonyl-4-methyl-phenyl)-ureido]-thiazol-5-yl}-acrylic acid ethyl ester (96 mg, 80%) was prepared from 1-(2-cyclopentanecarbonyl-4-methyl-phenyl)-3-(5-formyl-thiazol-2-yl)-urea (100 mg, 0.28 mmol) and (carbethoxymethylene)-triphenylphosphorane (0.12 g, 0.34 mmol) following the general procedure X

LC-MS (m/z): 428 ($M+1$)⁺, ¹H NMR (400 MHz, CDCl₃): δ 1.33 (t, 3H), 1.69 (m, 4H), 1.92 (m, 4H), 2.38 (s, 3H), 3.78 (m, 1H), 4.24 (q, 2H), 6.10 (d, 1H), 7.25 (d, 1H), 7.38 (d, 1H), 7.46 (m, 2H), 7.99 (s, 1H), 8.46 (d, 1H), 10.40 (br, 1H), 11.72 (br, 1H).

Example 345

3-{2-[3-(2-Cyclopentanecarbonyl-4-methyl-phenyl)-ureido]-thiazol-5-yl}-propionic acid ethyl ester




- 5 3-{2-[3-(2-Cyclopentanecarbonyl-4-methyl-phenyl)-ureido]-thiazol-5-yl}-propionic acid ethyl ester was prepared from 3-{2-[3-(2-cyclopentanecarbonyl-4-methyl-phenyl)-ureido]-thiazol-5-yl}-acrylic acid ethyl ester via hydrogenation under the conditions described in the general procedure C

LC-MS (*m/z*): 430 (*M*+1)⁺.

10 **Example 346**

1-[2-(2-Methoxy-benzoyl)-5-methyl-phenyl]-3-thiazol-2-yl-urea




- (2-Amino-4-methyl-phenyl)-(2-methoxy-phenyl)-methanone (723 g, 30%) was prepared from 2-amino-4-methyl-benzoic acid (1.51g, 10 mmol) following the general procedure S. 1-[2-(2-methoxy-benzoyl)-5-methyl-phenyl]-3-thiazol-2-yl-urea (129 mg, 70 %) was prepared from (2-amino-4-methyl-phenyl)-(2-methoxy-phenyl)-methanone (121 mg, 0.5 mmol) and 2-aminothiazole (0.060 g, 0.6 mmol) following the general procedure D.

- LC-MS (*m/z*): 368 (*M*+1)⁺; ¹H NMR (400 MHz, CDCl₃): δ 2.42 (s, 3H), 3.75 (s, 3H), 6.78 (d, 1H), 6.91 (d, 1H), 6.98 (d, 1H), 7.04 (t, 1H), 7.26 (t, 1H), 7.33 (d, 1H), 7.46 (m, 1H), 7.64 (d, 1H), 8.46 (s, 1H), 11.02 (br, 1H), 11.67 (br, 1H).

Example 347

1-[2-(2-Methoxy-benzoyl)-4-methyl-phenyl]-3-thiazol-2-yl-urea




(2-Amino-5-methyl-phenyl)-(2-methoxy-phenyl)-methanone (843 g, 35%) was prepared from 2-amino-5-methyl-benzoic acid (1.51 g, 0.1 mol) following the general procedure S. 1-[2-(2-methoxy-benzoyl)-4-methyl-phenyl]-3-thiazol-2-yl-urea (132 mg, 72%) was prepared from (2-amino-5-methyl-phenyl)-(2-methoxy-phenyl)-methanone (121 mg, 0.5 mmol) and 2-aminothiazole (0.060 g, 0.6 mmol) following the general procedure D.

LC-MS (m/z): 368 ($M+1$)⁺; ¹H NMR (400 MHz, CDCl₃): δ 2.23 (s, 3H), 3.76 (s, 3H), 6.91 (d, 1H), 7.01 (d, 1H), 7.05 (t, 1H), 7.23 (bs, 1H), 7.29 (d, 1H), 7.36 (d, 1H), 7.48 (t, 1H), 7.59 (d, 1H), 8.48 (s, 1H), 10.56 (br, 1H), 11.45 (br, 1H).

Example 348

1-[5-Fluoro-2-(2-methoxy-benzoyl)-phenyl]-3-thiazol-2-yl- urea




(2-Amino-4-fluoro-phenyl)-(2-methoxy-phenyl)-methanone (784 mg, 32 %) was prepared from 2-amino-4-fluoro-benzoic acid (1.55 g, 0.1 mol) following the general procedure S. 1-[5-Fluoro-2-(2-methoxy-benzoyl)-phenyl]-3-thiazol-2-yl- urea (127 mg, 68 %) was prepared from (2-amino-4-fluoro-phenyl)-(2-methoxy-phenyl)-methanone (123 mg, 0.5 mmol) and 2-aminothiazole (0.060 g, 0.6 mmol) following the general procedure D.

LC-MS (m/z): 372 ($M+1$)⁺; ¹H NMR (400 MHz, CDCl₃): δ 3.75 (s, 3H), 6.66 (m, 1H), 6.93 (d, 1H), 6.98 (t, 1H), 7.05 (t, 1H), 7.29 (d, 1H), 7.48 (m, 2H), 7.63 (d, 1H), 8.45 (dd, 1H), 11.18 (br, 1H), 11.84 (br, 1H).

5

Example 349

1-[5-Fluoro-2-(thiophene-2-carbonyl)-phenyl]-3-thiazol-2-yl- urea




(2-Amino-4-fluoro-phenyl)-thiophen-2-yl-methanone was (618 g, 28%) was prepared from 2-amino-4-fluoro-benzoic acid (1.55g, 10 mmol) following the general procedure S. N-[5-Fluoro-2-(thiophene-2-carbonyl)-phenyl]-N'-(thiazol-2-yl)urea (113 mg, 65 %) was prepared from (2-amino-4-fluoro-phenyl)-thiophen-2-yl-methanone (110 mg, 0.5 mmol) and 2-aminothiazole (0.060 g, 0.6 mmol) following the general procedure D.

LC-MS (m/z): 348 ($M+1$)⁺; ¹H NMR (400 MHz, DMSO- d_6): δ 7.03 (m, 1H), 7.12 (d, 1H), 7.27 (t, 1H), 7.35 (d, 1H), 7.64 (d, 1H), 7.80 (dd, 1H), 8.07 (d, 1H), 8.14 (d, 1H), 9.63 (br, 1H), 11.60 (br, 1H).

Example 350

N-[2-(2-Methoxy-benzoyl)-5-methoxy-phenyl]-N'-(thiazol-2-yl)urea




(2-Amino-4-methoxy-phenyl)-(2-methoxy-phenyl)-methanone (720 mg, 28 %) was prepared from 2-amino-4-methoxy-benzoic acid (1.67 g, 0.1 mol) following the general

procedure S. N-[2-(2-Methoxy-benzoyl)-5-methoxy-phenyl]-N'-(thiazol-2-yl)urea (131 mg, 68 %) was prepared from (2-amino-4-methoxy-phenyl)-(2-methoxy-phenyl)-methanone (129 mg, 0.5 mmol) and 2-aminothiazole (0.060 g, 0.6 mmol) following the general procedure D.

LC-MS (*m/z*): 384 (*M*+1)⁺; ¹H NMR (400 MHz, CDCl₃): δ 3.77 (s, 3H), 3.93 (s, 3H),
5 6.91 (d, 1H), 6.98 (d, 1H), 7.04 (t, 1H), 7.23 (d, 1H), 7.25 (d, 1H), 7.36 (d, 1H), 7.45 (m, 1H),
7.54 (br, 1H), 8.27 (d, 1H), 10.20 (br, 1H), 12.10 (br, 1H).

Example 351

(2-{3-[2-(2-Methoxy-benzoyl)-5-methyl-phenyl]-ureido}-thiazol-4-yl)-acetic acid ethyl ester



10


(2-{3-[2-(2-Methoxy-benzoyl)-5-methyl-phenyl]-ureido}-thiazol-4-yl)-acetic acid ethyl ester (163 mg, 72 %) was prepared from (2-amino-4-methyl-phenyl)-(2-methoxy-phenyl)-methanone (121 mg, 0.5 mmol) and (2-amino-thiazol-4-yl)-acetic acid

ethyl ester (111 mg, 0.6 mmol) following the general procedure D.

LC-MS (*m/z*): 454 (*M*+1)⁺; ¹H NMR (400 MHz, CDCl₃): δ 1.25 (t, 3H), 2.40 (s, 3H),
15 3.69 (s, 2H), 3.72 (s, 3H), 4.20 (q, 2H), 6.71 (s, 1H), 6.76 (d, 1H), 6.96 (d, 1H), 7.02 (t, 1H),
7.23 (d, 1H), 7.30 (d, 1H), 7.44 (t, 1H), 8.47 (s, 1H), 9.47 (br, 1H), 11.57 (br, 1H).

Example 352

(2-{3-[2-(2-Methoxy-benzoyl)-5-methyl -phenyl]-ureido}-thiazol-4-yl)-acetic acid




- (2-{3-[2-(2-Methoxy-benzoyl)-5-methyl -phenyl]-ureido}-thiazol-4-yl)-acetic acid (181mg, 85 %) was prepared from (2-{3-[2-(2-methoxy-benzoyl)-5-methyl -phenyl]-ureido}-thiazol-4-yl)-acetic acid ethyl ester (227 mg, 0.5 mmol) following the general procedure J.
- LC-MS (m/z): 426 ($M+1$)⁺; ¹H NMR (400 MHz, DMSO- d_6): δ 2.36 (s, 3H), 3.57 (s, 2H), 3.68 (s, 3H), 6.86 (s, 1H), 7.07 (t, 1H), 7.22 (m, 2H), 7.31 (d, 1H), 7.50 (t, 2H), 8.26 (s, 1H), 10.92 (br, 1H), 12.20 (br, 1H).

10

Example 353

2-(2-{3-[2-(2-Methoxy-benzoyl)- 5-methyl-phenyl]-ureido}-thiazol-4-yl)-N-methyl-acetamide




2-(2-{3-[2-(2-Methoxy-benzoyl)- 5-methyl-phenyl]-ureido}-thiazol-4-yl)-N-methyl-acetamide (164 mg, 75 %) was prepared from (2-{3-[2-(2-methoxy-benzoyl)-5-methyl - phenyl]-ureido}-thiazol-4-yl)-acetic acid (213 mg, 0.5 mmol) and 1 N solution of methylamine in THF (0.5 mL, 0.5 mmol) following the general procedure K.

- 5 LC-MS (*m/z*): 439 (*M*+1)⁺; ¹H NMR (400 MHz, CDCl₃): δ 2.41 (s, 3H), 2.74 (d, 3H), 3.62 (s, 2H), 3.73 (s, 3H), 6.66 (s, 1H), 6.79 (d, 1H), 6.97 (d, 1H), 7.03 (t, 1H), 7.24 (d, 1H), 7.32 (d, 1H), 7.46 (m, 1H), 8.45 (s, 1H), 11.60 (br, 1H), 11.82 (br, 1H).

Example 354

- 10 2-(2-{3-[2-(2-Methoxy-benzoyl)- 5-methyl-phenyl]-ureido}-thiazol-4-yl)-N-(methoxy-ethyl)-acetamide




- 15 2-(2-{3-[2-(2-Methoxy-benzoyl)- 5-methyl-phenyl]-ureido}-thiazol-4-yl)-N-(methoxy-ethyl)-acetamide (169 mg, 70 %) was prepared from (2-{3-[2-(2-methoxy-benzoyl)-5-methyl - phenyl]-ureido}-thiazol-4-yl)-acetic acid (213 mg, 0.5 mmol) and 2-methoxyethylamine (38 mg, 0.5 mmol) following the general procedure K.

- 20 LC-MS (*m/z*): 483 (*M*+1)⁺; ¹H NMR (400 MHz, CDCl₃): δ 2.41 (s, 3H), 3.33 (s, 3H), 3.48 (m, 4H), 3.61 (s, 2H), 3.73 (s, 3H), 6.65 (s, 1H), 6.76 (d, 1H), 6.97 (d, 1H), 7.03 (t, 1H), 7.25 (d, 1H), 7.31 (d, 1H), 7.42 (bs, 1H), 7.46 (m, 1H), 8.48 (s, 1H), 10.02 (br, 1H), 11.62 (br, 1H).

Example 355

N-(2-Methanesulfonyl-ethyl)-2-(2-{3-[2-(2-methoxy-benzoyl)- 5-methyl-phenyl]-ureido}-thiazol-4-yl)-acetamide




N-(2-Methanesulfonyl-ethyl)-2-(2-{3-[2-(2-methoxy-benzoyl)- 5-methyl-phenyl]-ureido}-thiazol-4-yl)-acetamide (181 mg, 68 %) was prepared from (2-{3-[2-(2-methoxy-benzoyl)-5-methyl -phenyl]-ureido}-thiazol-4-yl)-acetic acid (213 mg, 0.5 mmol) and 2-methanesulfonyl-ethylamine (61 mg, 0.5 mmol) following the general procedure K.

LC-MS (*m/z*): 531 (*M*+1)⁺; ¹H NMR (400 MHz, CDCl₃): δ 2.41 (s, 3H), 2.93 (s, 3H), 3.33 (t, 3H), 3.63 (s, 2H), 3.74 (s, 3H), 3.78 (m, 2H), 6.61 (s, 1H), 6.76 (d, 1H), 6.98 (d, 1H), 7.04 (t, 1H), 7.25 (d, 1H), 7.31 (d, 1H), 7.46 (m, 1H), 7.97 (t, 1H), 8.48 (s, 1H), 11.40 (br, 1H), 11.51 (br, 1H).

10

Example 356

(2-{3-[5-Fluoro-2-(2-Methoxy-benzoyl)-phenyl]-ureido}-thiazol-4-yl)-acetic acid ethyl ester




(2-{3-[5-Fluoro-2-(2-Methoxy-benzoyl)-phenyl]-ureido}-thiazol-4-yl)-acetic acid ethyl ester (164 mg, 72 %) was prepared from (2-amino-4-fluoro-phenyl)-(2-methoxy-phenyl)-methanone (123 mg, 0.5 mmol) and (2-amino-thiazol-4-yl)-acetic acid

ethyl ester (111 mg, 0.6 mmol) following the general procedure (D).

5 LC-MS (*m/z*): 458 (*M*+1)⁺; ¹H NMR (400 MHz, CDCl₃): δ 1.24 (t, 3H), 3.68 (s, 2H), 3.78 (s, 3H), 6.68 (s, 1H), 6.74 (d, 1H), 6.93 (d, 1H), 7.06 (t, 1H), 7.24 (m, 1H), 7.30 (d, 1H), 7.44 (t, 1H), 8.47 (s, 1H), 9.47 (br, 1H), 11.57 (br, 1H).

Example 357

10 (2-{3-[5-Fluoro-2-(2-methoxy-benzoyl)-phenyl]-ureido}-thiazol-4-yl)-acetic acid




(2-{3-[5-Fluoro-2-(2-methoxy-benzoyl)-phenyl]-ureido}-thiazol-4-yl)-acetic acid (161 mg, 75 %) was prepared from (2-{3-[5-fluoro-2-(2-methoxy-benzoyl)-phenyl]-ureido}-thiazol-4-yl)-acetic acid ethyl ester (229, 0.5 mmol) following the general procedure J.

15 LC-MS (*m/z*): 430 (*M*+1)⁺; ¹H NMR (400 MHz, DMSO-*d*₆): δ 3.57 (s, 2H), 3.69 (s, 3H), 6.89 (m, 2 H), 7.08 (t, 1H), 7.18 (d, 1H), 7.33 (d, 1H), 7.40 (m, 1H), 7.54 (m, 1H), 8.28 (d, 1H), 11.12 (br, 1H), 12.32 (br, 1H).

Example 358

20 2-(2-{3-[5-Fluoro-2-(2-methoxy-benzoyl)-phenyl]-ureido}-thiazol-4-yl)-N-methyl-acetamide




2-(2-{3-[5-Fluoro-2-(2-methoxy-benzoyl)-phenyl]-ureido}-thiazol-4-yl)-N-methyl-acetamide (144 mg, 65 %) was prepared from (2-{3-[5-fluoro-2-(2-methoxy-benzoyl)-phenyl]-ureido}-thiazol-4-yl)-acetic acid (215 mg, 0.5 mmol) and 1N solution of methyl amine in THF (0.5 mL, 0.5 mmol) following the general procedure K.

LC-MS (m/z): 443 ($M+1$)⁺; ¹H NMR (400 MHz, CDCl₃): δ 2.71 (d, 3H), 3.66 (s, 2H), 3.70 (s, s, 3H), 6.64 m, 2H), 6.85 (br, 1H), 7.36 (m, 2 H), 6.95 (d, 1 H), 7.02 (t, 1H), 7.24 (dd, 1H), 7.44 (m, 2H), 8.43 (dd, 1H), 10.72 (br, 1H), 11.84 (br, 1H).

10 Example 359


2-(2-{3-[5-Fluoro-2-(2-methoxy-benzoyl)-phenyl]-ureido}-thiazol-4-yl)- N-(2-methoxy-ethyl)-acetamide



2-(2-{3-[5-Fluoro-2-(2-methoxy-benzoyl)-phenyl]-ureido}-thiazol-4-yl)- N-(2-methoxy-ethyl)-acetamide (158 mg, 65 %) was prepared from (2-{3-[5-fluoro-2-(2-methoxy-benzoyl)-phenyl]-ureido}-thiazol-4-yl)-acetic acid (215 mg, 0.5 mmol) and 2-methoxyethylamine (38 mg, 0.5 mmol) following the general procedure K.

LC-MS (m/z): 487 ($M+1$)⁺; ¹H NMR (400 MHz, CDCl₃): δ 3.21 (s, 3H), 3.48 (m, 4H), 3.62 (s, 2 H), 3.73 (s, 3H), 6.62-6.66 (m, 2H), 6.97 (d, 1H), 7.04 (t, 1H), 7.28 (d, 1H), 7.42 (br, 1H), 7.42-7.50 (m, 2H), 8.46 (dd, 1H), 10.20 (br, 1H), 11.28 (br, 1H),

5


Example 360**1-[2-(Hydroxyimino-phenyl-methyl)-phenyl]-3-thiazol-2-yl-urea**

1-[2-(Hydroxyimino-phenyl-methyl)-phenyl]-3-thiazol-2-yl-urea (101 mg, 60 %) was prepared from (2-Amino-phenyl)-phenyl-methanone oxime (106 mg, 0.5 mmol) and 2-amino thiazole (0.06 mg, 0.6 mmol) following the general procedure D.

10

LC-MS (m/z): 339 ($M+1$)⁺; ¹H NMR (400 MHz, DMSO- d_6): δ 7.2 (dd, 1 H), 7.06 (d, 1H), 7.15 (m, 1 H), 7.30 (d, 1H), 7.34-7.43 (m, 6H), 8.03 (d, 1H), 8.05 (br, 1H), 11.07 (br, 1H), 11.63 (br, 1H),


15

EXAMPLE 361**1-[5-Chloro-2-(hydroxyimino-phenyl-methyl)-phenyl]-3-thiazol-2-yl-urea**

1-[5-Chloro-2-(hydroxyimino-phenyl-methyl)-phenyl]-3-thiazol-2-yl-urea (93 mg, 50%) was prepared from (2-Amino-4-chloro-phenyl)-phenyl-methanone oxime (123 mg, 0.5 mmol) and 2-amino thiazole (60 mg, 0.6 mmol) following the general procedure D.


20

LC-MS (m/z): 373 ($M+1$)⁺; ¹H NMR (400 MHz, DMSO- d_6): δ 7.04 (dd, 1 H), 7.12 (d, 1H), 7.16 (m, 1 H), 7.36 (d, 1H), 7.36-7.48 (m, 5H), 8.03 (d, 1H), 8.12 (br, 1H), 11.23 (br, 1H), 11.72 (br, 1H),

EXAMPLE 362**1-(2-Benzoyl-5-methyl-phenyl)-3-thiazol-2-yl-urea**

5 1-(2-Benzoyl-5-methyl-phenyl)-3-thiazol-2-yl-urea (60 mg, 65%) was prepared from (2-(2-amino-4-methyl-phenyl)-phenyl-methanone (55 mg, 0.25 mmol) following the general procedure D.


LC-MS (*m/z*): 338 (*M*+1)⁺; ¹H NMR (400 MHz, CDCl₃): δ 2.40 (s, 3H), 6.93 (s, 1H), 7.19 (t, 1H), 7.28 (s, 1H), 7.48 (t, 1H), 7.54-7.61 (m, 4H), 7.68 (dd, 1H), 8.54 (d, 1H), 10.64 (br, 1H), 11.22 (br, 1H).

EXAMPLE 363**1-[4-Bromo-2-(2-fluoro-benzoyl)-phenyl]-3-thiazol-2-yl-urea**

15 1-[4-Bromo-2-(2-fluoro-benzoyl)-phenyl]-3-thiazol-2-yl-urea (73 mg, 69%) was prepared from (2-amino-5-bromo-phenyl)-(2-fluoro-phenyl)-methanone (73 mg, 0.25 mmol) following the general procedure D.

LC-MS (*m/z*): 420 (*M*+1)⁺; ¹H NMR (400 MHz, CDCl₃): δ 6.93 (s, 1H), 7.19 (t, 1H), 7.28 (s, 1H), 7.48 (t, 1H), 7.54-7.61 (m, 3H), 7.68 (dd, 1H), 8.54 (d, 1H), 10.64 (br, 1H), 11.22 (br, 1H).

Example 364**1-[5-Chloro-2-(2-fluoro-benzoyl)-phenyl]-3-thiazol-2-yl-urea**




1-[5-Chloro-2-(2-fluorobenzoyl)phenyl]-3-thiazol-2-yl-urea (132 mg, 70%) was prepared from (2-amino-4-chlorophenyl)-(2-fluorophenyl)methanone (125 mg, 0.5 mmol) and 2-aminothiazole (60 mg, 0.6 mmol) following the general procedure D.

5 LC-MS (m/z): 376 ($M+1$)⁺; ¹H NMR (400 MHz, DMSO- d_6): δ 6.76 (s, 1H), 6.89 (s, 1H), 7.16 (m, 1H), 7.25 (d, 1H), 7.48 (m, 4H), 8.58 (br, 1H), 10.88 (br, 1H).

Example 365

1-[4-Fluoro-2-(2-methylsulfonylphenoxy)phenyl]-3-thiazol-2-yl-urea



10


4-Fluoro-2-(2-methylsulfonylphenoxy)-1-nitrobenzene (1.06 g, 68%) was prepared from 2-hydroxythioanisole (0.77 g, 5.5 mmol) and 2,5-difluoro-1-nitrobenzene (0.80 g, 5.0 mmol) following the general procedure A. This was reduced to 4-fluoro-2-(2-methanesulfonylphenoxy)phenylamine (0.52 g, 62%) following general procedure C. 1-[4-Fluoro-2-(2-methylsulfonylphenoxy)phenyl]-3-thiazol-2-yl-urea (117 mg, 60%) was prepared from 4-fluoro-2-(2-methylsulfonylphenoxy)aniline (125 mg, 0.5 mmol) and 2-aminothiazole (60 mg, 0.6 mmol) following the general procedure D.

15

LC-MS (m/z): 376 ($M+1$)⁺; ¹H NMR (400 MHz, CDCl₃): δ 2.41 (s, 3H), 6.41 (dd, 1H), 6.56 (br, 1H), 6.76 (d, 1H), 6.94 (d, 1H), 7.15-7.34 (m, 4H), 8.26 (dd, 1H), 9.88 (br, 1H),
20 11.12 (br, 1H),

Example 366

2-[4-Fluoro-2-(3-thiazol-2-yl-ureido)-phenoxy]-6-methoxy-benzoic acid methyl ester




2-(4-Fluoro-2-nitro-phenoxy)-6-methoxy-benzoic acid methyl ester (0.94 g, 58%) was prepared from 2-hydroxy-6-methoxy-benzoic acid methyl ester (1.01 g, 5.5 mmol) and 2,5-difluoro-1-nitro-benzene (0.80 g, 5.0 mmol) following the general procedure A. This was reduced to 2-(2-amino-4-fluoro-phenoxy)-6-methoxy-benzoic acid methyl ester (0.51 g, 60%) following general procedure C. 2-[4-Fluoro-2-(3-thiazol-2-yl-ureido)-phenoxy]-6-methoxy-benzoic acid methyl ester (129 mg, 62%) was prepared from 2-(2-amino-4-fluoro-phenoxy)-6-methoxy-benzoic acid methyl ester (145 mg, 0.5 mmol) and 2-aminothiazole (60 mg, 0.6 mmol) following the general procedure D.

LC-MS (m/z): 418 ($M+1$)⁺; ¹H NMR (400 MHz, CDCl₃): δ 3.62 (s, 3H), 3.64 (s, 3H), 6.31 (s, 1H), 6.45 (dd, 1H), 6.55 (s, 1H), 7.05 (d, 1H), 7.22 (t, 1H), 7.33 (br, 1H), 7.45 (d, 1H), 78.02 (d, 1H), 8.63 (br, 1H), 11.74 (br, 1H).

15

Example 367

2-[4-Fluoro-2-(3-thiazol-2-yl-ureido)-phenoxy]-6-methoxy-N-methyl-benzamide




2-[4-Fluoro-2-(3-thiazol-2-yl-ureido)-phenoxy]-6-methoxy-N-methyl-benzamide (150 mg, 72%) was prepared from 2-[4-fluoro-2-(3-thiazol-2-yl-ureido)-phenoxy]-6-methoxy-

benzoic acid (201 mg, 0.5 mmol) and 1N solution of methyl amine in THF (0.5 mL, 0.5 mmol) following the general procedure K.

LC-MS (m/z): 417 ($M+1$)⁺; ¹H NMR (400 MHz, DMSO- d_6): δ 3.42 (d, 3H), 3.67 (s, 3H), 6.51 (dd, 1H), 6.67 (m, 1H), 7.14 (m, 2H), 7.29 (m, 2H), 7.37 (d, 1H), 7.98 (dd, 1H), 8.11 (d, 1H), 9.13 (dd, 1H), 11.20 (br, 1H).

Example 368

1-[5-Fluoro-2-(2-methanesulfonyl-phenoxy)-phenyl]-3-thiazol-2-yl-urea




10 1-[5-Fluoro-2-(2-methanesulfonyl-phenoxy)-phenyl]-3-thiazol-2-yl-urea (153 mg, 75%) was prepared from 1-[5-fluoro-2-(2-methanesulfanyl-phenoxy)-phenyl]-3-thiazol-2-yl-urea (187 mg, 0.5 mmol) following the general procedure R.

LC-MS (m/z): 408 ($M+1$)⁺; ¹H NMR (400 MHz, CDCl₃): δ 3.44 (s, 3H), 6.80 (m, 1H), 6.84 (d, 1H), 7.03 (d, 1H), 7.27 (m, 2H), 7.33 (d, 1H), 7.56 (m, 1H), 8.05 (dd, 1H), 8.27 (dd, 1H), 9.13 (br, 1H), 11.12 (br, 1H).

Example 369

1-[2-(4,5-Dimethoxy-2-methyl-phenoxy)-phenyl]-3-thiazol-2-yl-urea




1-[2-(4,5-Dimethoxy-2-methyl-phenoxy)-phenyl]-3-thiazol-2-yl-urea (131 mg, 68%) was prepared from 2-(4,5-dimethoxy-2-methyl-phenoxy)-phenylamine (130 mg, 0.5 mmol) and 2-aminothiazole (60 mg, 0.6 mmol) following the general procedure D.

LC-MS (m/z): 386 ($M+1$)⁺; ¹H NMR (400 MHz, CDCl₃): δ 2.29 (s, 3H), 3.80 (s, 3H), 3.94 (s, 3H), 6.76 (s, 1H), 6.92 (d, 1H), 7.02 (t, 1H), 7.44 (d, 1H), 7.55 (t, 2H), 7.76 (d, 1H), 8.54 (d, 1H), 11.34 (br, 1H).

Example 370

3-[4-Fluoro-2-(3-thiazol-2-yl-ureido)-phenoxy]-benzoic acid methyl ester



10

3-(4-Fluoro-2-nitro-phenoxy)-benzoic acid methyl ester (0.73 g, 50%) was prepared from 3-hydroxy-benzoic acid methyl ester (0.84 g, 5.5 mmol) and 2,5-difluoro-1-nitro-benzene (0.80 g, 5.0 mmol) following the general procedure A. This was reduced to 2-(2-amino-4-fluoro-phenoxy)-benzoic acid methyl ester (0.37 g, 58%) following general procedure C. 3-[4-Fluoro-2-(3-thiazol-2-yl-ureido)-phenoxy]-benzoic acid methyl ester (145 mg, 75%) was prepared from 3-(2-amino-4-fluoro-phenoxy)-benzoic acid methyl ester (130 mg, 0.5 mmol) and 2-aminothiazole (60 mg, 0.6 mmol) following the general procedure D.


15

LC-MS (m/z): 388 ($M+1$)⁺; ¹H NMR (400 MHz, CDCl₃): δ 3.87 (s, 3H), 6.78 (m, 1H), 6.89 (d, 1H), 7.12 (m, 1H), 7.18 (br, 1H), 7.34 (m, 1H), 7.41 (m, 1H), 7.60 (m, 1H), 7.72 (d, 1H), 8.22 (dd, 1H), 10.50 (br, 1H).

20

Example 371

3-[4-Fluoro-2-(3-thiazol-2-yl-ureido)-phenoxy]-benzoic acid




3-[4-Fluoro-2-(3-thiazol-2-yl-ureido)-phenoxy]-benzoic acid (150 mg, 80%) was prepared from 3-[4-fluoro-2-(3-thiazol-2-yl-ureido)-phenoxy]-benzoic acid methyl ester (194 mg, 0.5 mmol) following the general procedure J

- 5 LC-MS (m/z): 374 ($M+1$)⁺; ¹H NMR (400 MHz, CDCl₃): δ 6.85 (m, 1H), 7.01 (dd, 1H), 7.13 (d, 1H), 7.26 (dd, 1H), 7.34 (d, 1H), 7.42 (m, 1H), 7.52 (t, 1H), 8.14 (dd, 1H), 9.47 (br, 1H), 11.28, br, 1H).

Example 372

- 10 2-[4-Fluoro-2-(3-thiazol-2-yl-ureido)-phenoxy]-benzoic acid methyl ester




- 15 2-(4-Fluoro-2-nitro-phenoxy)-benzoic acid methyl ester (0.81 g, 55%) was prepared from 2-hydroxy-benzoic acid methyl ester (0.84 g, 5.5 mmol) and 2,5-difluoro-1-nitro-benzene (0.80 g, 5.0 mmol) following the general procedure A. This was reduced to 2-(2-amino-4-fluoro-phenoxy)-benzoic acid methyl ester (0.43 g, 60%) following general procedure C. 2-[4-Fluoro-2-(3-thiazol-2-yl-ureido)-phenoxy]-benzoic acid methyl ester (125 mg, 65%) was prepared from 2-(2-amino-4-fluoro-phenoxy)-benzoic acid methyl ester (130 mg, 0.5 mmol) and 2-aminothiazole (60 mg, 0.6 mmol) following the general procedure D.

LC-MS (m/z): 388 ($M+1$)⁺; ¹H NMR (400 MHz, CDCl₃): δ 3.84 (s, 3H), 6.73 (m, 1H), 6.85 (d, 1H), 7.14 (m, 1H), 7.13 (br, 1H), 7.31 (m, 1H), 7.41 (m, 1H), 7.54 (m, 1H), 7.71 (d, 1H), 8.22 (dd, 1H), 10.54 (br, 1H).

5

Example 373

2-[4-Fluoro-2-(3-thiazol-2-yl-ureido)-phenoxy]-N-methyl-benzamide




2-[4-Fluoro-2-(3-thiazol-2-yl-ureido)-phenoxy]-N-methyl-benzamide was prepared from 2-[4-fluoro-2-(3-thiazol-2-yl-ureido)-phenoxy]-benzoic acid and 1N solution of methyl amine in THF (0.5 mL, 0.5 mmol) following the general procedure K.

LC-MS (m/z): 387 ($M+1$)⁺; ¹H NMR (400 MHz, CDCl₃): δ 3.53 (d, 3H), 6.73 (m, 1H), 5.86 (br, 1H), 6.88 (d, 1H), 7.16 (m, 1H), 7.18 (br, 1H), 7.35 (m, 1H), 7.41 (m, 1H), 7.60 (m, 1H), 7.76 (d, 1H), 8.23 (dd, 1H), 10.22 (br, 1H).

15

Example 374

N-{3-[4-Fluoro-2-(3-thiazol-2-yl-ureido)-phenoxy]-2-methoxy-phenyl}-methanesulfonamide




N-[3-(4-Fluoro-2-nitro-phenoxy)-2-methoxy-phenyl]-methanesulfonamide (0.86 g, 48%) was prepared from N-(3-hydroxy-2-methoxy-phenyl)-methanesulfonamide (1.2 g, 5.5 mmol) and 2,5-difluoro-1-nitro-benzene (0.80 g, 5.0 mmol) following the general procedure A. This was reduced to N-[3-(2-amino-4-fluoro-phenoxy)-2-methoxy-phenyl]-

5 methanesulfonamide (0.51 g, 70%) following general procedure C. N-[3-[4-Fluoro-2-(3-thiazol-2-yl-ureido)-phenoxy]-2-methoxy-phenyl]-methanesulfonamide (144 mg, 64%) was prepared from N-[3-(2-amino-4-fluoro-phenoxy)-2-methoxy-phenyl]-methanesulfonamide (123 mg, 0.5 mmol) and 2-aminothiazole (60 mg, 0.6 mmol) following the general procedure D.

10 LC-MS (m/z): 453 ($M+1$)⁺; ¹H NMR (400 MHz, CDCl₃): δ 3.27 (s, 3H), 3.89 (s, 3H), 6.52 (d, 1H), 6.72 (m, 1H), 6.95 (d, 1H), 7.04 (m, 1H), 7.16 (m, 1H), 7.34 (d, 1H), 7.45 (dd, 1H), 7.63 (d, 1H), 8.21 (dd, 1H), 8.44 (br, 1H), 8.85 (br, 1H).

Example 375

15 N-[4-(2-Fluoro-6-methoxy-phenoxy)-3-(3-thiazol-2-yl-ureido)-phenyl]-acetamide




N-[4-(2-Fluoro-6-methoxy-phenoxy)-3-(3-thiazol-2-yl-ureido)-phenyl]-acetamide (81 mg, 78%) was prepared from N-[3-amino-4-(2-fluoro-6-methoxy-phenoxy)-phenyl]-acetamide (72 mg, 0.25 mmol) following the general procedure D.

20 LC-MS (m/z): 417 ($M+1$)⁺; ¹H NMR (400 MHz, DMSO-d₆): δ 1.78 (s, 3H), 3.43 (s, 3H), 6.32 (m, 1H), 6.44 (dd, 1H), 6.48 (d, 1H), 6.63 (t, 1H), 6.86 (dd, 1H), 6.97 (d, 1H), 7.13 (d, 1H), 7.88 (d, 1H), 8.90 (br, 1H), 9.32 (br, 1H), and 10.46 (br, 1H).

Example 376

25 N-[4-(2-Fluoro-6-methoxy-phenoxy)-3-(3-thiazol-2-yl-ureido)-phenyl]-methanesulfonamide




N-[4-(2-Fluoro-6-methoxy-phenoxy)-3-(3-thiazol-2-yl-ureido)-phenyl]
methanesulfonamide (101 mg, 89%) was prepared from N-[3-amino-4-(2-fluoro-6-methoxy-
phenoxy)-phenyl]-methanesulfonamide (81 mg, 0.25 mmol) following the general procedure
5 L.

LC-MS (m/z): 453 ($M+1$)⁺; ¹H NMR (400 MHz, DMSO- d_6): δ 3.13 (s, 3H), 4.00 (s, 3H),
6.70 (d, 1H), 6.98 (dd, 1H), 7.14 (t, 1H), 7.21 (d, 1H), 7.38 (t, 1H), 7.46 (q, 1H), 7.55 (t, 1H),
8.14 (br, 1H), 8.38 (dd, 1H), 8.46 (d, 1H), and 10.86 (br, 1H).

10 Example 377

(2-{3-[2-(2-Fluoro-6-methoxy-phenoxy)-5-methanesulfonylamino-phenyl]-ureido}-
thiazol-4 yl)-acetic acid ethyl ester




(2-{3-[2-(2-Fluoro-6-methoxy-phenoxy)-5-methanesulfonylamino-phenyl]-ureido}-
15 thiazol-4 yl)-acetic acid ethyl ester (118 mg, 87%) was prepared from N-[3-amino-4-(2-fluoro-
6-methoxy-phenoxy)-phenyl]-methanesulfonamide (81 mg, 0.25 mmol) following the general
procedure D.

LC-MS (*m/z*): 539 (*M*+1)⁺; ¹H NMR (400 MHz, DMSO-*d*₆): δ 1.24 (t, 2H), 3.03 (s, 3H), 3.74 (s, 2H), 3.80 (s, 3H), 4.21 (q, 3H), 6.59 (d, 1H), 6.68 (s, 1H), 6.78 (s, 1H), 6.83 (dd, 1H), 7.12-7.19 (m, 2H), 8.12 (d, 1H), 8.42 (s, 1H), 9.60 (br, 1H), and 11.84 (br, 1H).

5

Example 378

(2-{3-[2-(2-Fluoro-6-methoxy-phenoxy)-5-methanesulfonylamino-phenyl]-ureido}-thiazol-4-yl)-acetic acid




(2-{3-[2-(2-Fluoro-6-methoxy-phenoxy)-5-methanesulfonylamino-phenyl]-ureido}-thiazol-4-yl)-acetic acid (85 mg, 83%) was prepared from (2-{3-[2-(2-fluoro-6-methoxy-phenoxy)-5-methanesulfonylamino-phenyl]-ureido}-thiazol-4-yl)-acetic acid ethyl ester (108 mg, 0.20 mmol) following the general procedure J.

LC-MS (*m/z*): 511 (*M*+1)⁺; ¹H NMR (400 MHz, CD₃OD): δ 3.12 (s, 3H), 3.54 (s, 2H), 3.81 (s, 3H), 6.54 (d, 1H), 6.64 (s, 1H), 6.84 (s, 1H), 6.92 (d, 1H), 7.11 (m, 1H), 7.16 (s, 1H), 7.44 (m, 1H), 8.46 (s, 1H), 9.40 (br, 1H), 10.83 (br, 1H), and 11.84 (br, 1H).

15

Example 379

N,N-[4-(2-Fluoro-6-methoxy-phenoxy)-3-(3-thiazol-2-yl-ureido)-phenyl]dimethanesulfonamide




N,N-[4-(2-Fluoro-6-methoxy-phenoxy)-3-(3-thiazol-2-yl-ureido)-phenyl]dimethanesulfonamide (115 mg, 87%) was prepared from N-[3-amino-4-(2-fluoro-6-methoxy-phenoxy)-phenyl]-dimethanesulfonamide (101 mg, 0.25 mmol) following the general procedure D.

LC-MS (m/z): 531 ($M+1$)⁺; ¹H NMR (400 MHz, DMSO- d_6): δ 3.22 (s, 3H), 3.24 (s, 3H), 4.11 (s, 3H), 6.78 (d, 1H), 6.89 (d, 1H), 7.11 (dd, 1H), 7.24 (d, 1H), 7.29 (m, 1H), 7.42 (t, 1H), 7.63 (m, 1H), 8.54 (d, 1H), 8.65 (dd, 1H), and 11.38 (br, 1H).

Example 380

N-[4-Isobutoxy-3-(3-thiazol-2-yl-ureido)-phenyl]-methanesulfonamide




N-[4-Isobutoxy-3-(3-thiazol-2-yl-ureido)-phenyl]-methanesulfonamide (77 mg, 80%) was prepared from N-(3-amino-4-isobutoxy-phenyl)-methanesulfonamide (64 mg, 0.25 mmol) following the general procedure D.

LC-MS (m/z): 385 ($M+1$)⁺; ¹H NMR (400 MHz, CDCl₃): δ 1.20 (d, 6H), 1.70 (m, 1H), 3.05 (d, 2H), 3.82 (s, 3H), 6.76 (dd, 1H), 6.86 (d, 1H), 6.94 (m, 1H), 7.36 (d, 1H), 7.42 (d, 1H), 8.42 (d, 1H), 9.56 (br, 1H), and 11.63 (br, 1H).

Example 381

1-[5-Amino-2-(2-fluoro-6-methoxy-phenoxy)-phenyl]-3-thiazol-2-yl-urea




1-[5-Amino-2-(2-fluoro-6-methoxy-phenoxy)-phenyl]-3-thiazol-2-yl-urea (3.52 g, 94%) was prepared from 3-amino-4-(2-fluoro-6-methoxy-phenoxy)-phenyl]-carbamic acid tert-butyl ester (3.48 g, 10.0 mmol) following the general procedure D and the corresponding *t*-BOC protected product [4-(2-fluoro-6-methoxy-phenoxy)-3-(3-thiazol-2-yl-ureido)-phenyl]-carbamic acid tert-butyl ester was subjected to deprotection conditions with hydrochloric acid (40 ml, 4.0 M in dioxane).

LC-MS (*m/z*): 375 (*M*+1)⁺; ¹H NMR (400 MHz, CD₃OD): δ 3.76 (s, 3H), 5.24 (br, 2H), 6.68 (d, 1H), 6.88 (q, 1H), 6.98 (d, 1H), 7.16 (m, 1H), 7.38 (dd, 1H), 7.63 (dd, 1H), 8.34 (dd, 1H), 8.44 (d, 1H), 8.86 (br, 1H), and 11.27 (br, 1H).

Example 382

Pyridine-2-carboxylic acid [4-(2-fluoro-6-methoxy-phenoxy)-3-(3-thiazol-2-yl-ureido)-phenyl]-amide




Pyridine-2-carboxylic acid [4-(2-fluoro-6-methoxy-phenoxy)-3-(3-thiazol-2-yl-ureido)-phenyl]-amide (98 mg, 82%) was prepared from 1-[5-amino-2-(2-fluoro-6-methoxy-phenoxy)-phenyl]-3-thiazol-2-yl-urea (94 mg, 0.25 mmol) following the general procedure K.

LC-MS (m/z): 480 ($M+1$)⁺; ¹H NMR (400 MHz, CDCl₃): δ 3.74 (s, 3H), 6.64 (d, 1H),
5 6.74 (d, 1H), 6.78 (s, 1H), 6.82 (m, 1H), 7.16 (q, 1H), 7.46 (dd, 1H), 7.48 (d, 1H), 7.54 (t, 1H),
7.72 (br, 1H), 7.84 (dd, 1H), 7.88 (t, 1H), 8.24 (d, 1H), 8.38 (br, 1H), 8.62 (d, 1H), and 10.06
(br, 1H).

Example 383

10 N-[4-(2-Fluoro-6-methoxy-phenoxy)-3-(3-thiazol-2-yl-ureido)-phenyl]-isonicotinamide




N-[4-(2-Fluoro-6-methoxy-phenoxy)-3-(3-thiazol-2-yl-ureido)-phenyl]-isonicotinamide (104 mg, 86%) was prepared from 1-[5-amino-2-(2-fluoro-6-methoxy-phenoxy)-phenyl]-3-thiazol-2-yl-urea (94 mg, 0.25 mmol) following the general procedure K.

15 LC-MS (m/z): 480 ($M+1$)⁺; ¹H NMR (400 MHz, CD₃OD): δ 3.58 (s, 3H), 6.32 (d, 1H),
6.62 (dd, 1H), 6.64 (d, 1H), 6.66 (d, 1H), 6.96 (m, 2H), 7.06 (d, 1H), 7.18 (dd, 1H), 7.60 (d,
2H), 8.12 (d, 1H), 8.40 (br, 1H), 8.48 (d, 1H), 9.62 (d, 1H), and 11.34 (br, 1H).

Example 384

20 5-Methyl-pyrazine-2-carboxylic acid [4-(2-fluoro-6-methoxy-phenoxy)-3-(3-thiazol-2-yl-ureido)-phenyl]-amide




5-Methyl-pyrazine-2-carboxylic acid [4-(2-fluoro-6-methoxy-phenoxy)-3-(3-thiazol-2-yl-ureido)-phenyl]-amide (103 mg, 83%) was prepared from 1-[5-amino-2-(2-fluoro-6-methoxy-phenoxy)-phenyl]-3-thiazol-2-yl-urea (94 mg, 0.25 mmol) following the general procedure K.

LC-MS (*m/z*): 495 (*M*+1)⁺; ¹H NMR (400 MHz, CDCl₃): δ 2.68 (s, 3H), 3.63 (s, 3H), 6.52 (d, 1H), 6.61 (d, 1H), 6.66 (d, 2H), 7.06 (q, 1H), 7.36 (s, 1H), 7.78 (d, 1H), 8.20 (br, 1H), 8.43 (s, 1H), 8.76 (br, 1H), 9.32 (s, 1H), 9.62 (s, 1H), and 11.80 (br, 1H).

Example 385

2,2,2-Trifluoro-ethanesulfonic acid [4-(2-fluoro-6-methoxy-phenoxy)-3-(3-thiazol-2-yl-ureido)-phenyl]-amide




2,2,2-Trifluoro-ethanesulfonic acid [4-(2-fluoro-6-methoxy-phenoxy)-3-(3-thiazol-2-yl-ureido)-phenyl]-amide (105 mg, 81%) was prepared from 1-[5-amino-2-(2-fluoro-6-methoxy-phenoxy)-phenyl]-3-thiazol-2-yl-urea (94 mg, 0.25 mmol) following the general procedure L.

LC-MS (m/z): 521 ($M+1$)⁺; ¹H NMR (400 MHz, CD₃OD): δ 3.75 (s, 3H), 3.86 (q, 2H), 6.48 (d, 1H), 6.80 (m, 2H), 6.88 (d, 1H), 7.16 (q, 1H), 7.28 (d, 1H), 7.47 (s, 1H), 8.08 (d, 1H), 8.40 (br, 1H), 9.62 (d, 1H), and 11.34 (br, 1H).

5

Example 386

N-[4-(2-Fluoro-6-methoxy-phenoxy)-3-(3-thiazol-2-yl-ureido)-phenyl]-C-methanesulfonyl-methanesulfonamide




10 N-[4-(2-Fluoro-6-methoxy-phenoxy)-3-(3-thiazol-2-yl-ureido)-phenyl]-C-methanesulfonyl-methanesulfonamide (102 mg, 77%) was prepared from 1-[5-amino-2-(2-fluoro-6-methoxy-phenoxy)-phenyl]-3-thiazol-2-yl-urea (94 mg, 0.25 mmol) following the general procedure L.

LC-MS (m/z): 530 ($M+1$)⁺; ¹H NMR (400 MHz, CD₃OD): δ 3.01 (s, 3H), 3.27 (s, 2H), 3.82 (s, 3H), 6.56 (d, 1H), 6.85 (t, 1H), 6.94 (d, 1H), 7.21 (q, 1H), 7.34 (d, 1H), 7.46 (m, 1H),
15 7.96 (s, 1H), 8.22 (d, 1H), 8.40 (br, 1H), 9.62 (br, 1H), and 10.94 (br, 1H).

Example 387

1-Methyl-1H-imidazole-4-sulfonic acid [4-(2-fluoro-6-methoxy-phenoxy)-3-(3-thiazol-2-yl-ureido)-phenyl]-amide




1-Methyl-1H-imidazole-4-sulfonic acid [4-(2-fluoro-6-methoxy-phenoxy)-3-(3-thiazol-2-yl-ureido)-phenyl]-amide (108 mg, 83%) was prepared from 1-[5-amino-2-(2-fluoro-6-methoxy-phenoxy)-phenyl]-3-thiazol-2-yl-urea (94 mg, 0.25 mmol) following the general procedure L.

LC-MS (m/z): 519 ($M+1$)⁺; ¹H NMR (400 MHz, CD₃OD): δ 3.00 (s, 3H), 3.78 (s, 3H), 6.46 (d, 1H), 6.74 (dd, 1H), 6.83 (d, 2H), 6.91 (d, 1H), 7.16-7.22 (m, 1H), 7.31 (d, 1H), 7.48 (m, 1H), 7.66 (s, 1H), 7.99 (d, 1H), 8.18 (d, 1H), 9.88 (br, 1H), and 11.24 (br, 1H).

10

Example 388

1,2-Dimethyl-1H-imidazole-4-sulfonic acid [4-(2-fluoro-6-methoxy-phenoxy)-3-(3-thiazol-2-yl-ureido)-phenyl]-amide



1,2-Dimethyl-1H-imidazole-4-sulfonic acid [4-(2-fluoro-6-methoxy-phenoxy)-3-(3-thiazol-2-yl-ureido)-phenyl]-amide (115 mg, 86%) was prepared from 1-[5-amino-2-(2-fluoro-


15

6-methoxy-phenoxy)-phenyl]-3-thiazol-2-yl-urea (94 mg, 0.25 mmol) following the general procedure L.

LC-MS (m/z): 533 ($M+1$)⁺; ¹H NMR (400 MHz, DMSO- d_6): δ 2.74 (s, 3H), 3.20 (s, 3H), 3.99 (s, 3H), 6.79 (d, 1H), 7.09 (dd, 1H), 7.34 (t, 1H), 7.44 (d, 1H), 7.48 (d, 1H), 7.67 (q, 1H), 7.78 (d, 1H), 8.01 (s, 1H), 8.36 (s, 1H), 8.46 (d, 1H), 9.88 (br, 1H), and 11.24 (br, 1H).

Example 389

1-[2-(2-Fluoro-6-methoxy-phenoxy)-5-methylamino-phenyl]-3-thiazol-2-yl-urea




10 1-[2-(2-Fluoro-6-methoxy-phenoxy)-5-methylamino-phenyl]-3-thiazol-2-yl-urea (85 mg, 88%) was prepared from 1-[5-amino-2-(2-fluoro-6-methoxy-phenoxy)-phenyl]-3-thiazol-2-yl-urea (94 mg, 0.25 mmol) following the general procedure O.

LC-MS (m/z): 389 ($M+1$)⁺; ¹H NMR (400 MHz, CDCl₃): δ 3.66 (s, 3H), 4.09 (s, 3H), 5.62 (br, 1H), 6.82 (d, 1H), 6.99 (s, 1H), 7.16 (m, 1H), 7.40 (d, 1H), 7.45-7.53 (m, 1H), 7.58 (d, 1H), 7.64 (d, 1H), 8.76 (d, 1H), 9.38 (br, 1H), and 11.27 (br, 1H).

Example 390

20 1-[2-(2-Fluoro-6-methoxy-phenoxy)-5-(1-methyl-3-thiazol-2-yl-ureido)-phenyl]-3-thiazol-2-yl-urea




1-[2-(2-Fluoro-6-methoxy-phenoxy)-5-(1-methyl-3-thiazol-2-yl-ureido)-phenyl]-3-thiazol-2-yl-urea (86 mg, 67%) was prepared from 4-(2-fluoro-6-methoxy-phenoxy)-N'1'-methyl-benzene-1,3-diamine (64 mg, 0.25 mmol) following the general procedure D.

- 5 LC-MS (m/z): 515 ($M+1$)⁺; ¹H NMR (400 MHz, CDCl₃): δ 3.32 (s, 3H), 3.73 (s, 3H), 5.40 (br, 1H), 6.46 (d, 2H), 6.54 (d, 1H), 6.74 (d, 1H), 6.80 (d, 1H), 7.08 (d, 2H), 7.22 (d, 1H), 7.33 (br, 1H), 8.38 (d, 1H), 9.03 (br, 1H), and 11.27 (br, 1H).

Example 391

- 10 1-[5-Dimethylamino-2-(2-fluoro-6-methoxy-phenoxy)-phenyl]-3-thiazol-2-yl-urea




1-[5-Dimethylamino-2-(2-fluoro-6-methoxy-phenoxy)-phenyl]-3-thiazol-2-yl-urea (85 mg, 85%) was prepared from 1-[5-amino-2-(2-fluoro-6-methoxy-phenoxy)-phenyl]-3-thiazol-2-yl-urea (94 mg, 0.25 mmol) following the general procedure O.

- 15 LC-MS (m/z): 403 ($M+1$)⁺; ¹H NMR (400 MHz, CDCl₃): δ 2.93 (s, 6H), 3.66 (s, 3H), 6.38 (d, 1H), 6.49 (t, 1H), 6.62-6.74 (m, 2H), 7.07 (t, 1H), 7.39 (d, 1H), 7.84 (s, 1H), 8.00 (br, 1H), 8.42 (br, 1H), and 11.52 (br, 1H).

Example 392

1-[2-(2-Fluoro-6-methoxy-phenoxy)-5-[(pyridin-3-ylmethyl)-amino]-phenyl]-3-thiazol-2-yl-urea




1-[2-(2-Fluoro-6-methoxy-phenoxy)-5-[(pyridin-3-ylmethyl)-amino]-phenyl]-3-thiazol-2-yl-urea (98 mg, 84%) was prepared from 1-[5-amino-2-(2-fluoro-6-methoxy-phenoxy)-phenyl]-3-thiazol-2-yl-urea (94 mg, 0.25 mmol) following the general procedure O.

LC-MS (m/z): 466 ($M+1$)⁺; ¹H NMR (400 MHz, CD₃OD): δ 3.16 (s, 2H), 3.62 (s, 3H), 6.42 (d, 1H), 6.68 (t, 1H), 6.78 (d, 1H), 6.88 (m, 1H), 7.05 (q, 1H), 7.21 (s, 1H), 7.41 (m, 1H), 7.52 (m, 1H), 7.81 (br, 1H), 8.00 (br, 1H), 8.16 (d, 1H), 8.42 (d, 1H), 8.56 (d, 1H), 8.70 (m, 1H), and 11.38 (br, 1H).

Example 393

1-[5-(Di-pyridin-2-yl-amino)-2-(2-fluoro-6-methoxy-phenoxy)-phenyl]-3-thiazol-2-yl-urea




1-[5-(Di-pyridin-2-yl-amino)-2-(2-fluoro-6-methoxy-phenoxy)-phenyl]-3-thiazol-2-yl-urea (111 mg, 84%) was prepared from 1-[5-amino-2-(2-fluoro-6-methoxy-phenoxy)-phenyl]-3-thiazol-2-yl-urea (94 mg, 0.25 mmol) following the general procedure O.

LC-MS (m/z): 529 ($M+1$)⁺; ¹H NMR (400 MHz, CD₃OD): δ 3.67 (s, 3H), 6.42 (d, 1H), 6.46 (d, 1H), 6.50 (d, 2H), 6.72 (q, 2H), 6.87 (m, 4H), 7.02-7.10 (m, 2H), 7.42 (m, 1H), 7.54 (m, 1H), 8.14 (d, 2H), 9.70 (br, 1H), and 11.58 (br, 1H).

Example 394

Propane-1-sulfonic acid [4-(2-fluoro-6-methoxy-phenoxy)-3-(3-thiazol-2-yl-ureido)-phenyl]-amide




Propane-1-sulfonic acid [4-(2-fluoro-6-methoxy-phenoxy)-3-(3-thiazol-2-yl-ureido)-phenyl]-amide (100 mg, 83%) was prepared from 1-[5-amino-2-(2-fluoro-6-methoxy-phenoxy)-phenyl]-3-thiazol-2-yl-urea (94 mg, 0.25 mmol) following the general procedure L.

LC-MS (m/z): 481 ($M+1$)⁺; ¹H NMR (400 MHz, Acetone- d_6): δ 1.02 (t, 3H), 1.84 (m, 2H), 3.08 (t, 2H), 3.82 (s, 3H), 6.56 (d, 1H), 6.88 (d, 1H), 6.96-7.00 (m, 1H), 7.24 (q, 1H), 7.34 (dd, 1H), 7.88 (s, 1H), 7.99 (br, 1H), 8.32 (d, 1H), 8.48 (d, 1H), 8.64 (br, 1H), and 11.12 (br, 1H).

Example 395

N-[4-(2-Fluoro-6-methoxy-phenoxy)-3-(3-thiazol-2-yl-ureido)-phenyl]-C-phenylmethanesulfonamide




N-[4-(2-Fluoro-6-methoxy-phenoxy)-3-(3-thiazol-2-yl-ureido)-phenyl]-C-phenylmethanesulfonamide (104 mg, 79%) was prepared from 1-[5-amino-2-(2-fluoro-6-methoxy-phenoxy)-phenyl]-3-thiazol-2-yl-urea (94 mg, 0.25 mmol) following the general procedure L.

5 LC-MS (m/z): 529 ($M+1$)⁺; ¹H NMR (400 MHz, CD₃OD): δ 3.36 (s, 2H), 3.83 (s, 3H), 6.54 (d, 1H), 6.86 (m, 2H), 6.94 (m, 2H), 7.22 (q, 1H), 7.38 (m, 5H), 7.42 (d, 1H), 7.96 (br, 1H), 8.06 (d, 1H), 9.34 (br, 1H), and 11.24 (br, 1H).

Example 396

10 N-{4-[4-(2-Fluoro-6-methoxy-phenoxy)-3-(3-thiazol-2-yl-ureido)-phenylsulfamoyl]-phenyl}-acetamide




15 N-{4-[4-(2-Fluoro-6-methoxy-phenoxy)-3-(3-thiazol-2-yl-ureido)-phenylsulfamoyl]-phenyl}-acetamide (113 mg, 79%) was prepared from 1-[5-amino-2-(2-fluoro-6-methoxy-phenoxy)-phenyl]-3-thiazol-2-yl-urea (94 mg, 0.25 mmol) following the general procedure L.

LC-MS (m/z): 572 ($M+1$)⁺; ¹H NMR (400 MHz, CD₃OD): δ 3.02 (s, 3H), 3.85 (s, 3H), 6.48 (d, 1H), 6.86 (m, 2H), 6.98 (br, 1H), 7.03 (m, 2H), 7.41 (m, 3H), 7.76 (m, 3H), 8.23 (d, 1H), 9.06 (br, 1H), 9.54 (br, 1H), and 11.24 (br, 1H).

5

Example 397

1-{2-(2-Fluoro-6-methoxy-phenoxy)-5-[(5-methyl-3H-imidazol-4-ylmethyl)-amino]-phenyl}-3-thiazol-2-yl-urea




10 1-{2-(2-Fluoro-6-methoxy-phenoxy)-5-[(5-methyl-3H-imidazol-4-ylmethyl)-amino]-phenyl}-3-thiazol-2-yl-urea (98 mg, 84%) was prepared from 1-[5-amino-2-(2-fluoro-6-methoxy-phenoxy)-phenyl]-3-thiazol-2-yl-urea (94 mg, 0.25 mmol) following the general procedure O.

LC-MS (m/z): 469 ($M+1$)⁺; ¹H NMR (400 MHz, CD₃OD): δ 2.29 (s, 3H), 2.55 (s, 2H), 3.68 (s, 3H), 6.46 (d, 1H), 6.86 (m, 2H), 6.96 (br, 1H), 7.08 (d, 1H), 7.41 (m, 1H), 7.49 (m, 15 2H), 7.54 (s, 1H), 7.64 (s, 1H), 7.72 (d, 1H), 9.46 (br, 1H), and 11.44 (br, 1H).

Example 398

4-Acetyl-N-[4-(2-fluoro-6-methoxy-phenoxy)-3-(3-thiazol-2-yl-ureido)-phenyl]-benzenesulfonamide




4-Acetyl-N-[4-(2-fluoro-6-methoxy-phenoxy)-3-(3-thiazol-2-yl-ureido)-phenyl]-benzenesulfonamide (98 mg, 84%) was prepared from 1-[5-amino-2-(2-fluoro-6-methoxy-phenoxy)-phenyl]-3-thiazol-2-yl-urea (94 mg, 0.25 mmol) following the general procedure L.

5 LC-MS (m/z): 557 ($M+1$)⁺; ¹H NMR (400 MHz, CD₃OD): δ 2.61 (s, 3H), 3.78 (s, 3H), 6.48 (d, 1H), 6.88 (m, 2H), 7.06 (m, 2H), 7.08 (d, 1H), 7.22 (br, 1H), 7.36 (d, 1H), 7.82 (m, 1H), 7.92 (d, 2H), 7.98 (t, 1H), 8.12 (d, 1H), 8.24 (br, 1H), and 11.88 (br, 1H).

Example 399

10 4-Cyano-N-[4-(2-fluoro-6-methoxy-phenoxy)-3-(3-thiazol-2-yl-ureido)-phenyl]benzenesulfonamide




15 4-Cyano-N-[4-(2-fluoro-6-methoxy-phenoxy)-3-(3-thiazol-2-yl-ureido)-phenyl]benzenesulfonamide (115 mg, 85%) was prepared from 1-[5-amino-2-(2-fluoro-6-methoxy-phenoxy)-phenyl]-3-thiazol-2-yl-urea (94 mg, 0.25 mmol) following the general procedure L.

LC-MS (m/z): 540 ($M+1$)⁺; ¹H NMR (400 MHz, CDCl₃): δ 3.76 (s, 3H), 6.52 (d, 1H), 6.80 (m, 1H), 6.88 (d, 1H), 7.08 (d, 2H), 7.12-7.18 (m, 1H), 7.30 (d, 1H), 7.38 (d, 1H), 7.54 (d, 1H), 7.71 (t, 1H), 7.91 (q, 1H), 8.01 (s, 1H), 8.08 (br, 1H), 8.22 (d, 1H), and 11.22 (br, 1H).

5

Example 400

N-[4-(2-Fluoro-6-methoxy-phenoxy)-3-(3-thiazol-2-yl-ureido)-phenyl]-4-isopropyl-benzenesulfonamide




10 N-[4-(2-Fluoro-6-methoxy-phenoxy)-3-(3-thiazol-2-yl-ureido)-phenyl]-4-isopropyl-benzenesulfonamide (103 mg, 74%) was prepared from 1-[5-amino-2-(2-fluoro-6-methoxy-phenoxy)-phenyl]-3-thiazol-2-yl-urea (94 mg, 0.25 mmol) following the general procedure L.

LC-MS (m/z): 557 ($M+1$)⁺; ¹H NMR (400 MHz, CDCl₃): δ 1.22 (d, 6H), 1.56 (m, 1H), 3.74 (s, 3H), 6.54 (d, 1H), 6.78 (m, 2H), 6.84 (d, 1H), 6.98 (m, 1H), 7.06 (d, 1H), 7.12-7.18 (m, 2H), 7.22 (d, 1H), 7.34 (d, 1H), 7.72 (d, 1H), 8.00 (br, 1H), 8.22 (d, 1H), 8.58 (br, 1H),
15 and 10.74 (br, 1H).

Example 401

20

N-{5-[4-(2-Fluoro-6-methoxy-phenoxy)-3-(3-thiazol-2-yl-ureido)-phenylsulfamoyl]-4-methyl-thiazol-2-yl}-acetamide




N-{5-[4-(2-Fluoro-6-methoxy-phenoxy)-3-(3-thiazol-2-yl-ureido)-phenylsulfamoyl]-4-methyl-thiazol-2-yl}-acetamide (121 mg, 82%) was prepared from 1-[5-amino-2-(2-fluoro-6-methoxy-phenoxy)-phenyl]-3-thiazol-2-yl-urea (94 mg, 0.25 mmol) following the general
5 procedure L.

LC-MS (*m/z*): 593 (*M*+1)⁺; ¹H NMR (400 MHz, CD₃OD): δ 2.32 (s, 3H), 3.00 (s, 3H), 3.77 (s, 3H), 4.04 (br, 1H), 6.48 (d, 1H), 6.84 (d, 1H), 6.88 (dd, 1H), 6.96 (d, 1H), 7.16-7.22 (m, 1H), 7.32 (d, 1H), 7.36 (d, 1H), 7.92 (d, 1H), 7.98 (br, 1H), 8.84 (br, 1H), and 10.74 (br, 1H).

10

Example 402

5-Bromo-6-chloro-pyridine-3-sulfonic acid [4-(2-fluoro-6-methoxy-phenoxy)-3-(3-thiazol-2-yl-ureido)-phenyl]-amide




5-Bromo-6-chloro-pyridine-3-sulfonic acid [4-(2-fluoro-6-methoxy-phenoxy)-3-(3-thiazol-2-yl-ureido)-phenyl]-amide (120 mg, 76%) was prepared from 1-[5-amino-2-(2-fluoro-6-methoxy-phenoxy)-phenyl]-3-thiazol-2-yl-urea (94 mg, 0.25 mmol) following the general procedure L.

LC-MS (m/z): 629 ($M+1$)⁺; ¹H NMR (400 MHz, CDCl₃): δ 3.78 (s, 3H), 6.56 (d, 1H), 6.88 (d, 1H), 6.78 (m, 1H), 6.88 (dd, 1H), 7.14 (d, 1H), 7.22 (d, 1H), 7.36 (d, 1H), 8.02 (d, 1H), 8.12 (br, 1H), 8.33 (d, 1H), 8.73 (d, 1H), 9.78 (br, 1H), and 10.74 (br, 1H).

10

Example 403

3,5-Dimethyl-isoxazole-4-sulfonic acid [4-(2-fluoro-6-methoxy-phenoxy)-3-(3-thiazol-2-yl-ureido)-phenyl]-amide



3,5-Dimethyl-isoxazole-4-sulfonic acid [4-(2-fluoro-6-methoxy-phenoxy)-3-(3-thiazol-2-yl-ureido)-phenyl]-amide (98 mg, 73%) was prepared from 1-[5-amino-2-(2-fluoro-6-


15

methoxy-phenoxy)-phenyl]-3-thiazol-2-yl-urea (94 mg, 0.25 mmol) following the general procedure L.

LC-MS (m/z): 534 ($M+1$)⁺; ¹H NMR (400 MHz, CDCl₃): δ 2.88 (s, 3H), 3.00 (s, 3H), 3.75 (s, 3H), 5.21 (br, 1H), 6.54 (d, 1H), 6.80-6.88 (m, 1H), 6.96 (m, 1H), 7.10-7.20 (m, 1H), 7.38 (d, 1H), 7.52 (d, 1H), 7.72 (br, 1H), 8.01 (d, 1H), 8.22 (d, 1H), and 10.74 (br, 1H).

Example 404

5-Chloro-1,3-dimethyl-1H-pyrazole-4-sulfonic acid [4-(2-fluoro-6-methoxy-phenoxy)-3-(3-thiazol-2-yl-ureido)-phenyl]-amide



10


5-Chloro-1,3-dimethyl-1H-pyrazole-4-sulfonic acid [4-(2-fluoro-6-methoxy-phenoxy)-3-(3-thiazol-2-yl-ureido)-phenyl]-amide (111 mg, 79%) was prepared from 1-[5-amino-2-(2-fluoro-6-methoxy-phenoxy)-phenyl]-3-thiazol-2-yl-urea (94 mg, 0.25 mmol) following the general procedure L.

LC-MS (m/z): 568 ($M+1$)⁺; ¹H NMR (400 MHz, CDCl₃): δ 2.32 (s, 3H), 2.63 (s, 3H), 3.74 (s, 3H), 6.48 (d, 1H), 6.76-6.84 (m, 1H), 6.94 (d, 1H), 7.06-7.18 (m, 1H), 7.22 (d, 1H), 7.38 (d, 1H), 7.52 (d, 1H), 7.94 (d, 1H), 8.08 (d, 1H), 8.20 (br, 1H), and 11.16 (br, 1H).

Example 405

6-Morpholin-4-yl-pyridine-3-sulfonic acid [4-(2-fluoro-6-methoxy-phenoxy)-3-(3-thiazol-2-yl-ureido)-phenyl]-amide

20




6-Morpholin-4-yl-pyridine-3-sulfonic acid [4-(2-fluoro-6-methoxy-phenoxy)-3-(3-thiazol-2-yl-ureido)-phenyl]-amide (115 mg, 76%) was prepared from 1-[5-amino-2-(2-fluoro-6-methoxy-phenoxy)-phenyl]-3-thiazol-2-yl-urea (94 mg, 0.25 mmol) following the
5 general procedure L.

LC-MS (*m/z*): 601 (*M*+1)⁺; ¹H NMR (400 MHz, CD₃OD): δ 2.15 (m, 4H), 3.60 (t, 2H), 3.70 (t, 2H), 3.76 (s, 3H), 6.42 (d, 1H), 6.78 (d, 2H), 6.84 (d, 1H), 6.96 (d, 1H), 7.04 (m, 2H), 7.20 (m, 1H), 7.34 (d, 1H), 7.79 (d, 1H), 8.02 (d, 1H), 8.39 (d, 1H), 8.52 (br, 1H), and 11.28 (br, 1H).

10

Example 406

2-Oxazol-5-yl-pyridine-3-sulfonic acid [4-(2-fluoro-6-methoxy-phenoxy)-3-(3-thiazol-2-yl-ureido)-phenyl]-amide




6-Oxazol-5-yl-pyridine-3-sulfonic acid [4-(2-fluoro-6-methoxy-phenoxy)-3-(3-thiazol-2-yl-ureido)-phenyl]-amide (96 mg, 66%) was prepared from 1-[5-amino-2-(2-fluoro-6-methoxy-phenoxy)-phenyl]-3-thiazol-2-yl-urea (94 mg, 0.25 mmol) following the general procedure L.

5 LC-MS (m/z): 583 ($M+1$)⁺; ¹H NMR (400 MHz, CD₃OD): δ 3.77 (s, 3H), 6.46 (d, 1H), 6.78-6.86 (m, 2H), 6.94 (d, 1H), 6.92-6.99 (m, 2H), 7.20 (q, 1H), 7.33 (dd, 1H), 7.55 (d, 1H), 7.80 (d, 1H), 7.86-7.92 (m, 2H), 8.05 (d, 1H), 8.18 (d, 1H), 9.88 (br, 1H), and 11.16 (br, 1H).

Example 407

10 2-(2-{3-[2-(2-Fluoro-6-methoxy-phenoxy)-5-methanesulfonylamino-phenyl]-ureido}-thiazol-4-yl)-N-methyl-acetamide




15 (2-{3-[2-(2-Fluoro-6-methoxy-phenoxy)-5-methanesulfonylamino-phenyl]-ureido}-thiazol-4-yl)-N-methyl-acetamide (47 mg, 87.9%) was prepared from (2-{3-[2-(2-fluoro-6-methoxy-phenoxy)-5-methanesulfonylamino-phenyl]-ureido}-thiazol-4-yl)-acetic acid (51 mg, 0.10 mmol) following the general procedure K.

LC-MS (m/z): 524 ($M+1$)⁺; ¹H NMR (400 MHz, CD₃OD): δ 2.93 (s, 3H), 3.48 (d, 3H), 3.56 (s, 2H), 4.02 (s, 3H), 4.82 (br, 1H), 6.76 (d, 1H), 7.04 (s, 1H), 7.09 (dd, 1H), 7.16 (t, 1H), 7.22 (d, 1H), 7.52 (q, 1H), 8.12 (br, 1H), 8.44 (d, 1H), 9.88 (br, 1H) and 11.84 (br, 1H).

20

Example 408

[4-(2-Fluoro-6-methoxy-phenoxy)-3-(3-thiazol-2-yl-ureido)-phenyl]-carbamic acid tert-butyl ester




[4-(2-Fluoro-6-methoxy-phenoxy)-3-(3-thiazol-2-yl-ureido)-phenyl]-carbamic acid tert-butyl ester (4.56 g, 96.2%) was prepared from 3-amino-4-(2-fluoro-6-methoxy-phenoxy)-phenyl]-carbamic acid tert-butyl ester (3.48 g, 10.0 mmol) following the general procedure D.

- 5 LC-MS (m/z): 475 ($M+1$)⁺; ¹H NMR (400 MHz, CD₃OD): δ 1.74 (s, 9H), 3.74 (s, 3H), 6.64 (d, 1H), 6.92 (dd, 1H), 7.02 (d, 1H), 7.14 (m, 1H), 7.34 (dd, 1H), 7.68 (dd, 1H), 7.88 (br, 1H), 8.32 (dd, 1H), 8.46 (d, 1H), 9.26 (br, 1H), 11.62 (br, 1H).


Example 409

- 10 1-(4-Methyl-2-propoxy-phenyl)-3-thiazol-2-yl-urea



- 3-Propoxy-4-nitrotoluene (0.79 g, 80%) was prepared from 1-propanol and 3-fluoro-4-nitrotoluene (0.77 g, 5.0 mmol) following the general procedure G. This was reduced to 4-methyl-2-propoxyaniline (0.54 g, 82%) following general procedure C. 1-(4-Methyl-2-propoxy-phenyl)-3-thiazol-2-yl-urea (220 mg, 75%) was prepared from 4-methyl-2-propoxyaniline (165 mg, 1.0 mmol) and 2-aminothiazole (100 mg, 1.0 mmol) following the general procedure D.


- LC-MS (m/z): 292 ($M+1$)⁺; ¹H NMR (400 MHz, DMSO- d_6): δ 1.03 (t, 3H), 1.80 (m, 2H), 2.25 (s, 3H), 3.98 (t, 2H), 6.70 (dd, 1H), 6.83 (d, 1H), 7.10 (d, 1H), 7.37 (d, 1H), 7.98 (d, 1H), 8.50 (br, 1H), 11.27 (br, 1H).
- 20

Example 410**1-(2-Butoxy-4-methyl-phenyl)-3-thiazol-2-yl-urea**

- 5 3-(1-Butoxy)-4-nitrotoluene (0.78 g, 75%) was prepared from 1-butanol and 3-fluoro-4-nitrotoluene (0.77 g, 5.0 mmol) following the general procedure G. This was reduced to 4-methyl-2-(1-butoxy)aniline (0.47 g, 70%) following general procedure C. 1-(2-Butoxy-4-methyl-phenyl)-3-thiazol-2-yl-urea (210 mg, 70%) was prepared from 4-methyl-2-(1-butoxy)aniline (179 mg, 1.0 mmol) and 2-aminothiazole (100 mg, 1.0 mmol) following the
- 10 general procedure D.

LC-MS (*m/z*): 306 (*M*+1)⁺; ¹H NMR (400 MHz, DMSO-*d*₆): δ 0.95 (t, 3H), 1.52 (m, 2H), 1.78 (m, 2H), 2.25 (s, 3H), 4.04 (t, 2H), 6.71 (dd, 1H), 6.86 (d, 1H), 7.11 (d, 1H), 7.36 (d, 1H), 7.96 (d, 1H), 8.50 (br, 1H), 11.26 (br, 1H).

15 **Example 411**

1-[4-Methyl-2-(3-methyl-butoxy)-phenyl]-3-thiazol-2-yl-urea


- 20 3-(3-Methyl-butoxy)-4-nitrotoluene (0.89 g, 80%) was prepared from 3-methylbutanol and 3-fluoro-4-nitrotoluene (0.77 g, 5.0 mmol) following the general procedure G. This was reduced to 4-methyl-2-(3-methylbutoxy)aniline (0.55 g, 72%) following general procedure C. 1-[4-Methyl-2-(3-methyl-butoxy)-phenyl]-3-thiazol-2-yl-urea (240 mg, 76%) was prepared

from 4-methyl-2-(3-methyl-butoxy)aniline (193 mg, 1.0 mmol) and 2-aminothiazole (100 mg, 1.0 mmol) following the general procedure D.

LC-MS (*m/z*): 320 (*M*+1)⁺; ¹H NMR (400 MHz, DMSO-*d*₆): δ 0.94 (d, 6H), 1.71 (m, 2H), 1.85 (m, 1H), 2.26 (s, 3H), 4.04 (t, 2H), 6.70 (d, 1H), 6.87 (s, 1H), 7.11 (d, 1H), 7.36 (d, 1H), 7.96 (d, 1H), 8.50 (br, 1H), 11.26 (br, 1H).

Example 412

1-(2-Allyloxy-4-methyl-phenyl)-3-thiazol-2-yl-urea




3-Allyloxy-4-nitrotoluene (0.67 g, 70%) was prepared from allyl alcohol and 3-fluoro-4-nitrotoluene (0.77 g, 5.0 mmol) following the general procedure G. This was reduced to 4-methyl-2-(allyloxy)aniline (0.40 g, 70%) following general procedure B. 1-(2-Allyloxy-4-methyl-phenyl)-3-thiazol-2-yl-urea (196 mg, 68%) was prepared from 4-methyl-2-(2-methylpropoxy)aniline (163 mg, 1.0 mmol) and 2-aminothiazole (100 mg, 1.0 mmol) following the general procedure D.

LC-MS (*m/z*): 290 (*M*+1)⁺; ¹H NMR (400 MHz, DMSO-*d*₆): δ 2.25 (s, 3H), 4.64 (d, 2H), 5.30 (m, 1H), 5.49 (m, 1H), 6.09 (m, 1H), 6.74 (dd, 1H), 6.87 (s, 1H), 7.10 (d, 1H), 7.36 (dd, 1H), 7.97 (d, 1H), 8.60 (br, 1H), 11.15 (br, 1H).

20

Example 413

1-[4-Methyl-2-(2-methyl-allyloxy)-phenyl]-3-thiazol-2-yl-urea




3-(2-Methyl-allyloxy)-4-nitrotoluene (0.78 g, 75%) was prepared from methallyl alcohol and 3-fluoro-4-nitrotoluene (0.77 g, 5.0 mmol) following the general procedure G.

This was reduced to 4-methyl-2-(2-methyl-allyloxy)aniline (0.47 g, 70%) following general procedure B. 1-[4-Methyl-2-(2-methyl-allyloxy)-phenyl]-3-thiazol-2-yl-urea (210 mg, 70%) was prepared from 4-methyl-2-(2-methyl-allyloxy)aniline (179 mg, 1.0 mmol) and 2-aminothiazole (100 mg, 1.0 mmol) following the general procedure D.

- 5 LC-MS (*m/z*): 304 (*M*+1)⁺; ¹H NMR (400 MHz, DMSO-*d*₆): δ 1.83 (s, 3H), 2.25 (s, 3H), 4.54 (s, 2H), 5.00 (s, 1H), 5.15 (s, 1H), 6.72 (d, 1H), 6.86 (s, 1H), 7.10 (d, 1H), 7.36 (d, 1H), 7.97 (d, 1H), 8.60 (br, 1H), 11.21 (br, 1H).

Example 414

- 10 1-[4-Methyl-2-(3-methyl-but-2-enyloxy)-phenyl]-3-thiazol-2-yl-urea




- 3-(3-methyl-but-2-enyloxy)-4-nitrotoluene (0.8 g, 72%) was prepared from 3-methylbut-2-en-1-ol and 3-fluoro-4-nitrotoluene (0.77 g, 5.0 mmol) following the general procedure G. This was reduced to 4-methyl-2-(3-methyl-but-2-enyloxy)aniline (0.43 g, 62%) following general procedure B. 1-[4-Methyl-2-(3-methyl-but-2-enyloxy)-phenyl]-3-thiazol-2-yl-urea (206 mg, 65%) was prepared from 4-methyl-2-(3-methyl-but-2-enyloxy)aniline (191 mg, 1.0 mmol) and 2-aminothiazole (100 mg, 1.0 mmol) following the general procedure D.

- 20 LC-MS (*m/z*): 318 (*M*+1)⁺; ¹H NMR (400 MHz, DMSO-*d*₆): δ 1.75 (d, 6H), 2.25 (s, 3H), 4.60 (d, 2H), 5.26 (t, 1H), 6.71 (d, 1H), 6.89 (s, 1H), 7.09 (d, 1H), 7.35 (d, 1H), 7.96 (dd, 1H), 8.56 (br, 1H), 11.17 (br, 1H).

Example 415

1-(2-Isopropoxy-4-methyl-phenyl)-3-thiazol-2-yl-urea




3-(2-Isopropoxy)-4-nitrotoluene (0.72 g, 74%) was prepared from 2-propanol and 3-fluoro-4-nitrotoluene (0.77 g, 5.0 mmol) following the general procedure G. This was reduced to 2-isopropoxy-4-methyl-aniline (0.42 g, 70%) following general procedure C. 1-(2-isopropoxy-4-methyl-phenyl)-3-thiazol-2-yl-urea (180 mg, 62%) was prepared from 2-isopropoxy-4-methyl-aniline (165 mg, 1.0 mmol) and 2-aminothiazole (100 mg, 1.0 mmol) following the general procedure D.

LC-MS (*m/z*): 292 (*M*+1)⁺; ¹H NMR (400 MHz, DMSO-*d*₆): δ 1.32 (d, 6H), 2.24 (s, 3H), 4.63 (m, 1H), 6.70 (dd, 1H), 6.88 (d, 1H), 7.10 (d, 1H), 7.37 (d, 1H), 7.97 (d, 1H), 8.60 (br, 1H), 11.27 (br, 1H).

EXAMPLE 416

1-(2-Cyclopentyloxy-4-methyl-phenyl)-3-thiazol-2-yl-urea




3-Cyclopentyloxy-4-nitrotoluene (0.68 g, 62%) was prepared from cyclopentanol and 3-fluoro-4-nitrotoluene (0.77 g, 5.0 mmol) following the general procedure G. This was reduced to 2-cyclopentyloxy-4-methyl-aniline (0.43 g, 70%) following general procedure C. 1-(2-Cyclopentyloxy-4-methyl-phenyl)-3-thiazol-2-yl-urea (220 mg, 70%) was prepared from 2-cyclopentyloxy-4-methyl-aniline (191 mg, 1.0 mmol) and 2-aminothiazole (100 mg, 1.0 mmol) following the general procedure D.

LC-MS (*m/z*): 318 (*M*+1)⁺; ¹H NMR (400 MHz, DMSO-*d*₆): δ 1.60 (m, 2H), 1.80 (m, 4H), 1.93 (m, 2H), 2.25 (s, 3H), 4.86 (m, 1H), 6.68 (dd, 1H), 6.81 (d, 1H), 7.10 (d, 1H), 7.37 (d, 1H), 7.97 (d, 1H), 8.40 (br, 1H), 11.31 (br, 1H).

Example 417

1-(2-Cyclopropylmethoxy-4-methyl-phenyl)-3-thiazol-2-yl-urea




3-Cyclopropylmethoxy-4-nitrotoluene (0.77 g, 75%) was prepared from cyclopropylmethanol and 3-fluoro-4-nitrotoluene (0.77 g, 5.0 mmol) following the general procedure G. This was reduced to 2-cyclopropylmethoxy-4-methyl-aniline (0.47 g, 71%) following general procedure C. 1-(2-Cyclopropylmethoxy-4-methyl-phenyl)-3-thiazol-2-yl-urea (218 mg, 72%) was prepared from 2-cyclopropylmethoxy-4-methyl-aniline (177 mg, 1.0 mmol) and 2-aminothiazole (100 mg, 1.0 mmol) following the general procedure D.

LC-MS (*m/z*): 304 (*M*+1)⁺; ¹H NMR (400 MHz, DMSO-*d*₆): δ 0.37 (m, 2H), 0.59 (m, 2H), 1.29 (m, 1H), 2.24 (s, 3H), 3.88 (d, 2H), 6.71 (d, 1H), 6.84 (d, 1H), 7.10 (d, 1H), 7.37 (d, 1H), 7.96 (d, 1H), 8.60 (br, 1H), 11.25 (br, 1H).

Example 418

1-[4-Methyl-2-(tetrahydro-pyran-2-ylmethoxy)-phenyl]-3-thiazol-2-yl-urea




4-Nitro-3-(tetrahydro-pyran-2-ylmethoxy)-toluene (0.80 g, 64%) was prepared from tetrahydropyran-2-ylmethanol and 3-fluoro-4-nitrotoluene (0.77 g, 5.0 mmol) following the general procedure G. This was reduced to 4-methyl-2-(tetrahydro-pyran-2-ylmethoxy)-aniline (0.50 g, 71%) following general procedure C. 1-[4-Methyl-2-(tetrahydro-pyran-2-ylmethoxy)-phenyl]-3-thiazol-2-yl-urea (246 mg, 71%) was prepared from 4-methyl-2-(tetrahydro-pyran-2-ylmethoxy)-aniline (221 mg, 1.0 mmol) and 2-aminothiazole (100 mg, 1.0 mmol) following the general procedure D.

LC-MS (m/z): 348 ($M+1$)⁺; ¹H NMR (400 MHz, DMSO- d_6): δ 1.34 (m, 1H), 1.50 (m, 2H), 1.55 (m, 1H), 1.75 (d, 1H), 1.83 (m, 1H), 2.25 (s, 3H), 3.43 (m, 1H), 3.90 (m, 2H), 4.01 (m, 1H), 6.72 (d, 1H), 6.86 (d, 1H), 7.10 (d, 1H), 7.37 (d, 1H), 7.96 (d, 1H), 8.50 (br, 1H), 11.28 (br, 1H).

5

Example 419

1-[4-Methyl-2-(tetrahydro-furan-2-ylmethoxy)-phenyl]-3-thiazol-2-yl-urea




4-Nitro-3-(tetrahydro-furan-2-ylmethoxy)-toluene (0.88 g, 75%) was prepared from tetrahydro-furan-2-ylmethanol and 3-fluoro-4-nitrotoluene (0.77 g, 5.0 mmol) following the general procedure G. This was reduced to 4-methyl-2-(tetrahydro-furan-2-ylmethoxy)-aniline (0.50 g, 65%) following general procedure C. 1-[4-Methyl-2-(tetrahydro-furan-2-ylmethoxy)-phenyl]-3-thiazol-2-yl-urea (236 mg, 71%) was prepared from 4-methyl-2-(tetrahydro-furan-2-ylmethoxy)-aniline (207 mg, 1.0 mmol) and 2-aminothiazole (100 mg, 1.0 mmol) following the general procedure D.

15

LC-MS (m/z): 334 ($M+1$)⁺.**Example 420**

1-(2-Cyclopentylmethoxy-4-methyl-phenyl)-3-thiazol-2-yl-urea



20


3-Cyclopentylmethoxy-4-nitrotoluene (0.82 g, 70%) was prepared from cyclopentylmethanol and 3-fluoro-4-nitrotoluene (0.77 g, 5.0 mmol) following the general

procedure G. This was reduced to 2-cyclopentylmethoxy-4-methyl-aniline (0.58 g, 81%) following general procedure C. 1-(2-Cyclopentylmethoxy-4-methyl-phenyl)-3-thiazol-2-yl-urea (225 mg, 68%) was prepared from 2-cyclopentylmethoxy-4-methyl-aniline (205 mg, 1.0 mmol) and 2-aminothiazole (100 mg, 1.0 mmol) following the general procedure D.

5 LC-MS (m/z): 332 ($M+1$)⁺;

Example 421

{2-[3-(2-Cyclopropylmethoxy-4-methyl-phenyl)-ureido]-thiazol-4-yl}-acetic acid



10 {2-[3-(2-Cyclopropylmethoxy-4-methyl-phenyl)-ureido]-thiazol-4-yl}-acetic acid ethyl ester (253 mg, 65%) was prepared from 2-cyclopropylmethoxy-4-methyl-aniline (177 mg, 1.0 mmol) and (2-amino-thiazol-4-yl)-acetic acid ethyl ester (186 mg, 1.0 mmol) following the general procedure D.


LC-MS (m/z): 390 ($M+1$)⁺, ¹H NMR (400 MHz, DMSO- d_6): δ 0.36 (m, 2H), 0.59 (m, 2H), 1.18 (t, 3H), 1.28 (m, 1H), 2.24 (s, 3H), 3.64 (s, 2H), 3.88 (d, 2H), 4.08 (q, 2H), 6.71 (dd, 1H), 6.83 (d, 1H), 6.86 (d, 1H), 7.94 (d, 1H), 8.40 (br, 1H), 11.36 (br, 1H).

{2-[3-(2-Cyclopropylmethoxy-4-methyl-phenyl)-ureido]-thiazol-4-yl}-acetic acid (165 mg, 92%) was prepared from {2-[3-(2-cyclopropylmethoxy-4-methyl-phenyl)-ureido]-thiazol-4-yl}-acetic acid ethyl ester (195 mg, 0.5 mmol) following the general procedure J.

20 LC-MS (m/z): 362 ($M+1$)⁺.

Example 422

{2-[3-(2-Cyclopentylloxy-4-methyl-phenyl)-ureido]-thiazol-4-yl}-acetic acid




{2-[3-(2-Cyclopentyloxy-4-methyl-phenyl)-ureido]-thiazol-4-yl}-acetic acid ethyl ester (250 mg, 62%) was prepared from 2-cyclopentyloxy-4-methyl-aniline (191 mg, 1.0 mmol) and (2-amino-thiazol-4-yl)-acetic acid ethyl ester (186 mg, 1.0 mmol) following the general procedure D. {2-[3-(2-Cyclopentyloxy-4-methyl-phenyl)-ureido]-thiazol-4-yl}-acetic acid (170 mg, 91%) was prepared from {2-[3-(2-cyclopropylmethoxy-4-methyl-phenyl)-ureido]-thiazol-4-yl}-acetic acid ethyl ester (200 mg, 0.5 mmol) following the general procedure J.

LC-MS (m/z): 376 ($M+1$)⁺.

10

Example 423

1-(2-Isobutoxy-4-methanesulfonylmethyl-phenyl)-3-thiazol-2-yl-urea




2-Isobutoxy-4-methanesulfonylmethyl-1-nitro-benzene (0.93 g, 65%) was prepared from 2-methylpropanol and 2-fluoro-4-methanesulfonylmethyl-1-nitro-benzene (1.16 g, 5.0 mmol) following the general procedure G. This was reduced to 2-isobutoxy-4-methanesulfonylmethyl-aniline (0.46 g, 56%) following general procedure B. 1-(2-Isobutoxy-4-methanesulfonylmethyl-phenyl)-3-thiazol-2-yl-urea (250 mg, 65%) was prepared from 2-isobutoxy-4-methanesulfonylmethyl-aniline (257 mg, 1.0 mmol) and 2-aminothiazole (100 mg, 1.0 mmol) following the general procedure D.

20

LC-MS (m/z): 384 ($M+1$)⁺; ¹H NMR (400 MHz, DMSO- d_6): δ 1.06 (d, 6H), 2.14 (m, 1H), 2.88 (s, 3H), 3.11 (s, 2H), 4.39 (d, 2H), 6.19 (s, 1H), 6.89 (d, 1H), 7.14 (s, 1H), 7.39 (d, 1H), 8.13 (d, 1H), 8.50 (br, 1H), 11.24 (br, 1H).

Example 424

(2-{2-[3-(2-Isobutoxy-4-methyl-phenyl)-ureido]-thiazol-4-yl}-acetylamino)-acetic acid




- 5 (2-{2-[3-(2-Isobutoxy-4-methyl-phenyl)-ureido]-thiazol-4-yl}-acetylamino)-acetic acid (90 mg, 86%) was prepared from (2-{2-[3-(2-isobutoxy-4-methyl-phenyl)-ureido]-thiazol-4-yl}-acetylamino)-acetic acid methyl ester (108 mg, 0.25 mmol) following the general procedure J. (2-{2-[3-(2-Isobutoxy-4-methyl-phenyl)-ureido]-thiazol-4-yl}-acetylamino)-acetic acid methyl ester (150 mg, 70%) was in turn prepared from {2-[3-(2-isobutoxy-4-methyl-phenyl)-ureido]-thiazol-4-yl}-acetic acid (181 mg, 0.5 mmol) and glycine methyl ester hydrochloride following the general procedure K.

LC-MS (*m/z*): 421 (*M*+1)⁺.

Example 425

- 15 1-(5-Formyl-2-isobutoxy-phenyl)-3-thiazol-2-yl-urea




4-(2-Methylpropoxy)-3-nitrobenzaldehyde (0.83 g, 75%) was prepared from 2-methylpropanol (0.46 ml, 5.0 mmol) and 4-fluoro-3-nitrobenzaldehyde (0.77 g, 5.0 mmol) following the general procedure G. This was reduced to 5-formyl-2-(2-methylpropoxy)-aniline (0.49 g,

68%) following general procedure B. 1-(5-Formyl-2-isobutoxy-phenyl)-3-thiazol-2-yl-urea (208 mg, 65%) was prepared from 5-formyl-2-(2-methylpropoxy)aniline (193 mg, 1.0 mmol) and 2-aminothiazole (100 mg, 1.0 mmol) following the general procedure D.

LC-MS (m/z): 320 ($M+1$)⁺; ¹H NMR (400 MHz, DMSO- d_6): δ 1.06 (d, 6H), 2.16 (m, 1H), 3.96 (d, 2H), 7.16 (d, 1H), 7.25 (d, 1H), 7.40 (dd, 1H), 7.61 (d, 1H), 8.69 (s, 1H), 9.86 (s, 1H), 10.75 (br, 1H), 11.51 (br, 1H).

Example 426

2-{2-[3-(2-Isobutoxy-4-methyl-phenyl)-ureido]-thiazol-4-yl}-N-(2-methoxy-ethyl)-acetamide




2-{2-[3-(2-Isobutoxy-4-methyl-phenyl)-ureido]-thiazol-4-yl}-N-(2-methoxy-ethyl)-acetamide (130 mg, 62%) was prepared from {2-[3-(2-isobutoxy-4-methyl-phenyl)-ureido]-thiazol-4-yl}-acetic acid (181 mg, 0.5 mmol) and 2-methoxyethylamine following the general procedure K.

LC-MS (m/z): 421 ($M+1$)⁺.

Example 427

1-(5-Fluoro-2-isobutoxy-4-methyl-phenyl)-3-thiazol-2-yl-urea




2-Fluoro-5-(2-methylpropoxy)-4-nitrotoluene (0.73 g, 65%) was prepared from 2-methylpropanol (0.46 ml, 5.0 mmol) and 2,5-difluoro-4-nitrotoluene (0.86 g, 5.0 mmol) following the general procedure G. This was reduced to 5-fluoro-4-methyl-2-(2-methylpropoxy)-aniline (0.45 g, 70%) following general procedure C. 1-(5-Fluoro-2-isobutoxy-4-methyl-phenyl)-3-thiazol-2-yl-urea (205 mg, 64%) was prepared from 5-fluoro-4-methyl-2-(2-methylpropoxy)-aniline (197 mg, 1.0 mmol) and 2-aminothiazole (100 mg, 1.0 mmol) following the general procedure D.

LC-MS (*m/z*): 324 (*M*+1)⁺; ¹H NMR (400 MHz, DMSO-*d*₆): δ 1.03 (d, 6H), 2.16 (m, 1H), 2.17 (s, 3H), 3.80 (d, 2H), 6.92 (d, 1H), 7.14 (d, 1H), 7.38 (d, 1H), 7.93 (d, 1H), 8.50 (br, 1H), 11.44 (br, 1H).

Example 428

1-(5-Fluoro-2-isobutoxy-phenyl)-3-thiazol-2-yl-urea



15


5-Fluoro-2-(2-methylpropoxy)-4-nitrobenzene (0.83 g, 78%) was prepared from 2-methylpropanol (0.46 ml, 5.0 mmol) and 2,5-difluoro-4-nitrobenzene (0.80 g, 5.0 mmol) following the general procedure G. This was reduced to 5-fluoro-2-(2-methylpropoxy)-aniline (0.58 g, 81%) following general procedure C. 1-(5-Fluoro-2-isobutoxy-phenyl)-3-thiazol-2-yl-urea (220 mg, 72%) was prepared from 5-fluoro-2-(2-methylpropoxy)-aniline (183 mg, 1.0 mmol) and 2-aminothiazole (100 mg, 1.0 mmol) following the general procedure D.

20

LC-MS (m/z): 310 ($M+1$)⁺; ¹H NMR (400 MHz, DMSO- d_6): δ 1.03 (d, 6H), 2.11 (m, 1H), 3.80 (d, 2H), 6.79 (m, 1H), 6.99 (m, 1H), 7.15 (d, 1H), 7.39 (d, 1H), 8.01 (dd, 1H), 8.60 (br, 1H), 11.52 (br, 1H).

5 Example 429

1-(2-Isobutylsulfanyl-4-methyl-phenyl)-3-thiazol-2-yl-urea




- 3-(Isobutylsulfanyl)-4-nitrotoluene (0.75 g, 67%) was prepared from 2-methyl-propanethiol and 3-fluoro-4-nitrotoluene (0.77 g, 5.0 mmol) following the general procedure
- 10 A. This was reduced to 2-isobutylsulfanyl-4-methyl-aniline (0.46 g, 63%) following general procedure B. 1-(2-Isobutylsulfanyl-4-methyl-phenyl)-3-thiazol-2-yl-urea (218 mg, 68%) was prepared from 2-isobutylsulfanyl-4-methyl-aniline (195 mg, 1.0 mmol) and 2-aminothiazole (100 mg, 1.0 mmol) following the general procedure D.

- LC-MS (m/z): 318 ($M+1$)⁺, ¹H NMR (400 MHz, DMSO- d_6): δ 0.85 (m, 6H), 1.40 (m, 1H), 2.26 (s, 3H), 2.80 (m, 2H), 7.11 (s, 1H), 7.29 (s, 1H), 7.38 (d, 1H), 7.89 (d, 1H), 8.65 (br, 1H), 11.28 (br, 1H).

Example 430

{2-[3-(2-Isobutylsulfanyl-4-methyl-phenyl)-ureido]-thiazol-4-yl}-acetic acid ethyl ester



20


{2-[3-(2-Isobutylsulfanyl-4-methyl-phenyl)-ureido]-thiazol-4-yl}-acetic acid ethyl ester (276 mg, 68%) was prepared from 2-isobutylsulfanyl-4-methyl-aniline (195 mg, 1.0

mmol) and ethyl 2-amino-4-thiazolylacetate (186 mg, 1.0 mmol) following the general procedure D. Hydrolysis of this ester following general procedure J gave (2-[3-(2-isobutylsulfanyl-4-methyl-phenyl)-ureido]-thiazol-4-yl)-acetic acid (230 mg, 90%).

LC-MS (m/z): 380 ($M+1$)⁺, ¹H NMR (400 MHz, DMSO- d_6): δ 0.85 (m, 6H), 1.49 (m, 1H), 2.26 (s, 3H), 2.83 (m, 2H), 3.56 (s, 2H), 6.85 (s, 1H), 7.09 (d, 1H), 7.29 (d, 1H), 7.89 (d, 1H), 8.65 (br, 1H), 11.28 (br, 2H).

Example 431

1-(2-Cyclopentanesulfinyl-4-methyl-phenyl)-3-thiazol-2-yl-urea



10


3-Cyclopentanesulfinyl-4-nitrotoluene (0.47 g, 2.0 mmol) was oxidized to 3-cyclopentanesulfonyl-4-nitrotoluene (0.46 g, 91%) following general procedure R (using one equivalent of *m*-CPBA). 3-Cyclopentanesulfonyl-4-nitrotoluene was reduced to 2-cyclopentanesulfonyl-4-methyl-aniline (0.33 g, 81%) following general procedure C. 1-(2-cyclopentanesulfonyl-4-methyl-phenyl)-3-thiazol-2-yl-urea (225 mg, 65%) was prepared from 2-cyclopentanesulfonyl-4-methyl-aniline (223 mg, 1.0 mmol) and 2-aminothiazole (100 mg, 1.0 mmol) following the general procedure D.

LC-MS (m/z): 350 ($M+1$)⁺; ¹H NMR (400 MHz, DMSO- d_6): δ 1.58 (m, 6H), 1.78 (m, 1H), 1.94 (m, 1H), 2.34 (s, 3H), 3.43 (m, 1H), 7.11 (d, 1H), 7.33 (dd, 1H), 7.36 (dd, 1H), 7.43 (s, 1H), 7.80 (d, 1H), 8.93 (br, 1H), 11.25 (br, 1H).

20

Example 432

1-(2-Cyclopentanesulfonyl-4-methyl-phenyl)-3-thiazol-2-yl-urea




3-Cyclopentanesulfonyl-4-nitrotoluene (0.83 g, 70%) was prepared from cyclopentyl mercaptan and 3-fluoro-4-nitrotoluene (0.77 g, 5.0 mmol) following the general procedure A. This was oxidized to 3-cyclopentanesulfonyl-4-nitrotoluene (0.84 g, 90%) following general

5 procedure R. 3-Cyclopentanesulfonyl-4-nitrotoluene was reduced to 2-cyclopentanesulfonyl-4-methyl-aniline (0.53 g, 70%) following general procedure C. 1-(2-Cyclopentanesulfonyl-4-methyl-phenyl)-3-thiazol-2-yl-urea (225 mg, 62%) was prepared from 2-cyclopentanesulfonyl-4-methyl-aniline (239 mg, 1.0 mmol) and 2-aminothiazole (100 mg, 1.0 mmol) following the general procedure D.

10 LC-MS (*m/z*): 366 (*M*+1)⁺; ¹H NMR (400 MHz, DMSO-*d*₆): δ 1.58 (m, 2H), 1.78 (m, 2H), 1.86 (m, 2H), 2.04 (m, 2H), 2.38 (s, 3H), 3.54 (m, 1H), 6.94 (d, 1H), 7.43 (dd, 1H), 7.61 (d, 1H), 7.67 (d, 1H), 8.18 (d, 1H), 9.20 (br, 1H), 11.25 (br, 1H).

Example 433

15 {2-[3-(2-Cyclopentanesulfonyl-4-methyl-phenyl)-ureido]-thiazol-4-yl}-acetic acid ethyl ester




20 {2-[3-(2-Cyclopentanesulfonyl-4-methyl-phenyl)-ureido]-thiazol-4-yl}-acetic acid ethyl ester (305 mg, 68%) was prepared from 2-cyclopentanesulfonyl-4-methyl-aniline (239 mg, 1.0 mmol) and ethyl 2-amino-4-thiazolylacetate (186 mg, 1.0 mmol) following the general procedure D.

LC-MS (*m/z*): 452 (*M*+1)⁺, ¹H NMR (400 MHz, DMSO-*d*₆): δ 1.19 (t, 3H), 1.58 (m, 4H), 1.84 (m, 4H), 2.35 (s, 3H), 3.65 (s, 2H), 3.77 (m, 1H), 4.08 (q, 2H), 6.91 (s, 1H), 7.52 (d, 1H), 7.63 (s, 1H), 8.00 (d, 1H), 8.91 (br, 1H), 11.89 (br, 1H).

25 Example 434

{2-[3-(2-Cyclopentanesulfonyl-4-methyl-phenyl)-ureido]-thiazol-4-yl}-acetic acid




Following the general procedure J, {2-[3-(2-cyclopentanesulfonyl-4-methyl-phenyl)-ureido]-thiazol-4-yl)-acetic acid ethyl ester (225 mg, 0.5 mmol) was hydrolyzed to afford {2-[3-(4-methyl-2-[2-methylpropoxy]phenyl)-ureido]-thiazol-4-yl)-acetic acid (193 mg, 92%).

- 5 LC-MS (*m/z*): 424 (*M*+1)⁺, ¹H NMR (400 MHz, DMSO-*d*₆): δ 1.60 (m, 4H), 1.84 (m, 4H), 2.36 (s, 3H), 3.57 (s, 2H), 3.77 (m, 1H), 6.87 (s, 1H), 7.52 (d, 1H), 7.63 (s, 1H), 8.01 (d, 1H), 8.95 (br, 1H), 12.32 (br, 2H).

Example 435

- 10 2-[2-[3-(2-Cyclopentanesulfonyl-4-methyl-phenyl)-ureido]-thiazol-4-yl]-N-methyl-acetamide




- 15 2-[2-[3-(2-Cyclopentanesulfonyl-4-methyl-phenyl)-ureido]-thiazol-4-yl]-N-methyl-acetamide (30 mg, 68%) was prepared from {2-[3-(4-methyl-2-[2-methylpropoxy]phenyl)-ureido]-thiazol-4-yl)-acetic acid (42 mg, 0.1 mmol) and methylamine following the general procedure K.

LC-MS (*m/z*): 437 (*M*+1)⁺.

Example 436

2-{2-[3-(2-Cyclopentanesulfonyl-4-methyl-phenyl)-ureido]-thiazol-4-yl}-N-(2-methoxy-ethyl)-acetamide




2-{2-[3-(2-Cyclopentanesulfonyl-4-methyl-phenyl)-ureido]-thiazol-4-yl}-N-methyl-
5 acetamide (30 mg, 68%) was prepared from {2-[3-(4-methyl-2-[2-methylpropoxy]phenyl)-
ureido]-thiazol-4-yl}-acetic acid (42 mg, 0.1 mmol) and 2-methoxyethylamine following the
general procedure K.

LC-MS (*m/z*): 480 (*M*+1)⁺, ¹H NMR (400 MHz, DMSO-*d*₆): δ 1.60 (m, 4H), 1.84 (m,
4H), 2.36 (s, 3H), 3.21 (m, 4H), 3.24 (s, 3H), 3.43 (s, 2H), 3.76 (m, 1H), 6.81 (s, 1H), 7.52 (d,
10 1H), 7.63 (s, 1H), 8.05 (br, 2H), 8.95 (br, 1H), 11.85 (br, 1H).

Example 437

1-(2-Cyclopentylamino-4-methyl-phenyl)-3-thiazol-2-yl-urea




15 2-Cyclopentylamino-4-methyl-aniline (0.64 g, 68%) was prepared from
cyclopentylamine (1.0 ml, 10.0 mmol) and 3-fluoro-4-nitrotoluene (0.77 g, 5.0 mmol) following
the general procedure W. 1-(2-Cyclopentylamino-4-methyl-phenyl)-3-thiazol-2-yl-urea (195
mg, 62%) was prepared from 2-cyclopentylamino-4-methyl-aniline (190 mg, 1.0 mmol) and 2-
aminothiazole (100 mg, 1.0 mmol) following the general procedure D.

LC-MS (m/z): 317 ($M+1$)⁺, ¹H NMR (400 MHz, CDCl₃): δ 1.46 (m, 2H), 1.59 (m, 2H), 1.68 (m, 2H), 2.01 (m, 2H), 2.32 (s, 3H), 3.78 (m, 1H), 6.51 (d, 1H), 6.56 (s, 1H), 6.88 (d, 1H), 7.07 (br, 1H), 7.31 (d, 1H), 9.30 (br, 1H), 10.72 (br, 1H).

5 **Example 438**

1-(2-Isobutylamino-4-methyl-phenyl)-3-thiazol-2-yl-urea




2-Isobutylamino-4-methyl-aniline (0.61 g, 69%) was prepared from isobutylamine (0.73 g, 10.0 mmol) and 3-fluoro-4-nitrotoluene (0.77 g, 5.0 mmol) following the general
10 procedure W. 1-(2-Isobutylamino-4-methyl-phenyl)-3-thiazol-2-yl-urea (196 mg, 65%) was prepared from 2-Isobutylamino-4-methyl-aniline (178 mg, 1.0 mmol) and 2-aminothiazole (100 mg, 1.0 mmol) following the general procedure D.

LC-MS (m/z): 305 ($M+1$)⁺, ¹H NMR (400 MHz, CDCl₃): δ 0.95 (d, 6H), 1.88 (m, 1H), 2.31 (s, 3H), 2.92 (d, 2H), 4.15 (br, 1H), 6.51 (m, 2H), 6.87 (d, 1H), 7.08 (m, 1H), 7.31 (d,
15 1H), 9.30 (br, 1H), 10.72 (br, 1H).

Example 439

1-[4-Methyl-2-(methyl-propyl-amino)-phenyl]-3-thiazol-2-yl-urea




4-Methyl-2-(methyl-propyl-amino)-aniline (0.56 g, 63%) was prepared from N-methyl-propylamine (0.73 g, 10.0 mmol) and 3-fluoro-4-nitrotoluene (0.77 g, 5.0 mmol) following the general procedure W. 1-[4-Methyl-2-(methyl-propyl-amino)-phenyl]-3-thiazol-
20 2-yl-urea (215 mg, 71%) was prepared from 4-methyl-2-(methyl-propyl-amino)-aniline

(178 mg, 1.0 mmol) and 2-aminothiazole (100 mg, 1.0 mmol) following the general procedure D.

LC-MS (m/z): 305 ($M+1$)⁺, ¹H NMR (400 MHz, DMSO- d_6): δ 0.85 (t, 3H), 1.40 (m, 2H), 2.24 (s, 3H), 2.54 (s, 3H), 2.79 (t, 2H), 6.87 (dd, 1H), 7.02 (d, 1H), 7.09 (d, 1H), 7.37 (d, 1H),
5 7.99 (d, 1H), 8.80 (br, 1H), 11.36 (br, 1H).

Example 440

1-[2-(Butyl-methyl-amino)-4-methyl-phenyl]-3-thiazol-2-yl-urea




10 4-Methyl-2-(butyl-methyl-amino)-aniline (0.62 g, 65%) was prepared from N-methyl-butylamine (0.87 g, 10.0 mmol) and 3-fluoro-4-nitrotoluene (0.77 g, 5.0 mmol) following the general procedure W. 1-[2-(Butyl-methyl-amino)-4-methyl-phenyl]-3-thiazol-2-yl-urea (220 mg, 69%) was prepared from 4-methyl-2-(butyl-methyl-amino)-aniline (192 mg, 1.0 mmol) and 2-aminothiazole (100 mg, 1.0 mmol) following the general procedure
15 D.

LC-MS (m/z): 319 ($M+1$)⁺, ¹H NMR (400 MHz, DMSO- d_6): δ 0.82 (t, 3H), 1.27 (m, 2H), 1.38 (m, 2H), 2.24 (s, 3H), 2.54 (s, 3H), 2.82 (t, 2H), 6.87 (d, 1H), 7.02 (s, 1H), 7.09 (d, 1H), 7.37 (d, 1H), 7.97 (d, 1H), 8.80 (br, 1H), 11.37 (br, 1H).

20 Example 441

1-(2-Diethylamino-4-methyl-phenyl)-3-thiazol-2-yl-urea




2-(Diethyl-amino)-4-methyl-aniline (0.63 g, 71%) was prepared from diethylamine (0.73 g, 10.0 mmol) and 3-fluoro-4-nitrotoluene (0.77 g, 5.0 mmol) following the general procedure W. 1-(2-Diethylamino-4-methyl-phenyl)-3-thiazol-2-yl-urea (188 mg, 62%) was prepared from 2-(diethyl-amino)-4-methyl-aniline (178 mg, 1.0 mmol) and 2-aminothiazole (100 mg, 1.0 mmol) following the general procedure D.

LC-MS (m/z): 305 ($M+1$)⁺, ¹H NMR (400 MHz, DMSO- d_6): δ 0.89 (t, 6H), 2.24 (s, 3H), 2.90 (q, 4H), 6.92 (dd, 1H), 7.04 (d, 1H), 7.09 (d, 1H), 7.37 (d, 1H), 8.04 (d, 1H), 9.00 (br, 1H), 11.44 (br, 1H).

10 Example 442

1-[2-(Cyclopropylmethyl-propyl-amino)-4-methyl-phenyl]-3-thiazol-2-yl-urea




2-(Cyclopropylmethyl-propyl-amino)-4-methyl-aniline (0.68 g, 63%) was prepared from N-propyl-cyclopropanemethylamine (1.13 g, 10.0 mmol) and 3-fluoro-4-nitrotoluene (0.77 g, 5.0 mmol) following the general procedure W. 1-[4-Methyl-2-(methyl-propyl-amino)-phenyl]-3-thiazol-2-yl-urea (210 mg, 61%) was prepared from 4-methyl-2-(methyl-propyl-amino)-aniline (218 mg, 1.0 mmol) and 2-aminothiazole (100 mg, 1.0 mmol) following the general procedure D.

LC-MS (m/z): 345 ($M+1$)⁺, ¹H NMR (400 MHz, DMSO- d_6): δ -0.24 (m, 2H), 0.30 (m, 2H), 0.80 (m, 4H), 1.32 (m, 2H), 2.24 (s, 3H), 2.69 (d, 2H), 2.91 (t, 2H), 6.90 (dd, 1H), 7.07 (m, 2H), 7.37 (d, 1H), 8.01 (m, 1H), 9.00 (br, 1H), 11.51 (br, 1H).

Example 443

1-[2-(Isobutyl-methyl-amino)-4-methyl-phenyl]-3-thiazol-2-yl-urea




2-(Isobutyl-methyl-amino)-4-methyl-aniline (0.62 g, 65%) was prepared from N-isobutyl-methylamine (0.87 g, 10.0 mmol) and 3-fluoro-4-nitrotoluene (0.77 g, 5.0 mmol) following the general procedure W. 1-[2-(Isobutyl-methyl-amino)-4-methyl-phenyl]-3-thiazol-2-yl-urea (216 mg, 68%) was prepared from 2-(isobutyl-methyl-amino)-4-methyl-aniline (192 mg, 1.0 mmol) and 2-aminothiazole (100 mg, 1.0 mmol) following the general procedure D.

LC-MS (m/z): 319 ($M+1$)⁺, ¹H NMR (400 MHz, CDCl₃): δ 0.86 (d, 6H), 1.68 (m, 1H), 2.31 (s, 3H), 2.58 (s, 3H), 2.66 (d, 2H), 6.88 (d, 1H), 6.96 (m, 2H), 7.44 (d, 1H), 8.12 (d, 1H), 9.20 (br, 1H), 11.36 (br, 1H).

Example 444

(2-{3-[2-(Isobutyl-methyl-amino)-4-methyl-phenyl]-ureido}-thiazol-4-yl)-acetic acid ethyl ester




(2-{3-[2-(Isobutyl-methyl-amino)-4-methyl-phenyl]-ureido}-thiazol-4-yl)-acetic acid ethyl ester (290 mg, 72%) was prepared from 2-(isobutyl-methyl-amino)-4-methyl-aniline (192 mg, 1.0 mmol) and ethyl 2-amino-4-thiazolylacetate (186 mg, 1.0 mmol) following the general procedure D. Hydrolysis of this ester following the general procedure J gave (2-{3-[2-(isobutyl-methyl-amino)-4-methyl-phenyl]-ureido}-thiazol-4-yl)-acetic acid (250 mg, 92%).

LC-MS (m/z): 377 ($M+1$)⁺, ¹H NMR (400 MHz, DMSO- d_6): δ 0.85 (d, 6H), 1.60 (m, 1H), 2.21 (s, 3H), 2.66 (m, 5H), 3.53 (s, 2H), 6.81 (s, 1H), 6.85 (d, 1H), 6.99 (s, 1H), 7.95 (d, 1H), 8.60 (br, 1H), 11.55 (br, 1H), 12.35 (br, 1H).

5

Example 445

2-(2-{3-[2-(Isobutyl-methyl-amino)-4-methyl-phenyl]-ureido}-thiazol-4-yl)-N-methyl-acetamide




2-(2-{3-[2-(Isobutyl-methyl-amino)-4-methyl-phenyl]-ureido}-thiazol-4-yl)-N-methyl-acetamide (63 mg, 65%) was prepared from (2-{3-[2-(isobutyl-methyl-amino)-4-methyl-phenyl]-ureido}-thiazol-4-yl)-acetic acid (95 mg, 0.25 mmol) and methylamine following the general procedure K.

LC-MS (m/z): 390 ($M+1$)⁺, ¹H NMR (400 MHz, CDCl₃): δ 0.85 (d, 6H), 1.67 (m, 1H), 2.29 (s, 3H), 2.63 (s, 3H), 2.75 (s, 2H), 2.80 (s, 3H), 3.62 (s, 2H), 6.50 (br, 1H), 6.62 (s, 1H), 6.93 (m, 2H), 8.04 (d, 1H), 8.68 (br, 1H), 10.00 (br, 1H).

15

Example 446

2-(2-{3-[2-(Isobutyl-methyl-amino)-4-methyl-phenyl]-ureido}-thiazol-4-yl)-N-(2-methoxy-ethyl)-acetamide




2-(2-{3-[2-(isobutyl-methyl-amino)-4-methyl-phenyl]-ureido}-thiazol-4-yl)-N-(2-methoxy-ethyl)-acetamide (78 mg, 72%) was prepared from (2-{3-[2-(isobutyl-methyl-amino)-4-methyl-phenyl]-ureido}-thiazol-4-yl)-acetic acid (95 mg, 0.25 mmol) and 2-methoxymethylamine following the general procedure K.

LC-MS (m/z): 434 ($M+1$)⁺.

Example 447

1-(4-Methyl-2-pyrrolidin-1-yl-phenyl)-3-thiazol-2-yl-urea



10


4-Methyl-2-(pyrrolidin-1-yl)aniline (0.57 g, 65%) was prepared from pyrrolidine (0.71 g, 10.0 mmol) and 3-fluoro-4-nitrotoluene (0.77 g, 5.0 mmol) following the general procedure W. 1-(4-Methyl-2-pyrrolidin-1-yl-phenyl)-3-thiazol-2-yl-urea (217 mg, 72%) was prepared from 4-methyl-2-(pyrrolidin-1-yl)aniline (176 mg, 1.0 mmol) and 2-aminothiazole (100 mg, 1.0 mmol) following the general procedure D.

15

LC-MS (m/z): 303 ($M+1$)⁺, ¹H NMR (400 MHz, DMSO- d_6): δ 1.89 (m, 4H), 2.23 (s, 3H), 3.02 (m, 4H), 6.75 (dd, 1H), 6.88 (d, 1H), 7.09 (d, 1H), 7.36 (dd, 1H), 7.70 (d, 1H), 8.40 (br, 1H), 11.07 (br, 1H).

Example 448

1-[2-(2,3-Dihydro-indol-1-yl)-5-fluoro-phenyl]-3-thiazol-2-yl-urea




2-(2,3-Dihydro-indol-1-yl)-5-fluoro-aniline (0.59 g, 52%) was prepared from
5 indoline (0.6 g, 5.0 mmol) and 2,5-difluoro-nitrobenzene (0.80 g, 5.0 mmol) following the
general procedure W. 1-[2-(2,3-Dihydro-indol-1-yl)-5-fluoro-phenyl]-3-thiazol-2-yl-urea
(198 mg, 56%) was prepared from 2-(2,3-dihydro-indol-1-yl)-5-fluoro-aniline (228 mg, 1.0
mmol) and 2-aminothiazole (100 mg, 1.0 mmol) following the general procedure D.

LC-MS (m/z): 355 ($M+1$)⁺, ¹H NMR (400 MHz, DMSO- d_6): δ 3.16 (t, 2H), 3.70 (br, 2H),
10 6.10 (d, 1H), 6.73 (m, 1H), 6.94 (m, 2H), 7.14 (d, 1H), 7.22 (d, 1H), 7.34 (m, 2H), 8.13 (dd,
1H), 9.01 (br, 1H), 11.32 (br, 1H).

Example 449

1-(4-Methyl-2-piperidin-1-yl-phenyl)-3-thiazol-2-yl-urea




4-Methyl-2-(piperidin-1-yl)-aniline (0.63 g, 67%) was prepared from piperidine
15 (0.85 g, 10.0 mmol) and 3-fluoro-4-nitrotoluene (0.77 g, 5.0 mmol) following the general
procedure W. 1-(4-Methyl-2-piperidin-1-yl-phenyl)-3-thiazol-2-yl-urea (214 mg, 68%) was
prepared from 4-methyl-2-(piperidin-1-yl)-aniline (190 mg, 1.0 mmol) and 2-aminothiazole
20 (100 mg, 1.0 mmol) following the general procedure D.

LC-MS (m/z): 317 ($M+1$)⁺, ¹H NMR (400 MHz, CDCl₃): δ 1.49 (m, 2H), 1.69 (m, 4H), 2.30 (s, 3H), 2.82 (t, 4H), 6.92 (m, 3H), 7.47 (d, 1H), 7.96 (d, 1H), 8.90 (br, 1H), 11.36 (br, 1H).

5 EXAMPLE 450

{2-[3-(4-Methyl-2-piperidin-1-yl-phenyl)-ureido]-thiazol-4-yl}-acetic acid




{2-[3-(4-Methyl-2-piperidin-1-yl-phenyl)-ureido]-thiazol-4-yl}-acetic acid ethyl ester (285 mg, 71%) was prepared from 4-methyl-2-(piperidin-1-yl)-aniline (190 mg, 1.0 mmol) and ethyl 2-amino-4-thiazolylacetate (186 mg, 1.0 mmol) following the general procedure D.
Hydrolysis of this ester following general procedure J gave {2-[3-(4-methyl-2-piperidin-1-yl-phenyl)-ureido]-thiazol-4-yl}-acetic acid (238 mg, 90%).

LC-MS (m/z): 375 ($M+1$)⁺; ¹H NMR (400 MHz, DMSO-*d*₆): δ 1.53 (br, 2H), 1.75 (m, 4H), 2.23 (s, 3H), 2.70 (m, 4H), 3.55 (s, 2H), 6.83 (s, 1H), 6.87 (d, 1H), 6.99 (s, 1H), 7.92 (d, 1H), 8.40 (br, 1H), 11.28 (br, 1H), 12.42 (br, 1H).

Example 451

1-[2-(2-Fluoro-6-methoxy-phenoxy)-4-methyl-phenyl]-3-thiazol-2-yl-urea




3-(2-fluoro-6-methoxyphenoxy)-nitrotoluene (0.9 g, 65%) was prepared from 2-fluoro-6-methoxyphenol (0.78 g, 5.5 mmol) and 3-fluoro-4-nitrotoluene (0.77 g, 5.0 mmol) following the general procedure A. This compound was reduced to 4-fluoro-2-(2-fluoro-6-

methoxyphenoxy)aniline (0.58 g, 72%) following the general procedure C. 1-[2-(2-Fluoro-6-methoxy-phenoxy)-4-methyl-phenyl]-3-thiazol-2-yl-urea (122 mg, 65%) was prepared from 2-(2-fluoro-6-methoxy-phenoxy)-4-methyl-aniline (124 mg, 0.5 mmol) and 2-aminothiazole (60 mg, 0.6 mmol) following the general procedure D.

- 5 LCMS (m/z): 374 ($M + 2$)*; ^1H NMR (400 MHz, DMSO- d_6): δ 2.13 (s, 3H), 3.79 (s, 3H), 6.30 (s, 1H), 6.83 (d, 1H), 7.07 (m, 3H), 7.34 (m, 2H), 8.06 (d, 1H), 8.97 (br, 1H), 10.93 (s, 1H).

EXAMPLE 452

- 10 1-[2-(2-Fluoro-6-methoxy-phenoxy)-4-(4-methyl-piperazin-1-ylmethyl)-phenyl]-3-thiazol-2-yl-urea




- 15 3-(2-fluoro-6-methoxyphenoxy)-nitrotoluene (0.28 g, 1.0 mmol) was heated with N-bromosuccinimide (200 mg, 1.1 mmol) and dibenzoyl peroxide (10 mg) in CCl_4 (10 mL) at 90 °C for 3 hr. The reaction mixture was cooled, filtered and the filtrate was concentrated. To the residue was added N-methylpiperazine (110 mg, 1.1 mmol) in THF (5 mL) and the mixture was heated at 60 °C for 4 hr. The volatiles were removed under reduced pressure. The residue was dissolved in ethyl acetate and washed (brine), dried (Na_2SO_4) and concentrated.
- 20 The resulting nitrobenzene was reduced to 2-(2-fluoro-6-methoxy-phenoxy)-4-(4-methyl-piperazin-1-ylmethyl)-aniline (224 mg, 65%) following the general procedure C. 1-[2-(2-Fluoro-6-methoxy-phenoxy)-4-(4-methyl-piperazin-1-ylmethyl)-phenyl]-3-thiazol-2-yl-urea (130 mg, 55%) was prepared from 2-(2-fluoro-6-methoxy-phenoxy)-4-(4-methyl-piperazin-1-ylmethyl)-aniline (172 mg, 0.5 mmol) and 2-aminothiazole (60 mg, 0.6 mmol) following the
- 25 general procedure D.

LCMS (m/z): 472 ($M + 2$)*; ^1H NMR (400 MHz, DMSO- d_6): δ 2.12 (s, 3H), 2.24 (br, 8H), 3.28 (s, 2H), 3.77 (s, 3H), 6.46 (s, 1H), 6.89 (d, 1H), 7.07 (m, 3H), 7.34 (m, 2H), 8.11 (d, 1H), 9.02 (br, 1H), 10.96 (s, 1H).

Example 453

(2-{3-[2-(2-Fluoro-6-methoxy-phenoxy)-4-methyl-phenyl]-ureido}-thiazol-4-yl)-acetic acid




(2-{3-[2-(2-Fluoro-6-methoxy-phenoxy)-4-methyl-phenyl]-ureido}-thiazol-4-yl)-acetic acid ethyl ester (330 mg, 72%) was prepared from 2-(2-fluoro-6-methoxyphenoxy)-aniline (247 mg, 1.0 mmol) and ethyl 2-amino-4-thiazolylacetate (186 mg, 1.0 mmol) following the general procedure D. Hydrolysis of this ester following general procedure J gave (2-{3-[2-(2-fluoro-6-methoxy-phenoxy)-4-methyl-phenyl]-ureido}-thiazol-4-yl)-acetic acid (283 mg, 91%).

LCMS (m/z): 432 ($M + 2$)⁺; ¹H NMR (400 MHz, DMSO- d_6): δ 2.13 (s, 3H), 3.55 (s, 2H), 3.79 (s, 3H), 6.30 (s, 1H), 6.83 (d, 1H), 6.88 (s, 1H), 7.03 (m, 2H), 7.34 (m, 1H), 8.04 (d, 1H), 9.05 (br, 1H), 11.02 (br, 2H).

Example 454

1-[2-(2-Fluoro-6-methoxy-phenoxy)-5-formyl-phenyl]-3-thiazol-2-yl-urea



15

4-(2-Fluoro-6-methoxy-phenoxy)-3-nitrobenzaldehyde (1.0 g, 70%) was prepared from 2-fluoro-6-methoxyphenol (0.78 g, 5.5 mmol) and 4-fluoro-3-nitrobenzaldehyde (0.84 g, 5.0 mmol) following the general procedure A. This was reduced to 5-formyl-2-(2-fluoro-6-methoxy-phenoxy)-aniline (0.57 g, 62%) following general procedure B. 1-[2-(2-Fluoro-6-methoxy-phenoxy)-5-formyl-phenyl]-3-thiazol-2-yl-urea (265 mg, 69%) was prepared from 5-


20

formyl-2-(2-fluoro-6-methoxy-phenoxy)-aniline (261 mg, 1.0 mmol) and 2-aminothiazole (100 mg, 1.0 mmol) following the general procedure D.

LC-MS (*m/z*): 388 (*M*+1)⁺, ¹H NMR (400 MHz, DMSO-*d*₆): δ 3.80 (s, 3H), 6.71 (d, 1H), 7.14 (m, 3H), 7.39 (m, 2H), 7.52 (dd, 1H), 8.78 (d, 1H), 9.31 (br, 1H), 9.90 (s, 1H), 11.04 (s, 1H).

Example 455

1-[2-(2-Fluoro-6-methoxy-phenoxy)-5-morpholin-4-ylmethyl-phenyl]-3-thiazol-2-yl-urea




1-[2-(2-Fluoro-6-methoxy-phenoxy)-5-morpholin-4-ylmethyl-phenyl]-3-thiazol-2-yl-urea (35 mg, 78%) was prepared from 1-[2-(2-fluoro-6-methoxy-phenoxy)-5-formyl-phenyl]-3-thiazol-2-yl-urea (39 mg, 0.1 mmol) and morpholine following the general procedure O.

LC-MS (*m/z*): 459 (*M*+1)⁺, ¹H NMR (400 MHz, DMSO-*d*₆): δ 2.24 (s, 4H), 3.30 (s, 2H), 3.48 (s, 4H), 3.77 (s, 3H), 6.48 (s, 1H), 6.93 (d, 1H), 7.10 (m, 3H), 7.37 (m, 2H), 8.12 (d, 1H), 9.04 (br, 1H), 10.96 (br, 1H).

Example 456

1-[2-(2-Fluoro-6-methoxy-phenoxy)-5-methanesulfonylmethyl-phenyl]-3-thiazol-2-yl-urea




4-(2-Fluoro-6-methoxyphenoxy)-3-nitrotoluene (0.28 g, 1.0 mmol) was heated with N-bromosuccinimide (200 mg, 1.1 mmol) and dibenzoyl peroxide (10 mg) in CCl₄ (10 mL) at 90 °C for 3 hr. The reaction mixture was cooled, filtered and the filtrate was concentrated. To this residue was added sodium methanesulfinate (110 mg, 1.1 mmol) in THF (5 mL) and the mixture was heated at 60 °C for 4 hr. The volatiles were removed under reduced pressure. The residue was dissolved in ethyl acetate and was washed (brine), dried (Na₂SO₄) and concentrated. The resulting nitrobenzene was reduced to 2-(2-fluoro-6-methoxy-phenoxy)-5-methanesulfonylmethyl-aniline (210 mg, 65%) following the general procedure B. 1-[2-(2-Fluoro-6-methoxy-phenoxy)-5-methanesulfonylmethyl-phenyl]-3-thiazol-2-yl-urea (138 mg, 61%) was prepared from 2-(2-fluoro-6-methoxy-phenoxy)-5-methanesulfonylmethyl-aniline (162 mg, 0.5 mmol) and 2-aminothiazole (60 mg, 0.6 mmol) following the general procedure D.

LCMS (*m/z*): 452 (M+1)⁺; ¹H NMR (400 MHz, DMSO-*d*₆): δ 2.93 (s, 3H), 3.35 (s, 2H), 3.80 (s, 3H), 6.52 (d, 1H), 6.93 (d, 1H), 7.07 (m, 3H), 7.35 (m, 2H), 8.31 (s, 1H), 9.22 (br, 1H), 11.02 (s, 1H).

Example 457

{5-Chloro-2-[3-(2-cyclopentanecarbonyl-4-methyl-phenyl)-ureido]-thiazol-4-yl}-acetic acid ethyl ester




- 5 To a solution of {2-[3-(2-cyclopentanecarbonyl-4-methyl-phenyl)-ureido]-thiazol-4-yl}-acetic acid ethyl ester (0.05 g, 0.12 mmol) in 2 ml of acetonitrile was added N-chlorosuccinimide (0.018 g, 0.13 mmol). The reaction mixture was stirred under nitrogen in a pressure bottle excluded from light for 1.5 hour at 80°C. The mixture was diluted with 15 ml of DCM, washed (1N HCl, saturated NaHCO₃, water), dried (MgSO₄) and concentrated under
10 reduced pressure. The crude product was purified by preparative HPLC to afford the title compound (0.08 g, 15%).

HPLC-MS (Method A): m/z = 450, 452 ($M+1$)⁺; R_t = 5.55 min.

- ¹H NMR (300 MHz, CDCl₃): δ 1.29 (t, J = 7.2 Hz, 3H), 1.71 (m, 4H), 1.90 (m, 4H), 2.37 (s, 3H), 3.70 (s, 2H), 3.75 (m, 1H), 4.24 (q, J = 7.2 Hz, 2H), 7.35 (d, J = 7.9 Hz, 1H), 7.70 (s,
15 1H), 8.42 (d, J = 8.3 Hz, 1H) and 11.59 (s, 1H)

Example 458

{5-Bromo-2-[3-(2-cyclopentanecarbonyl-4-methyl-phenyl)-ureido]-thiazol-4-yl}-acetic acid ethyl ester



- 20 To a solution of {2-[3-(2-cyclopentanecarbonyl-4-methyl-phenyl)-ureido]-thiazol-4-yl}-acetic acid ethyl ester (0.05 g, 0.12 mmol) in 2 ml of acetonitrile was added N-bromosuccinimide (0.024 g, 0.13 mmol). The reaction mixture was stirred under nitrogen in a


pressure bottle excluded from light overnight at 25°C. The mixture was diluted with 15 ml of DCM, washed (1N HCl, saturated NaHCO₃, water), dried (MgSO₄) and concentrated under reduced pressure. The crude product was purified by preparative HPLC to afford the title compound (0.013 g, 21%).

5 HPLC-MS (Method A): m/z = 494, 496 (M+1)⁺; R_t = 5.45 min.

¹H NMR (300 MHz, CDCl₃): δ 1.29 (t, J = 7.2 Hz, 3H), 1.71 (m, 4H), 1.92 (m, 4H), 2.38 (s, 3H), 3.73 (s, 2H), 3.77 (m, 1H), 4.23 (q, J = 7.2 Hz, 2H), 7.36 (dd, J = 1.3, 8.5 Hz, 1H), 7.73 (s, 1H), 8.37 (d, J = 8.3 Hz, 1H) and 11.81 (s, 1H).

Example 459

10 {5-Chloro-2-[3-(2-cyclopentanecarbonyl-4-methyl-phenyl)-ureido]-thiazol-4-yl}-acetic acid




To a suspension of {2-[3-(2-cyclopentanecarbonyl-4-methyl-phenyl)-ureido]-thiazol-4-yl}-acetic acid (0.50 g, 1.29 mmol) in 20 ml of acetonitrile was added N-chlorosuccinimide (0.19 g, 1.42 mmol). The reaction mixture was stirred under nitrogen, in a pressure bottle excluded from light, for 2h at 80°C. The mixture was concentrated under reduced pressure, dissolved in DMSO and purified by preparative HPLC to afford the title compound (0.128 g, 24%).

15 HPLC-MS (Method A): m/z = 422, 424 (M+1)⁺; R_t = 5.84 min.

¹H NMR (300 MHz, DMSO-*d*₆): δ 1.64 (m, 4H), 1.77 (m, 2H), 1.90 (m, 2H), 1.90 (m, 2H), 2.34 (s, 3H), 3.56 (s, 2H), 3.88 (m, 1H), 7.40 (dd, J = 1.5, 8.7 Hz, 1H), 7.86 (d, J = 1.1 Hz, 1H), 8.15 (d, J = 8.3 Hz, 1H) and 10.69 (s, 1H)

Example 460

{5-Bromo-2-[3-(2-cyclopentanecarbonyl-4-methyl-phenyl)-ureido]-thiazol-4-yl}-acetic acid




To a suspension of {2-[3-(2-cyclopentanecarbonyl-4-methyl-phenyl)-ureido]-thiazol-4-yl}-acetic acid ethyl ester (0.20 g, 0.52 mmol) in 10 ml of acetic acid was added N-bromosuccinimide (0.10 g, 0.57 mmol). The reaction mixture was stirred under nitrogen in a pressure bottle excluded from light overnight at 25°C. The mixture was concentrated under reduced pressure, dissolved in DMSO and purified by preparative HPLC to afford the title compound (0.137 g, 57%).

HPLC-MS (Method A): m/z = 466, 468 ($M+1$)⁺; R_t = 4.59 min.

¹H NMR (300 MHz, DMSO-*d*₆): δ 1.64 (m, 4H), 1.76 (m, 2H), 1.89 (m, 2H), 2.34 (s, 3H), 3.55 (s, 2H), 3.88 (m, 1H), 7.40 (d, J = 8.3 Hz, 1H), 7.87 (s, 1H), 8.15 (d, J = 8.7 Hz, 1H), 10.71 (s, 1H) and 12.15 (s, 1H)

Example 461

Bromo-2-[3-(2-cyclopentanecarbonyl-4-methyl-phenyl)-ureido]-thiazol-4-yl}-acetic acid ethyl ester



To a solution of {2-[3-(2-cyclopentanecarbonyl-4-methyl-phenyl)-ureido]-thiazol-4-yl}-acetic acid ethyl ester (0.05 g, 0.12 mmol) in 2 ml of acetonitrile was added N-bromosuccinimide (0.024 g, 0.13 mmol). The reaction mixture was stirred under nitrogen in a pressure bottle excluded from light overnight at 25°C. The mixture was diluted with 15 ml of


DCM, washed (1N HCl, saturated NaHCO₃, water), dried (MgSO₄) and concentrated under reduced pressure. The crude product was purified by preparative HPLC to afford the title compound (0.010 g, 17%).

HPLC-MS (Method A): m/z = 494, 496 (M+1)⁺; R_t = 5.19 min.

- 5 ¹H NMR (300 MHz, CDCl₃): δ 1.33 (t, J = 7.2 Hz, 3H), 1.71 (m, 4H), 1.92 (m, 4H), 2.37 (s, 3H), 3.76 (m, 1H), 4.30 (dq, J = 2.3, 7.2 Hz, 2H), 5.45 (s, 1H), 7.17 (s, 1H), 7.36 (d, J = 8.7 Hz, 1H), 7.73 (s, 1H), 8.40 (d, J = 8.7 Hz, 1H) and 11.83 (s, 1H)

Example 462 (General procedure D)

- 10 {2-[3-(2-Cyclopentanecarbonyl-phenyl)-ureido]-thiazol-4-yl}-acetic acid ethyl ester




- To a stirred solution of phenyl-carbamic acid tert-butyl ester (1.2 g, 6.2 mmol) in Et₂O (12 mL) under a nitrogen atmosphere was added dropwise a 1.7 M solution of t-BuLi in pentan (8.4 mL, 14.3 mmol) over a 10-min period at -20°C. The mixture was stirred at -10°C for 2.5 h and then cyclopentanecarboxylic acid methoxy-methyl-amide (1.07 g, 6.8 mmol) was added over 5 min. The mixture was allowed to warm to room temperature and stirred for further 1 h before it was quenched with aqueous NH₄Cl. The organic phase was isolated and the aqueous phase was extracted with CH₂Cl₂, and the combined organic phases were dried and concentrated *in vacuo*. The crude material was dissolved CH₂Cl₂ (18 mL) and TFA (6 mL) and stirred at room temperature for 2 h. The reaction mixture was evaporated *in vacuo*, and added aqueous NaHCO₃ to pH 7 and extracted with CH₂Cl₂. The combined organic phases were dried and concentrated *in vacuo* to give 1.03 g (68%) of (2-amino-phenyl)-cyclopentyl-methanone as an oil.

- The title compound (41 mg, 16 %) was prepared from 2-amino-phenyl-cyclopentyl-methanone (120 mg, 0.63 mmol) and ethyl-2-amino-4-thiazolyl acetate (118 mg, 0.63 mmol) following the general procedure D.

¹H NMR (400MHz; CDCl₃): δ 1.28 (t, 3H), 1.66-1.77 (m, 4H), 1.84-1.96 (m, 4H), 3.70 (s, 2H), 3.74 (br t, 1H), 4.20 (q, 2H), 6.72 (s, 1H), 7.10 (t, 1H), 7.53 (t, 1H), 7.92 (d, 1H), 8.57 (br s, 1H), 11.78 (s, 1H); HPLC-MS : m/z = 424.0 (M+23); R_t = 4.45 min.

Example 463 (General procedure J)

{2-[3-(2-Cyclopentanecarbonyl-phenyl)-ureido]-thiazol-4-yl}-acetic acid



5

The title compound (9 mg, 27 %) was prepared from {2-[3-(2-cyclopentanecarbonyl-phenyl)-ureido]-thiazol-4-yl}-acetic acid ethyl ester (208 mg, 0.5 mmol) following the general procedure J.


¹H NMR (400MHz; Acetone d₆): δ 1.85-2.11 (m, 8H), 3.71 (s, 2H), 3.94 (br t, 1H), 6.86 (s, 1H), 7.18 (t, 1H), 7.79 (t, 1H), 8.09 (d, 1H), 8.56 (d, 1H), 11.36 (s, 1H); HPLC-MS : m/z = 396.1 (M+23); R_t = 3.87 min.

10

Example 464 (General procedure AA)

{2-[3-(2-Cyclopentanecarbonyl-4-methoxy-phenyl)-ureido]-thiazol-4-yl}-acetic acid

15 ethyl ester



To a stirred solution of (4-methoxy-phenyl)-carbamic acid tert-butyl ester (2.0 g, 9.0 mmol) in Et₂O (20 mL) under a nitrogen atmosphere was added dropwise a 1.7 M solution of t-BuLi in pentan (12.1 mL, 20.6 mmol) over a 10-min period at -20 °C. The mixture was stirred at -15 °C for 1.5 h and then cyclopentanecarboxylic acid methoxy-methyl-amide (1.55 g, 9.9 mmol) was added over 5 min. The mixture was allowed to warm to room temperature and stirred for further 1 h before it was quenched with aqueous NH₄Cl. The organic phase was isolated and the aqueous phase was extracted with CH₂Cl₂, and the combined organic phases were dried and concentrated *in vacuo*. The crude material was

20


dissolved CH_2Cl_2 (18 mL) and TFA (6 mL) and stirred at room temperature for 2 h. The reaction mixture was evaporated *in vacuo*, and added aqueous NaHCO_3 to pH 7 and extracted with CH_2Cl_2 , dried and concentrated to give a crude product which was purified by flash chromatography (Quad flash 25, EtOAc-heptane 1:20 \rightarrow 1:4). This gave 0.65 g (31%) of (2-amino-5-methoxy-phenyl)-cyclopentyl-methanone as an oil.

The title compound (158 mg, 80 %) was prepared from (2-amino-5-methoxy-phenyl)-cyclopentyl-methanone (100 mg, 0.46 mmol) and ethyl-2-amino-4-thiazolyl acetate (85 mg, 0.46 mmol) following the general procedure AA.

^1H NMR (400MHz; CDCl_3): δ 1.28 (t, 3H), 1.66-1.74 (m, 4H), 1.86-1.96 (m, 4H), 3.68 (br t, 1H), 3.70 (s, 2H), 3.84 (s, 3H), 4.20 (q, 2H), 6.71 (s, 1H), 7.10 (dd, 1H), 7.42 (d, 1H), 8.45 (br s, 1H), 9.13 (br s, 1H), 11.31 (br s, 1H); HPLC-MS : m/z = 454.1 ($M+23$); R_t = 5.48 min.

Example 465 (General procedure J)

{2-[3-(2-Cyclopentanecarbonyl-4-methoxy-phenyl)-ureido]-thiazol-4-yl}-acetic acid




The title compound (92 mg, 86 %) was prepared from {2-[3-(2-cyclopentanecarbonyl-4-methoxy-phenyl)-ureido]-thiazol-4-yl}-acetic acid ethyl ester (112 mg, 0.26 mmol) following the general procedure J.

^1H NMR (400MHz; $\text{DMSO}-d_6$): δ 1.58-1.65 (m, 4H), 1.70-1.92 (m, 4H), 3.57 (s, 2H), 3.82 (s, 3H), 3.88 (br p, 1H), 6.84 (s, 1H), 7.19 (dd, 1H), 7.44 (s, 1H), 8.08 (d, 1H), 10.22 (s, 1H), 12.05 (br s, 2H); HPLC-MS : m/z = 426.1 ($M+23$); R_t = 4.49 min.

Example 466 (General procedure AA)

{2-[3-(2-Cyclopropanecarbonyl-4-methyl-phenyl)-ureido]-thiazol-4-yl}-acetic acid ethyl ester




To a stirred solution of p-tolyl-carbamic acid tert-butyl ester (1.0 g, 4.8 mmol) in Et₂O (10 mL) under a nitrogen atmosphere was added dropwise a 1.7 M solution of t-BuLi in pentan (6.5 mL, 11.1 mmol) over a 10-min period at -20°C. The mixture was stirred at -10°C for 2.5 h and then cyclopropanecarboxylic acid methoxy-methyl-amide (0.92 g, 6.3 mmol) was added over 5 min. The mixture was allowed to warm to room temperature and stirred for further 1 h before it was quenched with aqueous NH₄Cl. The organic phase was isolated and the aqueous phase was extracted with CH₂Cl₂, and the combined organic phases were dried and concentrated *in vacuo*. The crude material was dissolved CH₂Cl₂ (15 mL) and TFA (15 mL) and stirred at room temperature for 3 h. The reaction mixture was evaporated *in vacuo*, and added aqueous NaHCO₃ to pH 7 and extracted with CH₂Cl₂. The combined organic phases were dried and concentrated *in vacuo* to give 0.80 g crude (2-amino-5-methyl-phenyl)-cyclopropyl-methanone as a yellow oil.

The title compound (190 mg, 43 %) was prepared from crude (2-amino-5-methyl-phenyl)-cyclopropyl-methanone (200 mg, 1.14 mmol) and ethyl-2-amino-4-thiazolyl acetate (212 mg, 1.14 mmol) following the general procedure AA.

¹H NMR (400MHz; CDCl₃): δ 1.04-1.10 (m, 2H), 1.24-1.30 (m, 5H), 2.39 (s, 3H), 2.62-2.72 (m, 1H), 3.73 (s, 2H), 4.20 (q, 2H), 6.78 (s, 1H), 7.37 (dd, 1H), 7.89 (d, 1H), 8.37 (d, 1H), 9.42 (br s, 2H), 11.50 (s, 1H); HPLC-MS : *m/z* = 410.0 (M+23); R_t = 4.11 min.

Example 467 (General procedure J)

{2-[3-(2-Cyclopropanecarbonyl-4-methyl-phenyl)-ureido]-thiazol-4-yl}-acetic acid




The title compound (111 mg, 99 %) was prepared from {2-[3-(2-cyclopentanecarbonyl-phenyl)-ureido]-thiazol-4-yl}-acetic acid ethyl ester (120 mg, 0.31 mmol) following the general procedure J.

^1H NMR (400MHz; DMSO d_6): δ 1.04 (br s, 4H), 2.38 (s, 3H), 2.76 (br s, 1H), 3.58 (s, 2H), 6.86 (s, 1H), 6.92 (d, 1H), 8.02 (s, 1H), 8.14 (d, 1H), 10.46 (s, 1H), 12.05 (br s, 2H); HPLC-MS : m/z = 382.0 ($M+23$); R_t = 3.38 min.

5 **Example 468** (General procedure AA)

{2-[3-(2-Cyclobutanecarbonyl-4-methyl-phenyl)-ureido]-thiazol-4-yl}-acetic acid ethyl ester




Crude (2-amino-5-methyl-phenyl)-cyclobutyl-methanone (80%, oil) was prepared in a similar fashion as described for (2-amino-5-methyl-phenyl)-cyclopropyl-methanone using cyclobutanecarboxylic acid methoxy-methyl-amide instead of cyclopropanecarboxylic acid methoxy-methyl-amide.

The title compound (210 mg, 49 %) was prepared from crude (2-amino-5-methyl-phenyl)-cyclobutyl-methanone (200 mg, 1.06 mmol) and ethyl-2-amino-4-thiazolyl acetate (197 mg, 1.06 mmol) following the general procedure AA.

^1H NMR (400MHz; CDCl_3): δ 1.30 (t, 3H), 1.83-2.50 (m, 6H), 2.36 (s, 3H), 3.76 (s, 2H), 4.05 (p, 1H), 4.22 (q, 2H), 6.80 (s, 1H), 7.35 (d, 1H), 7.52 (s, 1H), 8.41 (br d, 1H), 11.9 (br s, 1H); HPLC-MS : m/z = 424.1 ($M+23$); R_t = 4.48 min.

20 **Example 469** (General procedure J)

{2-[3-(2-Cyclobutanecarbonyl-4-methyl-phenyl)-ureido]-thiazol-4-yl}-acetic acid




The title compound (112 mg, 73 %) was prepared from {2-[3-(2-cyclobutanecarbonyl-4-methyl-phenyl)-ureido]-thiazol-4-yl}-acetic acid ethyl ester (165 mg, 0.41 mmol) following the general procedure J.

^1H NMR (400MHz; DMSO d_6): δ 1.70-2.30 (m, 6H), 2.32 (s, 3H), 3.56 (s, 2H), 4.19 (br t, 1H), 6.86 (s, 1H), 7.39 (d, 1H), 7.63 (s, 1H), 8.22 (d, 1H), 10.81 (s, 1H), 12.15 (br s, 2H); HPLC-MS : m/z = 396.1 (M+23); R_t = 3.75 min.

5

Example 470 (General procedure AA)

{2-[3-(2-Cyclohexanecarbonyl-4-methyl-phenyl)-ureido]-thiazol-4-yl}-acetic acid ethyl ester



Crude (2-amino-5-methyl-phenyl)-cyclohexyl-methanone (75%, oil) was prepared in a similar fashion as described for (2-amino-5-methyl-phenyl)-cyclopropyl-methanone using cyclohexanecarboxylic acid methoxy-methyl-amide instead of cyclopropanecarboxylic acid methoxy-methyl-amide.


The title compound (188 mg, 48 %) was prepared from crude (2-amino-5-methyl-phenyl)-cyclohexyl-methanone (200 mg, 0.92 mmol) and ethyl-2-amino-4-thiazolyl acetate (171 mg, 1.06 mmol) following the general procedure AA.

^1H NMR (400MHz; CDCl_3): δ 1.30 (t, 3H), 1.37-1.59 (m, 5H), 1.71-1.91 (m, 5H), 2.37 (s, 3H), 3.30 (br s, 1H), 3.73 (s, 2H), 4.22 (q, 2H), 6.78 (s, 1H), 7.34 (d, 1H), 7.68 (s, 1H), 8.39 (br d, 1H), 11.78 (br s, 1H); HPLC-MS : m/z = 452.2 (M+23); R_t = 4.92 min.

20

Example 471 (General procedure J)

{2-[3-(2-Cyclohexanecarbonyl-4-methyl-phenyl)-ureido]-thiazol-4-yl}-acetic acid




The title compound (100 mg, 89 %) was prepared from {2-[3-(2-cyclohexanecarbonyl-4-methyl-phenyl)-ureido]-thiazol-4-yl}-acetic acid ethyl ester (112 mg, 0.28 mmol) following the general procedure J.

¹H NMR (400MHz; DMSO d₆): δ 1.13-1.45 (m, 5H), 1.62-1.83 (m, 5H), 2.34 (s, 3H), 3.42 (br s, 1H), 3.57 (s, 2H), 6.83 (s, 1H), 7.39 (d, 1H), 7.81 (s, 1H), 8.15 (d, 1H), 10.50 (s, 1H), 12.07 (br s, 2H); HPLC-MS : *m/z* = 402.0 (M+1); R_t = 4.18 min.

Example 472

{2-[(2-Cyclopentanecarbonyl-4-methyl-phenylamino)oxalyl]-amino]-thiazol-4-yl}-acetic acid ethyl ester



To a solution of ethyl-2-aminothiazole-4-acetate (2 g) in dichloromethane (80 mL) at ice bath temperature was added ethyl oxalyl chloride (1.2 mL) and the reaction stirred 1 h. The reaction mixture was washed with water, and the organic phase concentrated to give N-(4-ethoxycarbonylmethyl-thiazol-2-yl)-oxalamic acid ethyl ester (1.2 g).


To N-(4-ethoxycarbonylmethyl-thiazol-2-yl)-oxalamic acid ethyl ester (1 g) was added THF (5 mL) and 2 N lithium hydroxide (4 mL). A white precipitate formed and after 1h the reaction mixture was acidified to pH5 and the white precipitate recovered by filtration, and triturated with a 1:1 mixture of ether and ethyl acetate to remove unreacted starting material. The solid residue was dried in a vacuum oven to give N-(4-ethoxycarbonylmethyl-thiazol-2-yl)-oxalamic acid (300 mg).

N-(4-ethoxycarbonylmethyl-thiazol-2-yl)-oxalamic acid (80 mg) was dissolved in DMF (5 mL) and (2-amino-5-methyl-phenyl)-cyclopentyl-methanone was added followed by PyBOP. The reaction was stirred for 24 h and water was added. The aqueous phase was extracted with ether/ethyl acetate and the organic phase dried, concentrated and purified by flash chromatography to give the title compound (15 mg).

¹H NMR (CDCl₃): 1.25 (3H, t), 1.6-2.1 (8H, m), 2.45 (3H, s), 3.65-3.75 (1H, m), 3.7 (1H, s), 4.2 (4H, q), 6.9 (1H), 7.4 (1H, dd), 7.8 (1H, s), 8.65 (1H, d), 10.5 (1H, bs), 13.3 (1H, s).

Example 473

{2-[(2-Cyclopentanecarbonyl-4-methyl-phenylamino)oxalyl]-amino}-thiazol-4-yl]-acetic acid




5 To {2-[(2-cyclopentanecarbonyl-4-methyl-phenylamino)oxalyl]-thiazol-4-yl]-acetic acid ethyl ester (15 mg) in methanol (2 mL) was added 1N sodium hydroxide (1N) and the mixture was stirred 3 h at room temperature. The mixture was acidified with 1N HCl, concentrated under reduced pressure and purified by HPLC to give the title compound (4 mg).

10 LCMS (m/z): 416 (M+1)

¹H NMR (CDCl₃ + MeOD): 1.6-2.1 (8H, m), 2.45 (3H, s), 3.7 (1H, s), 3.70-3.85 (1H, m), 6.9 (1H), 7.4 (1H, dd), 7.8 (1H, s), 8.65 (1H, d), 13.3 (1H, s).

Example 474

15 [2-[3-(2-Cyclopentanecarbonyl-4-methyl-phenyl)-ureido]-5-(pyridin-2-ylsulfanyl)-thiazol-4-yl]-acetic acid ethyl ester




To a solution of {5-chloro-2-[3-(2-cyclopentanecarbonyl-4-methyl-phenyl)-ureido]-thiazol-4-yl]-acetic acid ethyl ester (0.1 g, 0.22 mmol) in DMF (4 mL) was added sodium hydrogen carbonate (0.075 g, 0.89 mmol) and 2-pyridylthiole (0.027 g, 0.24 mmol). The reaction mixture was stirred under nitrogen in a pressure bottle excluded from light for 2.0 hour at 80°C. The crude product was purified by flash chromatography

(isopropanole/heptane 1:4) and recrystallized (dichloromethane/heptane) to afford the title compound (0.02 g).

¹H NMR (300 MHz, CDCl₃): δ 1.20 (t, 3H), 1.60-1.82 (m, 4H), 1.82-2.02 (m, 4H), 2.38 (s, 3H), 3.68-3.80 (m, 1H), 3.85 (s, 2H), 4.12 (q, 2H), 7.00 (t, 2H), 7.35 (d, 1H), 7.50 (t, 1H), 7.73 (s, 1H), 8.40 (m, 2H), 11.80 (s, 1H).

Example 475 (General procedure (AA))

{2-[3-(2-Cyclopentanecarbonyl-4-trifluoromethyl-phenyl)-ureido]-thiazol-4-yl}-acetic acid ethyl ester



10


To a stirred solution of (4-trifluoromethyl-phenyl)-carbamic acid tert-butyl ester (15 mmol) in Et₂O (40 mL) under a nitrogen atmosphere was added dropwise a 1.7 M solution of t-BuLi in pentan (38 mmol) over a 10-min period at -20 °C. The mixture was stirred at -15 °C for 1.5 h and then cyclopentanecarboxylic acid methoxy-methyl-amide (20 mmol) was added over 5 min. The mixture was allowed to warm to room temperature and stirred for further 1 h before it was quenched with aqueous NH₄Cl. The organic phase was isolated and the aqueous phase was extracted with CH₂Cl₂, and the combined organic phases were dried and concentrated *in vacuo*. The crude material was dissolved in CH₂Cl₂ (25 mL) and TFA (25 mL) and stirred at room temperature for 2 h. The reaction mixture was evaporated *in vacuo*, and added aqueous NaHCO₃ to pH 7 and extracted with CH₂Cl₂, dried and concentrated to give a crude product which was purified by flash chromatography (Quad flash 40, EtOAc- Heptane 0:1 ->1:3). This gave 70% of (2-amino-5-trifluoromethyl-phenyl)-cyclopentyl-methanone as an oil.

The title compound (15%) was prepared from (2-amino-5-trifluoromethyl-phenyl)-cyclopentyl-methanone and ethyl-2-amino-4-thiazolyl acetate following the general procedure AA.

¹H NMR (400MHz; CDCl₃): δ 1.28 (t, 3H), 1.64-1.99 (m, 8H), 3.70 (br t, 1H), 3.74 (s, 2H), 4.20 (q, 2H), 6.74 (s, 1H), 7.73 (d, 1H), 8.13 (s, 1H), 8.74 (br s, 1H), 9.53 (br s, 1H), 11.82 (br s, 1H); HPLC-MS : *m/z* = 492.1 (M+23); R_t = 5.02 min.

Example 476 (General procedure J)

{2-[3-(2-Cyclopentanecarbonyl-4-trifluoromethyl-phenyl)-ureido]-thiazol-4-yl}-acetic acid



5

The title compound (13 mg, 33%) was prepared from {2-[3-(2-cyclopentanecarbonyl-4-trifluoromethyl-phenyl)-ureido]-thiazol-4-yl}-acetic acid ethyl ester (42 mg, 0.09 mmol) following the general procedure J.


¹H NMR (400MHz; DMSO d₆): δ 1.65- 1.95 (m, 8H), 3.60 (s, 2H), 3.99 (br s, 1H), 6.94 (s, 1H), 7.90 (d, 1H), 8.25 (br s, 1H), 8.54 (br s, 1H), 10.94 (br s, 1H), 12.25 (br s, 2H); HPLC-MS : m/z = 426.1 (M+23); R_t = 4.49 min.

10

Example 477 (General procedure (AA))

{2-[3-(4-Chloro-2-cyclopentanecarbonyl-phenyl)-ureido]-thiazol-4-yl}-acetic acid ethyl ester

15



To a stirred solution of (4-chloro-phenyl)-carbamic acid tert-butyl ester (35 mmol) in Et₂O (80 mL) under a nitrogen atmosphere was added dropwise a 1.7 M solution of t-BuLi in pentan (88 mmol) over a 10-min period at -20 °C. The mixture was stirred at 0 °C for 1.5 h and then cyclopentanecarboxylic acid methoxy-methyl-amide (45 mmol) was added over 5 min. The mixture was allowed to warm to room temperature and stirred for further 1 h before it was quenched with aqueous NH₄Cl. The organic phase was isolated and the aqueous phase was extracted with CH₂Cl₂, and the combined organic phases were dried and concentrated *in vacuo*. The crude material was dissolved in CH₂Cl₂ (50 mL) and TFA (50 mL)

20

and stirred at room temperature for 1 h. The reaction mixture was evaporated *in vacuo*, and added aqueous NaHCO₃ to pH 7 and extracted with CH₂Cl₂, dried and concentrated to give a crude product which was purified by flash chromatography (Quad flash 65, EtOAc-Heptane 0:1 → 1:3). This gave 75% of (2-amino-5-chloro-phenyl)-cyclopentyl-methanone as


5 an oil.

The title compound (30%) was prepared from (2-amino-5-chloro-phenyl)-cyclopentyl-methanone and ethyl-2-amino-4-thiazolyl acetate following the general procedure AA.

¹H NMR (400MHz; CDCl₃): δ 1.28 (t, 3H), 1.62-1.99 (m, 8H), 3.3.67 (br t, 1H), 3.73 (s, 2H), 4.20 (q, 2H), 6.72 (s, 1H), 7.46 (dd, 1H), 7.84 (d, 1H), 8.56 (br s, 1H), 9.50 (br s, 1H), 11.58 (br s, 1H); HPLC-MS : *m/z* = 458.0 (M+23); R_t = 4.86 min.

Example 478 (General procedure J)

{2-[3-(4-Chloro-2-cyclopentanecarbonyl-phenyl)-ureido]-thiazol-4-yl}-acetic acid



15

The title compound (39 mg, 85%) was prepared from {2-[3-(4-chloro-2-cyclopentanecarbonyl-phenyl)-ureido]-thiazol-4-yl}-acetic acid ethyl ester (49 mg, 0.11 mmol) following the general procedure J.


¹H NMR (400MHz; DMSO d₆): δ 1.67- 1.92 (m, 8H), 3.59 (s, 2H), 3.89 (br s, 1H), 6.88 (s, 1H), 7.63 (d, 1H), 8.04 (br s, 1H), 8.32 (br s, 1H), 10.64 (s, 1H), 12.20 (br s, 2H); HPLC-MS : *m/z* = 430.0 (M+23); R_t = 4.11 min.

20

Example 479 (General procedure (AA))

{2-[3-(2-Cyclopentanecarbonyl-4-fluoro-phenyl)-ureido]-thiazol-4-yl}-acetic acid ethyl

25 ester



To a stirred solution of (4-fluoro-phenyl)-carbamic acid tert-butyl ester (9.5 mmol) in Et₂O (20 mL) under a nitrogen atmosphere was added dropwise a 1.7 M solution of t-BuLi in pentan (22 mmol) over a 10-min period at -30 °C. The mixture was stirred at -20 °C for 1.5 h and then cyclopentanecarboxylic acid methoxy-methyl-amide (10.5 mmol) was added over 5 min. The mixture was allowed to warm to room temperature and stirred for further 1 h before it was quenched with aqueous NH₄Cl. The organic phase was isolated and the aqueous phase was extracted with CH₂Cl₂, and the combined organic phases were dried and concentrated *in vacuo*. The crude material was dissolved in CH₂Cl₂ (25 mL) and TFA (25 mL) and stirred at room temperature for 2 h. The reaction mixture was evaporated *in vacuo*, and added aqueous NaHCO₃ to pH 7 and extracted with CH₂Cl₂, dried and concentrated to give a crude product which was purified by flash chromatography (Quad flash 40, EtOAc-Heptane 0:1 → 1:3). This gave 60% of (2-amino-5-fluoro-phenyl)-cyclopentyl-methanone as an oil.


The title compound (26%) was prepared from (2-amino-5-fluoro-phenyl)-cyclopentyl-methanone and ethyl-2-amino-4-thiazolyl acetate following the general procedure AA.

¹H NMR (400MHz; CDCl₃): δ 1.28 (t, 3H), 1.63-1.97 (m, 8H), 3.64 (br t, 1H), 3.73 (s, 2H), 4.20 (q, 2H), 6.72 (s, 1H), 7.22-7.29 (m, 1H), 7.58 (dd, 1H), 8.55 (br s, 1H), 9.40 (br s, 1H), 11.45 (br s, 1H); HPLC-MS : *m/z* = 442.0 (M+23); R_t = 4.49 min.

20

Example 480 (General procedure J)

{2-[3-(2-Cyclopentanecarbonyl-4-fluoro-phenyl)-ureido]-thiazol-4-yl}-acetic acid



The title compound (75 mg, 87%) was prepared from {2-[3-(2-cyclopentanecarbonyl-4-fluoro-phenyl)-ureido]-thiazol-4-yl}-acetic acid ethyl ester (92 mg, 0.22 mmol) following the general procedure J.

¹H NMR (400MHz; DMSO d₆): δ 1.65- 1.94 (m, 8H), 3.57 (s, 2H), 3.84 (br s, 1H),
5 6.88 (s, 1H), 7.45 (t, 1H), 7.85 (br s, 1H), 8.25 (br s, 1H), 10.48 (s, 1H), 12.10 (br s, 2H);
HPLC-MS : *m/z* = 414.1 (M+23); R_t = 3.74 min.

Using the methods disclosed herein the following compounds may also be made:

- 1-(5-Chloro-thiazol-2-yl)-3-(2-cyclopentanecarbonyl-4-methyl-phenyl)-urea
- 10 1-(5-Chloro-thiazol-2-yl)-3-(2-cyclopentanecarbonyl-phenyl)-urea
- 1-(4-Chloro-2-cyclopentanecarbonyl-phenyl)-3-(5-chloro-thiazol-2-yl)-urea
- 1-(5-Chloro-thiazol-2-yl)-3-(2-cyclopentanecarbonyl-4-fluoro-phenyl)-urea
- 1-(5-Chloro-thiazol-2-yl)-3-(2-cyclopentanecarbonyl-4-trifluoromethyl-phenyl)-urea
- 1-(5-Chloro-thiazol-2-yl)-3-(2-cyclopentanecarbonyl-4-methoxy-phenyl)-urea
- 15 1-(5-Chloro-thiazol-2-yl)-3-(2-cyclopentanecarbonyl-4-dimethylamino-phenyl)-urea
- 1-(5-Chloro-thiazol-2-yl)-3-(2-cyclopentanecarbonyl-4-morpholin-4-yl-phenyl)-urea
- 1-(5-Chloro-thiazol-2-yl)-3-(3-cyclopentanecarbonyl-pyridin-4-yl)-urea
- 1-(5-Chloro-thiazol-2-yl)-3-(4-cyclopentanecarbonyl-pyridin-3-yl)-urea
- 1-(5-Chloro-thiazol-2-yl)-3-(3-cyclopentanecarbonyl-pyridin-2-yl)-urea
- 20 1-(5-Bromo-thiazol-2-yl)-3-(2-cyclopentanecarbonyl-4-methyl-phenyl)-urea
- 1-(5-Bromo-thiazol-2-yl)-3-(2-cyclopentanecarbonyl-phenyl)-urea
- 1-(5-Bromo-thiazol-2-yl)-3-(4-chloro-2-cyclopentanecarbonyl-phenyl)-urea
- 1-(5-Bromo-thiazol-2-yl)-3-(2-cyclopentanecarbonyl-4-fluoro-phenyl)-urea
- 1-(5-Bromo-thiazol-2-yl)-3-(2-cyclopentanecarbonyl-4-trifluoromethyl-phenyl)-urea
- 25 1-(5-Bromo-thiazol-2-yl)-3-(2-cyclopentanecarbonyl-4-methoxy-phenyl)-urea
- 1-(5-Bromo-thiazol-2-yl)-3-(2-cyclopentanecarbonyl-4-dimethylamino-phenyl)-urea
- 1-(5-Bromo-thiazol-2-yl)-3-(2-cyclopentanecarbonyl-4-morpholin-4-yl-phenyl)-urea
- 1-(5-Bromo-thiazol-2-yl)-3-(3-cyclopentanecarbonyl-pyridin-4-yl)-urea
- 1-(5-Bromo-thiazol-2-yl)-3-(4-cyclopentanecarbonyl-pyridin-3-yl)-urea
- 30 1-(5-Bromo-thiazol-2-yl)-3-(3-cyclopentanecarbonyl-pyridin-2-yl)-urea
- 1-(2-Cyclopentanecarbonyl-4-methyl-phenyl)-3-(5-methylsulfanyl-thiazol-2-yl)-urea
- 1-(2-Cyclopentanecarbonyl-phenyl)-3-(5-methylsulfanyl-thiazol-2-yl)-urea
- 1-(4-Chloro-2-cyclopentanecarbonyl-phenyl)-3-(5-methylsulfanyl-thiazol-2-yl)-urea
- 1-(2-Cyclopentanecarbonyl-4-fluoro-phenyl)-3-(5-methylsulfanyl-thiazol-2-yl)-urea
- 35 1-(2-Cyclopentanecarbonyl-4-trifluoromethyl-phenyl)-3-(5-methylsulfanyl-thiazol-2-yl)-urea

- 1-(2-Cyclopentanecarbonyl-4-methoxy-phenyl)-3-(5-methylsulfanyl-thiazol-2-yl)-urea
1-(2-Cyclopentanecarbonyl-4-dimethylamino-phenyl)-3-(5-methylsulfanyl-thiazol-2-yl)-urea
1-(2-Cyclopentanecarbonyl-4-morpholin-4-yl-phenyl)-3-(5-methylsulfanyl-thiazol-2-yl)-urea
1-(3-Cyclopentanecarbonyl-pyridin-4-yl)-3-(5-methylsulfanyl-thiazol-2-yl)-urea
5 1-(4-Cyclopentanecarbonyl-pyridin-3-yl)-3-(5-methylsulfanyl-thiazol-2-yl)-urea
1-(3-Cyclopentanecarbonyl-pyridin-2-yl)-3-(5-methylsulfanyl-thiazol-2-yl)-urea
1-(2-Cyclopentanecarbonyl-4-methyl-phenyl)-3-(5-methanesulfonyl-thiazol-2-yl)-urea
1-(2-Cyclopentanecarbonyl-phenyl)-3-(5-methanesulfonyl-thiazol-2-yl)-urea
1-(4-Chloro-2-cyclopentanecarbonyl-phenyl)-3-(5-methanesulfonyl-thiazol-2-yl)-urea
10 1-(2-Cyclopentanecarbonyl-4-fluoro-phenyl)-3-(5-methanesulfonyl-thiazol-2-yl)-urea
1-(2-Cyclopentanecarbonyl-4-trifluoromethyl-phenyl)-3-(5-methanesulfonyl-thiazol-2-yl)-urea
1-(2-Cyclopentanecarbonyl-4-methoxy-phenyl)-3-(5-methanesulfonyl-thiazol-2-yl)-urea
1-(2-Cyclopentanecarbonyl-4-dimethylamino-phenyl)-3-(5-methanesulfonyl-thiazol-2-yl)-urea
1-(2-Cyclopentanecarbonyl-4-morpholin-4-yl-phenyl)-3-(5-methanesulfonyl-thiazol-2-yl)-urea
15 1-(3-Cyclopentanecarbonyl-pyridin-4-yl)-3-(5-methanesulfonyl-thiazol-2-yl)-urea
1-(4-Cyclopentanecarbonyl-pyridin-3-yl)-3-(5-methanesulfonyl-thiazol-2-yl)-urea
1-(3-Cyclopentanecarbonyl-pyridin-2-yl)-3-(5-methanesulfonyl-thiazol-2-yl)-urea
1-(2-Cyclopentanecarbonyl-4-methyl-phenyl)-3-[5-(pyridin-2-ylsulfanyl)-thiazol-2-yl]-urea
1-(2-Cyclopentanecarbonyl-phenyl)-3-[5-(pyridin-2-ylsulfanyl)-thiazol-2-yl]-urea
20 1-(4-Chloro-2-cyclopentanecarbonyl-phenyl)-3-[5-(pyridin-2-ylsulfanyl)-thiazol-2-yl]-urea
1-(2-Cyclopentanecarbonyl-4-fluoro-phenyl)-3-[5-(pyridin-2-ylsulfanyl)-thiazol-2-yl]-urea
1-(2-Cyclopentanecarbonyl-4-trifluoromethyl-phenyl)-3-[5-(pyridin-2-ylsulfanyl)-thiazol-2-yl]-
urea
1-(2-Cyclopentanecarbonyl-4-methoxy-phenyl)-3-[5-(pyridin-2-ylsulfanyl)-thiazol-2-yl]-urea
25 1-(2-Cyclopentanecarbonyl-4-dimethylamino-phenyl)-3-[5-(pyridin-2-ylsulfanyl)-thiazol-2-yl]-
urea
1-(2-Cyclopentanecarbonyl-4-morpholin-4-yl-phenyl)-3-[5-(pyridin-2-ylsulfanyl)-thiazol-2-yl]-
urea
1-(3-Cyclopentanecarbonyl-pyridin-4-yl)-3-[5-(pyridin-2-ylsulfanyl)-thiazol-2-yl]-urea
30 1-(4-Cyclopentanecarbonyl-pyridin-3-yl)-3-[5-(pyridin-2-ylsulfanyl)-thiazol-2-yl]-urea
1-(3-Cyclopentanecarbonyl-pyridin-2-yl)-3-[5-(pyridin-2-ylsulfanyl)-thiazol-2-yl]-urea
1-(2-Cyclopentanecarbonyl-4-methyl-phenyl)-3-[5-(pyridine-2-sulfonyl)-thiazol-2-yl]-urea
1-(2-Cyclopentanecarbonyl-phenyl)-3-[5-(pyridine-2-sulfonyl)-thiazol-2-yl]-urea
1-(4-Chloro-2-cyclopentanecarbonyl-phenyl)-3-[5-(pyridine-2-sulfonyl)-thiazol-2-yl]-urea
35 1-(2-Cyclopentanecarbonyl-4-fluoro-phenyl)-3-[5-(pyridine-2-sulfonyl)-thiazol-2-yl]-urea

- 1-(2-Cyclopentanecarbonyl-4-trifluoromethyl-phenyl)-3-[5-(pyridine-2-sulfonyl)-thiazol-2-yl]-urea
- 1-(2-Cyclopentanecarbonyl-4-methoxy-phenyl)-3-[5-(pyridine-2-sulfonyl)-thiazol-2-yl]-urea
- 1-(2-Cyclopentanecarbonyl-4-dimethylamino-phenyl)-3-[5-(pyridine-2-sulfonyl)-thiazol-2-yl]-urea
- 5 1-(2-Cyclopentanecarbonyl-4-morpholin-4-yl-phenyl)-3-[5-(pyridine-2-sulfonyl)-thiazol-2-yl]-urea
- 1-(3-Cyclopentanecarbonyl-pyridin-4-yl)-3-[5-(pyridine-2-sulfonyl)-thiazol-2-yl]-urea
- 1-(4-Cyclopentanecarbonyl-pyridin-3-yl)-3-[5-(pyridine-2-sulfonyl)-thiazol-2-yl]-urea
- 10 1-(3-Cyclopentanecarbonyl-pyridin-2-yl)-3-[5-(pyridine-2-sulfonyl)-thiazol-2-yl]-urea
- 1-(2-Cyclopentanecarbonyl-4-methyl-phenyl)-3-[5-(1*H*-imidazol-2-ylsulfanyl)-thiazol-2-yl]-urea
- 1-(2-Cyclopentanecarbonyl-phenyl)-3-[5-(1*H*-imidazol-2-ylsulfanyl)-thiazol-2-yl]-urea
- 1-(4-Chloro-2-cyclopentanecarbonyl-phenyl)-3-[5-(1*H*-imidazol-2-ylsulfanyl)-thiazol-2-yl]-urea
- 1-(2-Cyclopentanecarbonyl-4-fluoro-phenyl)-3-[5-(1*H*-imidazol-2-ylsulfanyl)-thiazol-2-yl]-urea
- 15 1-(2-Cyclopentanecarbonyl-4-trifluoromethyl-phenyl)-3-[5-(1*H*-imidazol-2-ylsulfanyl)-thiazol-2-yl]-urea
- 1-(2-Cyclopentanecarbonyl-4-methoxy-phenyl)-3-[5-(1*H*-imidazol-2-ylsulfanyl)-thiazol-2-yl]-urea
- 1-(2-Cyclopentanecarbonyl-4-dimethylamino-phenyl)-3-[5-(1*H*-imidazol-2-ylsulfanyl)-thiazol-2-yl]-urea
- 20 1-(2-Cyclopentanecarbonyl-4-morpholin-4-yl-phenyl)-3-[5-(1*H*-imidazol-2-ylsulfanyl)-thiazol-2-yl]-urea
- 1-(3-Cyclopentanecarbonyl-pyridin-4-yl)-3-[5-(1*H*-imidazol-2-ylsulfanyl)-thiazol-2-yl]-urea
- 1-(4-Cyclopentanecarbonyl-pyridin-3-yl)-3-[5-(1*H*-imidazol-2-ylsulfanyl)-thiazol-2-yl]-urea
- 25 1-(3-Cyclopentanecarbonyl-pyridin-2-yl)-3-[5-(1*H*-imidazol-2-ylsulfanyl)-thiazol-2-yl]-urea
- 1-(2-Cyclopentanecarbonyl-4-methyl-phenyl)-3-[5-(1*H*-imidazole-2-sulfonyl)-thiazol-2-yl]-urea
- 1-(2-Cyclopentanecarbonyl-phenyl)-3-[5-(1*H*-imidazole-2-sulfonyl)-thiazol-2-yl]-urea
- 1-(4-Chloro-2-cyclopentanecarbonyl-phenyl)-3-[5-(1*H*-imidazole-2-sulfonyl)-thiazol-2-yl]-urea
- 1-(2-Cyclopentanecarbonyl-4-fluoro-phenyl)-3-[5-(1*H*-imidazole-2-sulfonyl)-thiazol-2-yl]-urea
- 30 1-(2-Cyclopentanecarbonyl-4-trifluoromethyl-phenyl)-3-[5-(1*H*-imidazole-2-sulfonyl)-thiazol-2-yl]-urea
- 1-(2-Cyclopentanecarbonyl-4-methoxy-phenyl)-3-[5-(1*H*-imidazole-2-sulfonyl)-thiazol-2-yl]-urea
- 1-(2-Cyclopentanecarbonyl-4-dimethylamino-phenyl)-3-[5-(1*H*-imidazole-2-sulfonyl)-thiazol-2-yl]-urea
- 35 2-yl]-urea

- 1-(2-Cyclopentanecarbonyl-4-morpholin-4-yl-phenyl)-3-[5-(1*H*-imidazole-2-sulfonyl)-thiazol-2-yl]-urea
- 1-(3-Cyclopentanecarbonyl-pyridin-4-yl)-3-[5-(1*H*-imidazole-2-sulfonyl)-thiazol-2-yl]-urea
- 1-(4-Cyclopentanecarbonyl-pyridin-3-yl)-3-[5-(1*H*-imidazole-2-sulfonyl)-thiazol-2-yl]-urea
- 5 1-(3-Cyclopentanecarbonyl-pyridin-2-yl)-3-[5-(1*H*-imidazole-2-sulfonyl)-thiazol-2-yl]-urea
- 1-(2-Cyclopentanecarbonyl-4-methyl-phenyl)-3-[5-(1-methyl-1*H*-tetrazol-5-ylsulfanyl)-thiazol-2-yl]-urea
- 1-(2-Cyclopentanecarbonyl-phenyl)-3-[5-(1-methyl-1*H*-tetrazol-5-ylsulfanyl)-thiazol-2-yl]-urea
- 1-(4-Chloro-2-cyclopentanecarbonyl-phenyl)-3-[5-(1-methyl-1*H*-tetrazol-5-ylsulfanyl)-thiazol-2-yl]-urea
- 10 2-yl]-urea
- 1-(2-Cyclopentanecarbonyl-4-fluoro-phenyl)-3-[5-(1-methyl-1*H*-tetrazol-5-ylsulfanyl)-thiazol-2-yl]-urea
- 1-(2-Cyclopentanecarbonyl-4-trifluoromethyl-phenyl)-3-[5-(1-methyl-1*H*-tetrazol-5-ylsulfanyl)-thiazol-2-yl]-urea
- 15 1-(2-Cyclopentanecarbonyl-4-methoxy-phenyl)-3-[5-(1-methyl-1*H*-tetrazol-5-ylsulfanyl)-thiazol-2-yl]-urea
- 1-(2-Cyclopentanecarbonyl-4-dimethylamino-phenyl)-3-[5-(1-methyl-1*H*-tetrazol-5-ylsulfanyl)-thiazol-2-yl]-urea
- 1-(2-Cyclopentanecarbonyl-4-morpholin-4-yl-phenyl)-3-[5-(1-methyl-1*H*-tetrazol-5-ylsulfanyl)-thiazol-2-yl]-urea
- 20 thiazol-2-yl]-urea
- 1-(3-Cyclopentanecarbonyl-pyridin-4-yl)-3-[5-(1-methyl-1*H*-tetrazol-5-ylsulfanyl)-thiazol-2-yl]-urea
- 1-(4-Cyclopentanecarbonyl-pyridin-3-yl)-3-[5-(1-methyl-1*H*-tetrazol-5-ylsulfanyl)-thiazol-2-yl]-urea
- 25 1-(3-Cyclopentanecarbonyl-pyridin-2-yl)-3-[5-(1-methyl-1*H*-tetrazol-5-ylsulfanyl)-thiazol-2-yl]-urea
- 1-(2-Cyclopentanecarbonyl-4-methyl-phenyl)-3-[5-(1-methyl-1*H*-tetrazole-5-sulfonyl)-thiazol-2-yl]-urea
- 1-(2-Cyclopentanecarbonyl-phenyl)-3-[5-(1-methyl-1*H*-tetrazole-5-sulfonyl)-thiazol-2-yl]-urea
- 30 1-(4-Chloro-2-cyclopentanecarbonyl-phenyl)-3-[5-(1-methyl-1*H*-tetrazole-5-sulfonyl)-thiazol-2-yl]-urea
- 1-(2-Cyclopentanecarbonyl-4-fluoro-phenyl)-3-[5-(1-methyl-1*H*-tetrazole-5-sulfonyl)-thiazol-2-yl]-urea
- 1-(2-Cyclopentanecarbonyl-4-trifluoromethyl-phenyl)-3-[5-(1-methyl-1*H*-tetrazole-5-sulfonyl)-thiazol-2-yl]-urea
- 35 thiazol-2-yl]-urea

- 1-(2-Cyclopentanecarbonyl-4-methoxy-phenyl)-3-[5-(1-methyl-1*H*-tetrazole-5-sulfonyl)-thiazol-2-yl]-urea
- 1-(2-Cyclopentanecarbonyl-4-dimethylamino-phenyl)-3-[5-(1-methyl-1*H*-tetrazole-5-sulfonyl)-thiazol-2-yl]-urea
- 5 1-(2-Cyclopentanecarbonyl-4-morpholin-4-yl-phenyl)-3-[5-(1-methyl-1*H*-tetrazole-5-sulfonyl)-thiazol-2-yl]-urea
- 1-(3-Cyclopentanecarbonyl-pyridin-4-yl)-3-[5-(1-methyl-1*H*-tetrazole-5-sulfonyl)-thiazol-2-yl]-urea.
- 1-(4-Cyclopentanecarbonyl-pyridin-3-yl)-3-[5-(1-methyl-1*H*-tetrazole-5-sulfonyl)-thiazol-2-yl]-urea
- 10 1-(3-Cyclopentanecarbonyl-pyridin-2-yl)-3-[5-(1-methyl-1*H*-tetrazole-5-sulfonyl)-thiazol-2-yl]-urea
- 1-(2-Cyclopentanecarbonyl-4-methyl-phenyl)-3-[5-(1*H*-tetrazole-5-sulfonyl)-thiazol-2-yl]-urea
- 1-(2-Cyclopentanecarbonyl-phenyl)-3-[5-(1*H*-tetrazole-5-sulfonyl)-thiazol-2-yl]-urea
- 15 1-(4-Chloro-2-cyclopentanecarbonyl-phenyl)-3-[5-(1*H*-tetrazole-5-sulfonyl)-thiazol-2-yl]-urea
- 1-(2-Cyclopentanecarbonyl-4-fluoro-phenyl)-3-[5-(1*H*-tetrazole-5-sulfonyl)-thiazol-2-yl]-urea
- 1-(2-Cyclopentanecarbonyl-4-trifluoromethyl-phenyl)-3-[5-(1*H*-tetrazole-5-sulfonyl)-thiazol-2-yl]-urea
- 1-(2-Cyclopentanecarbonyl-4-methoxy-phenyl)-3-[5-(1*H*-tetrazole-5-sulfonyl)-thiazol-2-yl]-urea
- 20 1-(2-Cyclopentanecarbonyl-4-dimethylamino-phenyl)-3-[5-(1*H*-tetrazole-5-sulfonyl)-thiazol-2-yl]-urea
- 1-(2-Cyclopentanecarbonyl-4-morpholin-4-yl-phenyl)-3-[5-(1*H*-tetrazole-5-sulfonyl)-thiazol-2-yl]-urea
- 25 1-(3-Cyclopentanecarbonyl-pyridin-4-yl)-3-[5-(1*H*-tetrazole-5-sulfonyl)-thiazol-2-yl]-urea
- 1-(4-Cyclopentanecarbonyl-pyridin-3-yl)-3-[5-(1*H*-tetrazole-5-sulfonyl)-thiazol-2-yl]-urea
- 1-(3-Cyclopentanecarbonyl-pyridin-2-yl)-3-[5-(1*H*-tetrazole-5-sulfonyl)-thiazol-2-yl]-urea
- 5-Chloro-2-[3-(2-cyclopentanecarbonyl-4-methyl-phenyl)-ureido]-thiazole-4-carboxylic acid
- 5-Chloro-2-[3-(2-cyclopentanecarbonyl-phenyl)-ureido]-thiazole-4-carboxylic acid
- 30 5-Chloro-2-[3-(4-chloro-2-cyclopentanecarbonyl-phenyl)-ureido]-thiazole-4-carboxylic acid
- 5-Chloro-2-[3-(2-cyclopentanecarbonyl-4-fluoro-phenyl)-ureido]-thiazole-4-carboxylic acid
- 5-Chloro-2-[3-(2-cyclopentanecarbonyl-4-trifluoromethyl-phenyl)-ureido]-thiazole-4-carboxylic acid
- 5-Chloro-2-[3-(2-cyclopentanecarbonyl-4-methoxy-phenyl)-ureido]-thiazole-4-carboxylic acid

- 5-Chloro-2-[3-(2-cyclopentanecarbonyl-4-dimethylamino-phenyl)-ureido]-thiazole-4-carboxylic acid
- 5-Chloro-2-[3-(2-cyclopentanecarbonyl-4-morpholin-4-yl-phenyl)-ureido]-thiazole-4-carboxylic acid
- 5 5-Chloro-2-[3-(3-cyclopentanecarbonyl-pyridin-4-yl)-ureido]-thiazole-4-carboxylic acid
- 5-Chloro-2-[3-(4-cyclopentanecarbonyl-pyridin-3-yl)-ureido]-thiazole-4-carboxylic acid
- 5-Chloro-2-[3-(3-cyclopentanecarbonyl-pyridin-2-yl)-ureido]-thiazole-4-carboxylic acid
- 5-Bromo-2-[3-(2-cyclopentanecarbonyl-4-methyl-phenyl)-ureido]-thiazole-4-carboxylic acid
- 5-Bromo-2-[3-(2-cyclopentanecarbonyl-phenyl)-ureido]-thiazole-4-carboxylic acid
- 10 5-Bromo-2-[3-(4-chloro-2-cyclopentanecarbonyl-phenyl)-ureido]-thiazole-4-carboxylic acid
- 5-Bromo-2-[3-(2-cyclopentanecarbonyl-4-fluoro-phenyl)-ureido]-thiazole-4-carboxylic acid
- 5-Bromo-2-[3-(2-cyclopentanecarbonyl-4-trifluoromethyl-phenyl)-ureido]-thiazole-4-carboxylic acid
- 5-Bromo-2-[3-(2-cyclopentanecarbonyl-4-methoxy-phenyl)-ureido]-thiazole-4-carboxylic acid
- 15 5-Bromo-2-[3-(2-cyclopentanecarbonyl-4-dimethylamino-phenyl)-ureido]-thiazole-4-carboxylic acid
- 5-Bromo-2-[3-(2-cyclopentanecarbonyl-4-morpholin-4-yl-phenyl)-ureido]-thiazole-4-carboxylic acid
- 5-Bromo-2-[3-(3-cyclopentanecarbonyl-pyridin-4-yl)-ureido]-thiazole-4-carboxylic acid
- 20 5-Bromo-2-[3-(4-cyclopentanecarbonyl-pyridin-3-yl)-ureido]-thiazole-4-carboxylic acid
- 5-Bromo-2-[3-(3-cyclopentanecarbonyl-pyridin-2-yl)-ureido]-thiazole-4-carboxylic acid
- 2-[3-(2-Cyclopentanecarbonyl-4-methyl-phenyl)-ureido]-5-methylsulfanyl-thiazole-4-carboxylic acid
- 2-[3-(2-Cyclopentanecarbonyl-phenyl)-ureido]-5-methylsulfanyl-thiazole-4-carboxylic acid
- 25 2-[3-(4-Chloro-2-cyclopentanecarbonyl-phenyl)-ureido]-5-methylsulfanyl-thiazole-4-carboxylic acid
- 2-[3-(2-Cyclopentanecarbonyl-4-fluoro-phenyl)-ureido]-5-methylsulfanyl-thiazole-4-carboxylic acid
- 2-[3-(2-Cyclopentanecarbonyl-4-trifluoromethyl-phenyl)-ureido]-5-methylsulfanyl-thiazole-4-carboxylic acid
- 30 2-[3-(2-Cyclopentanecarbonyl-4-methoxy-phenyl)-ureido]-5-methylsulfanyl-thiazole-4-carboxylic acid
- 2-[3-(2-Cyclopentanecarbonyl-4-dimethylamino-phenyl)-ureido]-5-methylsulfanyl-thiazole-4-carboxylic acid

- 2-[3-(2-Cyclopentanecarbonyl-4-morpholin-4-yl-phenyl)-ureido]-5-methylsulfanyl-thiazole-4-carboxylic acid
- 2-[3-(3-Cyclopentanecarbonyl-pyridin-4-yl)-ureido]-5-methylsulfanyl-thiazole-4-carboxylic acid
- 5 2-[3-(4-Cyclopentanecarbonyl-pyridin-3-yl)-ureido]-5-methylsulfanyl-thiazole-4-carboxylic acid
- 2-[3-(3-Cyclopentanecarbonyl-pyridin-2-yl)-ureido]-5-methylsulfanyl-thiazole-4-carboxylic acid
- 2-[3-(2-Cyclopentanecarbonyl-4-methyl-phenyl)-ureido]-5-methanesulfonyl-thiazole-4-carboxylic acid
- 10 2-[3-(2-Cyclopentanecarbonyl-phenyl)-ureido]-5-methanesulfonyl-thiazole-4-carboxylic acid
- 2-[3-(4-Chloro-2-cyclopentanecarbonyl-phenyl)-ureido]-5-methanesulfonyl-thiazole-4-carboxylic acid
- 2-[3-(2-Cyclopentanecarbonyl-4-fluoro-phenyl)-ureido]-5-methanesulfonyl-thiazole-4-carboxylic acid
- 15 2-[3-(2-Cyclopentanecarbonyl-4-trifluoromethyl-phenyl)-ureido]-5-methanesulfonyl-thiazole-4-carboxylic acid
- 2-[3-(2-Cyclopentanecarbonyl-4-methoxy-phenyl)-ureido]-5-methanesulfonyl-thiazole-4-carboxylic acid
- 20 2-[3-(2-Cyclopentanecarbonyl-4-dimethylamino-phenyl)-ureido]-5-methanesulfonyl-thiazole-4-carboxylic acid
- 2-[3-(2-Cyclopentanecarbonyl-4-morpholin-4-yl-phenyl)-ureido]-5-methanesulfonyl-thiazole-4-carboxylic acid
- 2-[3-(3-Cyclopentanecarbonyl-pyridin-4-yl)-ureido]-5-methanesulfonyl-thiazole-4-carboxylic acid
- 25 2-[3-(4-Cyclopentanecarbonyl-pyridin-3-yl)-ureido]-5-methanesulfonyl-thiazole-4-carboxylic acid
- 2-[3-(3-Cyclopentanecarbonyl-pyridin-2-yl)-ureido]-5-methanesulfonyl-thiazole-4-carboxylic acid
- 30 2-[3-(2-Cyclopentanecarbonyl-4-methyl-phenyl)-ureido]-5-(pyridin-2-ylsulfanyl)-thiazole-4-carboxylic acid
- 2-[3-(2-Cyclopentanecarbonyl-phenyl)-ureido]-5-(pyridin-2-ylsulfanyl)-thiazole-4-carboxylic acid
- 2-[3-(4-Chloro-2-cyclopentanecarbonyl-phenyl)-ureido]-5-(pyridin-2-ylsulfanyl)-thiazole-4-carboxylic acid
- 35

- 2-[3-(2-Cyclopentanecarbonyl-4-fluoro-phenyl)-ureido]-5-(pyridin-2-ylsulfanyl)-thiazole-4-carboxylic acid
- 2-[3-(2-Cyclopentanecarbonyl-4-trifluoromethyl-phenyl)-ureido]-5-(pyridin-2-ylsulfanyl)-thiazole-4-carboxylic acid
- 5 2-[3-(2-Cyclopentanecarbonyl-4-methoxy-phenyl)-ureido]-5-(pyridin-2-ylsulfanyl)-thiazole-4-carboxylic acid
- 2-[3-(2-Cyclopentanecarbonyl-4-dimethylamino-phenyl)-ureido]-5-(pyridin-2-ylsulfanyl)-thiazole-4-carboxylic acid
- 2-[3-(2-Cyclopentanecarbonyl-4-morpholin-4-yl-phenyl)-ureido]-5-(pyridin-2-ylsulfanyl)-thiazole-4-carboxylic acid
- 10 2-[3-(3-Cyclopentanecarbonyl-pyridin-4-yl)-ureido]-5-(pyridin-2-ylsulfanyl)-thiazole-4-carboxylic acid
- 2-[3-(4-Cyclopentanecarbonyl-pyridin-3-yl)-ureido]-5-(pyridin-2-ylsulfanyl)-thiazole-4-carboxylic acid
- 15 2-[3-(3-Cyclopentanecarbonyl-pyridin-2-yl)-ureido]-5-(pyridin-2-ylsulfanyl)-thiazole-4-carboxylic acid
- 2-[3-(2-Cyclopentanecarbonyl-4-methyl-phenyl)-ureido]-5-(pyridine-2-sulfonyl)-thiazole-4-carboxylic acid
- 2-[3-(2-Cyclopentanecarbonyl-phenyl)-ureido]-5-(pyridine-2-sulfonyl)-thiazole-4-carboxylic acid
- 20 acid
- 2-[3-(4-Chloro-2-cyclopentanecarbonyl-phenyl)-ureido]-5-(pyridine-2-sulfonyl)-thiazole-4-carboxylic acid
- 2-[3-(2-Cyclopentanecarbonyl-4-fluoro-phenyl)-ureido]-5-(pyridine-2-sulfonyl)-thiazole-4-carboxylic acid
- 25 2-[3-(2-Cyclopentanecarbonyl-4-trifluoromethyl-phenyl)-ureido]-5-(pyridine-2-sulfonyl)-thiazole-4-carboxylic acid
- 2-[3-(2-Cyclopentanecarbonyl-4-methoxy-phenyl)-ureido]-5-(pyridine-2-sulfonyl)-thiazole-4-carboxylic acid
- 2-[3-(2-Cyclopentanecarbonyl-4-dimethylamino-phenyl)-ureido]-5-(pyridine-2-sulfonyl)-thiazole-4-carboxylic acid
- 30 thiazole-4-carboxylic acid
- 2-[3-(2-Cyclopentanecarbonyl-4-morpholin-4-yl-phenyl)-ureido]-5-(pyridine-2-sulfonyl)-thiazole-4-carboxylic acid
- 2-[3-(3-Cyclopentanecarbonyl-pyridin-4-yl)-ureido]-5-(pyridine-2-sulfonyl)-thiazole-4-carboxylic acid

- 2-[3-(4-Cyclopentanecarbonyl-pyridin-3-yl)-ureido]-5-(pyridine-2-sulfonyl)-thiazole-4-carboxylic acid
- 2-[3-(3-Cyclopentanecarbonyl-pyridin-2-yl)-ureido]-5-(pyridine-2-sulfonyl)-thiazole-4-carboxylic acid
- 5 2-[3-(2-Cyclopentanecarbonyl-4-methyl-phenyl)-ureido]-5-(1*H*-imidazol-2-ylsulfanyl)-thiazole-4-carboxylic acid
- 2-[3-(2-Cyclopentanecarbonyl-phenyl)-ureido]-5-(1*H*-imidazol-2-ylsulfanyl)-thiazole-4-carboxylic acid
- 2-[3-(4-Chloro-2-cyclopentanecarbonyl-phenyl)-ureido]-5-(1*H*-imidazol-2-ylsulfanyl)-thiazole-
- 10 4-carboxylic acid
- 2-[3-(2-Cyclopentanecarbonyl-4-fluoro-phenyl)-ureido]-5-(1*H*-imidazol-2-ylsulfanyl)-thiazole-4-carboxylic acid
- 2-[3-(2-Cyclopentanecarbonyl-4-trifluoromethyl-phenyl)-ureido]-5-(1*H*-imidazol-2-ylsulfanyl)-thiazole-4-carboxylic acid
- 15 2-[3-(2-Cyclopentanecarbonyl-4-methoxy-phenyl)-ureido]-5-(1*H*-imidazol-2-ylsulfanyl)-thiazole-4-carboxylic acid
- 2-[3-(2-Cyclopentanecarbonyl-4-dimethylamino-phenyl)-ureido]-5-(1*H*-imidazol-2-ylsulfanyl)-thiazole-4-carboxylic acid
- 2-[3-(2-Cyclopentanecarbonyl-4-morpholin-4-yl-phenyl)-ureido]-5-(1*H*-imidazol-2-ylsulfanyl)-
- 20 thiazole-4-carboxylic acid
- 2-[3-(3-Cyclopentanecarbonyl-pyridin-4-yl)-ureido]-5-(1*H*-imidazol-2-ylsulfanyl)-thiazole-4-carboxylic acid
- 2-[3-(4-Cyclopentanecarbonyl-pyridin-3-yl)-ureido]-5-(1*H*-imidazol-2-ylsulfanyl)-thiazole-4-carboxylic acid
- 25 2-[3-(3-Cyclopentanecarbonyl-pyridin-2-yl)-ureido]-5-(1*H*-imidazol-2-ylsulfanyl)-thiazole-4-carboxylic acid
- 2-[3-(2-Cyclopentanecarbonyl-4-methyl-phenyl)-ureido]-5-(1*H*-imidazole-2-sulfonyl)-thiazole-4-carboxylic acid
- 2-[3-(2-Cyclopentanecarbonyl-phenyl)-ureido]-5-(1*H*-imidazole-2-sulfonyl)-thiazole-4-
- 30 carboxylic acid
- 2-[3-(4-Chloro-2-cyclopentanecarbonyl-phenyl)-ureido]-5-(1*H*-imidazole-2-sulfonyl)-thiazole-4-carboxylic acid
- 2-[3-(2-Cyclopentanecarbonyl-4-fluoro-phenyl)-ureido]-5-(1*H*-imidazole-2-sulfonyl)-thiazole-4-carboxylic acid

- 2-[3-(2-Cyclopentanecarbonyl-4-trifluoromethyl-phenyl)-ureido]-5-(1*H*-imidazole-2-sulfonyl)-thiazole-4-carboxylic acid
- 2-[3-(2-Cyclopentanecarbonyl-4-methoxy-phenyl)-ureido]-5-(1*H*-imidazole-2-sulfonyl)-thiazole-4-carboxylic acid
- 5 2-[3-(2-Cyclopentanecarbonyl-4-dimethylamino-phenyl)-ureido]-5-(1*H*-imidazole-2-sulfonyl)-thiazole-4-carboxylic acid
- 2-[3-(2-Cyclopentanecarbonyl-4-morpholin-4-yl-phenyl)-ureido]-5-(1*H*-imidazole-2-sulfonyl)-thiazole-4-carboxylic acid
- 2-[3-(3-Cyclopentanecarbonyl-pyridin-4-yl)-ureido]-5-(1*H*-imidazole-2-sulfonyl)-thiazole-4-
- 10 carboxylic acid
- 2-[3-(4-Cyclopentanecarbonyl-pyridin-3-yl)-ureido]-5-(1*H*-imidazole-2-sulfonyl)-thiazole-4-carboxylic acid
- 2-[3-(3-Cyclopentanecarbonyl-pyridin-2-yl)-ureido]-5-(1*H*-imidazole-2-sulfonyl)-thiazole-4-carboxylic acid
- 15 2-[3-(2-Cyclopentanecarbonyl-4-methyl-phenyl)-ureido]-5-(1-methyl-1*H*-tetrazol-5-ylsulfanyl)-thiazole-4-carboxylic acid
- 2-[3-(2-Cyclopentanecarbonyl-phenyl)-ureido]-5-(1-methyl-1*H*-tetrazol-5-ylsulfanyl)-thiazole-4-carboxylic acid
- 2-[3-(4-Chloro-2-cyclopentanecarbonyl-phenyl)-ureido]-5-(1-methyl-1*H*-tetrazol-5-ylsulfanyl)-
- 20 thiazole-4-carboxylic acid
- 2-[3-(2-Cyclopentanecarbonyl-4-fluoro-phenyl)-ureido]-5-(1-methyl-1*H*-tetrazol-5-ylsulfanyl)-thiazole-4-carboxylic acid
- 2-[3-(2-Cyclopentanecarbonyl-4-trifluoromethyl-phenyl)-ureido]-5-(1-methyl-1*H*-tetrazol-5-ylsulfanyl)-thiazole-4-carboxylic acid
- 25 2-[3-(2-Cyclopentanecarbonyl-4-methoxy-phenyl)-ureido]-5-(1-methyl-1*H*-tetrazol-5-ylsulfanyl)-thiazole-4-carboxylic acid
- 2-[3-(2-Cyclopentanecarbonyl-4-dimethylamino-phenyl)-ureido]-5-(1-methyl-1*H*-tetrazol-5-ylsulfanyl)-thiazole-4-carboxylic acid
- 2-[3-(2-Cyclopentanecarbonyl-4-morpholin-4-yl-phenyl)-ureido]-5-(1-methyl-1*H*-tetrazol-5-
- 30 ylsulfanyl)-thiazole-4-carboxylic acid
- 2-[3-(3-Cyclopentanecarbonyl-pyridin-4-yl)-ureido]-5-(1-methyl-1*H*-tetrazol-5-ylsulfanyl)-thiazole-4-carboxylic acid
- 2-[3-(4-Cyclopentanecarbonyl-pyridin-3-yl)-ureido]-5-(1-methyl-1*H*-tetrazol-5-ylsulfanyl)-thiazole-4-carboxylic acid

- 2-[3-(3-Cyclopentanecarbonyl-pyridin-2-yl)-ureido]-5-(1-methyl-1*H*-tetrazol-5-ylsulfanyl)-thiazole-4-carboxylic acid
- 2-[3-(2-Cyclopentanecarbonyl-4-methyl-phenyl)-ureido]-5-(1-methyl-1*H*-tetrazole-5-sulfonyl)-thiazole-4-carboxylic acid
- 5 2-[3-(2-Cyclopentanecarbonyl-phenyl)-ureido]-5-(1-methyl-1*H*-tetrazole-5-sulfonyl)-thiazole-4-carboxylic acid
- 2-[3-(4-Chloro-2-cyclopentanecarbonyl-phenyl)-ureido]-5-(1-methyl-1*H*-tetrazole-5-sulfonyl)-thiazole-4-carboxylic acid
- 2-[3-(2-Cyclopentanecarbonyl-4-fluoro-phenyl)-ureido]-5-(1-methyl-1*H*-tetrazole-5-sulfonyl)-thiazole-4-carboxylic acid
- 10 2-[3-(2-Cyclopentanecarbonyl-4-trifluoromethyl-phenyl)-ureido]-5-(1-methyl-1*H*-tetrazole-5-sulfonyl)-thiazole-4-carboxylic acid
- 2-[3-(2-Cyclopentanecarbonyl-4-methoxy-phenyl)-ureido]-5-(1-methyl-1*H*-tetrazole-5-sulfonyl)-thiazole-4-carboxylic acid
- 15 2-[3-(2-Cyclopentanecarbonyl-4-dimethylamino-phenyl)-ureido]-5-(1-methyl-1*H*-tetrazole-5-sulfonyl)-thiazole-4-carboxylic acid
- 2-[3-(2-Cyclopentanecarbonyl-4-morpholin-4-yl-phenyl)-ureido]-5-(1-methyl-1*H*-tetrazole-5-sulfonyl)-thiazole-4-carboxylic acid
- 2-[3-(3-Cyclopentanecarbonyl-pyridin-4-yl)-ureido]-5-(1-methyl-1*H*-tetrazole-5-sulfonyl)-thiazole-4-carboxylic acid
- 20 2-[3-(4-Cyclopentanecarbonyl-pyridin-3-yl)-ureido]-5-(1-methyl-1*H*-tetrazole-5-sulfonyl)-thiazole-4-carboxylic acid
- 2-[3-(3-Cyclopentanecarbonyl-pyridin-2-yl)-ureido]-5-(1-methyl-1*H*-tetrazole-5-sulfonyl)-thiazole-4-carboxylic acid
- 25 2-[3-(2-Cyclopentanecarbonyl-4-methyl-phenyl)-ureido]-5-(1*H*-tetrazole-5-sulfonyl)-thiazole-4-carboxylic acid
- 2-[3-(2-Cyclopentanecarbonyl-phenyl)-ureido]-5-(1*H*-tetrazole-5-sulfonyl)-thiazole-4-carboxylic acid
- 2-[3-(4-Chloro-2-cyclopentanecarbonyl-phenyl)-ureido]-5-(1*H*-tetrazole-5-sulfonyl)-thiazole-4-carboxylic acid
- 30 2-[3-(2-Cyclopentanecarbonyl-4-fluoro-phenyl)-ureido]-5-(1*H*-tetrazole-5-sulfonyl)-thiazole-4-carboxylic acid
- 2-[3-(2-Cyclopentanecarbonyl-4-trifluoromethyl-phenyl)-ureido]-5-(1*H*-tetrazole-5-sulfonyl)-thiazole-4-carboxylic acid

- 2-[3-(2-Cyclopentanecarbonyl-4-methoxy-phenyl)-ureido]-5-(1*H*-tetrazole-5-sulfonyl)-thiazole-4-carboxylic acid
- 2-[3-(2-Cyclopentanecarbonyl-4-dimethylamino-phenyl)-ureido]-5-(1*H*-tetrazole-5-sulfonyl)-thiazole-4-carboxylic acid
- 5 2-[3-(2-Cyclopentanecarbonyl-4-morpholin-4-yl-phenyl)-ureido]-5-(1*H*-tetrazole-5-sulfonyl)-thiazole-4-carboxylic acid
- 2-[3-(3-Cyclopentanecarbonyl-pyridin-4-yl)-ureido]-5-(1*H*-tetrazole-5-sulfonyl)-thiazole-4-carboxylic acid
- 2-[3-(4-Cyclopentanecarbonyl-pyridin-3-yl)-ureido]-5-(1*H*-tetrazole-5-sulfonyl)-thiazole-4-carboxylic acid
- 10 2-[3-(3-Cyclopentanecarbonyl-pyridin-2-yl)-ureido]-5-(1*H*-tetrazole-5-sulfonyl)-thiazole-4-carboxylic acid
- {5-Chloro-2-[3-(2-cyclopentanecarbonyl-4-methyl-phenyl)-ureido]-thiazol-4-yl}-acetic acid
- {5-Chloro-2-[3-(2-cyclopentanecarbonyl-phenyl)-ureido]-thiazol-4-yl}-acetic acid
- 15 {5-Chloro-2-[3-(4-chloro-2-cyclopentanecarbonyl-phenyl)-ureido]-thiazol-4-yl}-acetic acid
- {5-Chloro-2-[3-(2-cyclopentanecarbonyl-4-fluoro-phenyl)-ureido]-thiazol-4-yl}-acetic acid
- {5-Chloro-2-[3-(2-cyclopentanecarbonyl-4-trifluoromethyl-phenyl)-ureido]-thiazol-4-yl}-acetic acid
- {5-Chloro-2-[3-(2-cyclopentanecarbonyl-4-methoxy-phenyl)-ureido]-thiazol-4-yl}-acetic acid
- 20 {5-Chloro-2-[3-(2-cyclopentanecarbonyl-4-dimethylamino-phenyl)-ureido]-thiazol-4-yl}-acetic acid
- {5-Chloro-2-[3-(2-cyclopentanecarbonyl-4-morpholin-4-yl-phenyl)-ureido]-thiazol-4-yl}-acetic acid
- {5-Chloro-2-[3-(3-cyclopentanecarbonyl-pyridin-4-yl)-ureido]-thiazol-4-yl}-acetic acid
- 25 {5-Chloro-2-[3-(4-cyclopentanecarbonyl-pyridin-3-yl)-ureido]-thiazol-4-yl}-acetic acid
- {5-Chloro-2-[3-(3-cyclopentanecarbonyl-pyridin-2-yl)-ureido]-thiazol-4-yl}-acetic acid
- {5-Bromo-2-[3-(2-cyclopentanecarbonyl-4-methyl-phenyl)-ureido]-thiazol-4-yl}-acetic acid
- {5-Bromo-2-[3-(2-cyclopentanecarbonyl-phenyl)-ureido]-thiazol-4-yl}-acetic acid
- {5-Bromo-2-[3-(4-chloro-2-cyclopentanecarbonyl-phenyl)-ureido]-thiazol-4-yl}-acetic acid
- 30 {5-Bromo-2-[3-(2-cyclopentanecarbonyl-4-fluoro-phenyl)-ureido]-thiazol-4-yl}-acetic acid
- {5-Bromo-2-[3-(2-cyclopentanecarbonyl-4-trifluoromethyl-phenyl)-ureido]-thiazol-4-yl}-acetic acid
- {5-Bromo-2-[3-(2-cyclopentanecarbonyl-4-methoxy-phenyl)-ureido]-thiazol-4-yl}-acetic acid
- {5-Bromo-2-[3-(2-cyclopentanecarbonyl-4-dimethylamino-phenyl)-ureido]-thiazol-4-yl}-acetic acid
- 35 acid

- {5-Bromo-2-[3-(2-cyclopentanecarbonyl-4-morpholin-4-yl-phenyl)-ureido]-thiazol-4-yl}-acetic acid
- {5-Bromo-2-[3-(3-cyclopentanecarbonyl-pyridin-4-yl)-ureido]-thiazol-4-yl}-acetic acid
- {5-Bromo-2-[3-(4-cyclopentanecarbonyl-pyridin-3-yl)-ureido]-thiazol-4-yl}-acetic acid
- 5 {5-Bromo-2-[3-(3-cyclopentanecarbonyl-pyridin-2-yl)-ureido]-thiazol-4-yl}-acetic acid
- {2-[3-(2-Cyclopentanecarbonyl-4-methyl-phenyl)-ureido]-5-methylsulfanyl-thiazol-4-yl}-acetic acid
- {2-[3-(2-Cyclopentanecarbonyl-phenyl)-ureido]-5-methylsulfanyl-thiazol-4-yl}-acetic acid
- {2-[3-(4-Chloro-2-cyclopentanecarbonyl-phenyl)-ureido]-5-methylsulfanyl-thiazol-4-yl}-acetic acid
- 10 acid
- {2-[3-(2-Cyclopentanecarbonyl-4-fluoro-phenyl)-ureido]-5-methylsulfanyl-thiazol-4-yl}-acetic acid
- acid
- {2-[3-(2-Cyclopentanecarbonyl-4-trifluoromethyl-phenyl)-ureido]-5-methylsulfanyl-thiazol-4-yl}-acetic acid
- 15 {2-[3-(2-Cyclopentanecarbonyl-4-methoxy-phenyl)-ureido]-5-methylsulfanyl-thiazol-4-yl}-acetic acid
- acid
- {2-[3-(2-Cyclopentanecarbonyl-4-dimethylamino-phenyl)-ureido]-5-methylsulfanyl-thiazol-4-yl}-acetic acid
- {2-[3-(2-Cyclopentanecarbonyl-4-morpholin-4-yl-phenyl)-ureido]-5-methylsulfanyl-thiazol-4-yl}-acetic acid
- 20 yl}-acetic acid
- {2-[3-(3-Cyclopentanecarbonyl-pyridin-4-yl)-ureido]-5-methylsulfanyl-thiazol-4-yl}-acetic acid
- {2-[3-(4-Cyclopentanecarbonyl-pyridin-3-yl)-ureido]-5-methylsulfanyl-thiazol-4-yl}-acetic acid
- {2-[3-(3-Cyclopentanecarbonyl-pyridin-2-yl)-ureido]-5-methylsulfanyl-thiazol-4-yl}-acetic acid
- {2-[3-(2-Cyclopentanecarbonyl-4-methyl-phenyl)-ureido]-5-methanesulfonyl-thiazol-4-yl}-acetic acid
- 25 acid
- {2-[3-(2-Cyclopentanecarbonyl-phenyl)-ureido]-5-methanesulfonyl-thiazol-4-yl}-acetic acid
- {2-[3-(4-Chloro-2-cyclopentanecarbonyl-phenyl)-ureido]-5-methanesulfonyl-thiazol-4-yl}-acetic acid
- acid
- {2-[3-(2-Cyclopentanecarbonyl-4-fluoro-phenyl)-ureido]-5-methanesulfonyl-thiazol-4-yl}-acetic acid
- 30 acid
- {2-[3-(2-Cyclopentanecarbonyl-4-trifluoromethyl-phenyl)-ureido]-5-methanesulfonyl-thiazol-4-yl}-acetic acid
- {2-[3-(2-Cyclopentanecarbonyl-4-methoxy-phenyl)-ureido]-5-methanesulfonyl-thiazol-4-yl}-acetic acid

- {2-[3-(2-Cyclopentanecarbonyl-4-dimethylamino-phenyl)-ureido]-5-methanesulfonyl-thiazol-4-yl}-acetic acid
- {2-[3-(2-Cyclopentanecarbonyl-4-morpholin-4-yl-phenyl)-ureido]-5-methanesulfonyl-thiazol-4-yl}-acetic acid
- 5 {2-[3-(3-Cyclopentanecarbonyl-pyridin-4-yl)-ureido]-5-methanesulfonyl-thiazol-4-yl}-acetic acid
- {2-[3-(4-Cyclopentanecarbonyl-pyridin-3-yl)-ureido]-5-methanesulfonyl-thiazol-4-yl}-acetic acid
- {2-[3-(3-Cyclopentanecarbonyl-pyridin-2-yl)-ureido]-5-methanesulfonyl-thiazol-4-yl}-acetic acid
- 10 acid
- {2-[3-(2-Cyclopentanecarbonyl-4-methyl-phenyl)-ureido]-5-(pyridin-2-ylsulfanyl)-thiazol-4-yl}-acetic acid
- {2-[3-(2-Cyclopentanecarbonyl-phenyl)-ureido]-5-(pyridin-2-ylsulfanyl)-thiazol-4-yl}-acetic acid
- {2-[3-(4-Chloro-2-cyclopentanecarbonyl-phenyl)-ureido]-5-(pyridin-2-ylsulfanyl)-thiazol-4-yl}-acetic acid
- 15 acid
- {2-[3-(2-Cyclopentanecarbonyl-4-fluoro-phenyl)-ureido]-5-(pyridin-2-ylsulfanyl)-thiazol-4-yl}-acetic acid
- {2-[3-(2-Cyclopentanecarbonyl-4-trifluoromethyl-phenyl)-ureido]-5-(pyridin-2-ylsulfanyl)-thiazol-4-yl}-acetic acid
- 20 {2-[3-(2-Cyclopentanecarbonyl-4-methoxy-phenyl)-ureido]-5-(pyridin-2-ylsulfanyl)-thiazol-4-yl}-acetic acid
- {2-[3-(2-Cyclopentanecarbonyl-4-dimethylamino-phenyl)-ureido]-5-(pyridin-2-ylsulfanyl)-thiazol-4-yl}-acetic acid
- {2-[3-(2-Cyclopentanecarbonyl-4-morpholin-4-yl-phenyl)-ureido]-5-(pyridin-2-ylsulfanyl)-thiazol-4-yl}-acetic acid
- 25 acid
- {2-[3-(3-Cyclopentanecarbonyl-pyridin-4-yl)-ureido]-5-(pyridin-2-ylsulfanyl)-thiazol-4-yl}-acetic acid
- {2-[3-(4-Cyclopentanecarbonyl-pyridin-3-yl)-ureido]-5-(pyridin-2-ylsulfanyl)-thiazol-4-yl}-acetic acid
- 30 acid
- {2-[3-(3-Cyclopentanecarbonyl-pyridin-2-yl)-ureido]-5-(pyridin-2-ylsulfanyl)-thiazol-4-yl}-acetic acid
- {2-[3-(2-Cyclopentanecarbonyl-4-methyl-phenyl)-ureido]-5-(pyridine-2-sulfonyl)-thiazol-4-yl}-acetic acid
- {2-[3-(2-Cyclopentanecarbonyl-phenyl)-ureido]-5-(pyridine-2-sulfonyl)-thiazol-4-yl}-acetic acid

- [2-[3-(4-Chloro-2-cyclopentanecarbonyl-phenyl)-ureido]-5-(pyridine-2-sulfonyl)-thiazol-4-yl]-acetic acid
- [2-[3-(2-Cyclopentanecarbonyl-4-fluoro-phenyl)-ureido]-5-(pyridine-2-sulfonyl)-thiazol-4-yl]-acetic acid
- 5 [2-[3-(2-Cyclopentanecarbonyl-4-trifluoromethyl-phenyl)-ureido]-5-(pyridine-2-sulfonyl)-thiazol-4-yl]-acetic acid
- [2-[3-(2-Cyclopentanecarbonyl-4-methoxy-phenyl)-ureido]-5-(pyridine-2-sulfonyl)-thiazol-4-yl]-acetic acid
- [2-[3-(2-Cyclopentanecarbonyl-4-dimethylamino-phenyl)-ureido]-5-(pyridine-2-sulfonyl)-thiazol-4-yl]-acetic acid
- 10 [2-[3-(2-Cyclopentanecarbonyl-4-morpholin-4-yl-phenyl)-ureido]-5-(pyridine-2-sulfonyl)-thiazol-4-yl]-acetic acid
- [2-[3-(3-Cyclopentanecarbonyl-pyridin-4-yl)-ureido]-5-(pyridine-2-sulfonyl)-thiazol-4-yl]-acetic acid
- [2-[3-(4-Cyclopentanecarbonyl-pyridin-3-yl)-ureido]-5-(pyridine-2-sulfonyl)-thiazol-4-yl]-acetic acid
- 15 [2-[3-(3-Cyclopentanecarbonyl-pyridin-2-yl)-ureido]-5-(pyridine-2-sulfonyl)-thiazol-4-yl]-acetic acid
- [2-[3-(2-Cyclopentanecarbonyl-4-methyl-phenyl)-ureido]-5-(1*H*-imidazol-2-ylsulfanyl)-thiazol-4-yl]-acetic acid
- 20 [2-[3-(2-Cyclopentanecarbonyl-phenyl)-ureido]-5-(1*H*-imidazol-2-ylsulfanyl)-thiazol-4-yl]-acetic acid
- [2-[3-(4-Chloro-2-cyclopentanecarbonyl-phenyl)-ureido]-5-(1*H*-imidazol-2-ylsulfanyl)-thiazol-4-yl]-acetic acid
- 25 [2-[3-(2-Cyclopentanecarbonyl-4-fluoro-phenyl)-ureido]-5-(1*H*-imidazol-2-ylsulfanyl)-thiazol-4-yl]-acetic acid
- [2-[3-(2-Cyclopentanecarbonyl-4-trifluoromethyl-phenyl)-ureido]-5-(1*H*-imidazol-2-ylsulfanyl)-thiazol-4-yl]-acetic acid
- [2-[3-(2-Cyclopentanecarbonyl-4-methoxy-phenyl)-ureido]-5-(1*H*-imidazol-2-ylsulfanyl)-thiazol-4-yl]-acetic acid
- 30 [2-[3-(2-Cyclopentanecarbonyl-4-dimethylamino-phenyl)-ureido]-5-(1*H*-imidazol-2-ylsulfanyl)-thiazol-4-yl]-acetic acid
- [2-[3-(2-Cyclopentanecarbonyl-4-morpholin-4-yl-phenyl)-ureido]-5-(1*H*-imidazol-2-ylsulfanyl)-thiazol-4-yl]-acetic acid

- [2-[3-(3-Cyclopentanecarbonyl-pyridin-4-yl)-ureido]-5-(1*H*-imidazol-2-ylsulfanyl)-thiazol-4-yl]-acetic acid
- [2-[3-(4-Cyclopentanecarbonyl-pyridin-3-yl)-ureido]-5-(1*H*-imidazol-2-ylsulfanyl)-thiazol-4-yl]-acetic acid
- 5 [2-[3-(3-Cyclopentanecarbonyl-pyridin-2-yl)-ureido]-5-(1*H*-imidazol-2-ylsulfanyl)-thiazol-4-yl]-acetic acid
- [2-[3-(2-Cyclopentanecarbonyl-4-methyl-phenyl)-ureido]-5-(1*H*-imidazole-2-sulfonyl)-thiazol-4-yl]-acetic acid
- [2-[3-(2-Cyclopentanecarbonyl-phenyl)-ureido]-5-(1*H*-imidazole-2-sulfonyl)-thiazol-4-yl]-acetic acid
- 10 acid
- [2-[3-(4-Chloro-2-cyclopentanecarbonyl-phenyl)-ureido]-5-(1*H*-imidazole-2-sulfonyl)-thiazol-4-yl]-acetic acid
- [2-[3-(2-Cyclopentanecarbonyl-4-fluoro-phenyl)-ureido]-5-(1*H*-imidazole-2-sulfonyl)-thiazol-4-yl]-acetic acid
- 15 [2-[3-(2-Cyclopentanecarbonyl-4-trifluoromethyl-phenyl)-ureido]-5-(1*H*-imidazole-2-sulfonyl)-thiazol-4-yl]-acetic acid
- [2-[3-(2-Cyclopentanecarbonyl-4-methoxy-phenyl)-ureido]-5-(1*H*-imidazole-2-sulfonyl)-thiazol-4-yl]-acetic acid
- [2-[3-(2-Cyclopentanecarbonyl-4-dimethylamino-phenyl)-ureido]-5-(1*H*-imidazole-2-sulfonyl)-thiazol-4-yl]-acetic acid
- 20 thiazol-4-yl]-acetic acid
- [2-[3-(2-Cyclopentanecarbonyl-4-morpholin-4-yl-phenyl)-ureido]-5-(1*H*-imidazole-2-sulfonyl)-thiazol-4-yl]-acetic acid
- [2-[3-(3-Cyclopentanecarbonyl-pyridin-4-yl)-ureido]-5-(1*H*-imidazole-2-sulfonyl)-thiazol-4-yl]-acetic acid
- 25 [2-[3-(4-Cyclopentanecarbonyl-pyridin-3-yl)-ureido]-5-(1*H*-imidazole-2-sulfonyl)-thiazol-4-yl]-acetic acid
- [2-[3-(3-Cyclopentanecarbonyl-pyridin-2-yl)-ureido]-5-(1*H*-imidazole-2-sulfonyl)-thiazol-4-yl]-acetic acid
- [2-[3-(2-Cyclopentanecarbonyl-4-methyl-phenyl)-ureido]-5-(1-methyl-1*H*-tetrazol-5-ylsulfanyl)-thiazol-4-yl]-acetic acid
- 30 thiazol-4-yl]-acetic acid
- [2-[3-(2-Cyclopentanecarbonyl-phenyl)-ureido]-5-(1-methyl-1*H*-tetrazol-5-ylsulfanyl)-thiazol-4-yl]-acetic acid
- [2-[3-(4-Chloro-2-cyclopentanecarbonyl-phenyl)-ureido]-5-(1-methyl-1*H*-tetrazol-5-ylsulfanyl)-thiazol-4-yl]-acetic acid

- [2-[3-(2-Cyclopentanecarbonyl-4-fluoro-phenyl)-ureido]-5-(1-methyl-1*H*-tetrazol-5-ylsulfanyl)-thiazol-4-yl]-acetic acid
- [2-[3-(2-Cyclopentanecarbonyl-4-trifluoromethyl-phenyl)-ureido]-5-(1-methyl-1*H*-tetrazol-5-ylsulfanyl)-thiazol-4-yl]-acetic acid
- 5 [2-[3-(2-Cyclopentanecarbonyl-4-methoxy-phenyl)-ureido]-5-(1-methyl-1*H*-tetrazol-5-ylsulfanyl)-thiazol-4-yl]-acetic acid
- [2-[3-(2-Cyclopentanecarbonyl-4-dimethylamino-phenyl)-ureido]-5-(1-methyl-1*H*-tetrazol-5-ylsulfanyl)-thiazol-4-yl]-acetic acid
- [2-[3-(2-Cyclopentanecarbonyl-4-morpholin-4-yl-phenyl)-ureido]-5-(1-methyl-1*H*-tetrazol-5-ylsulfanyl)-thiazol-4-yl]-acetic acid
- 10 [2-[3-(3-Cyclopentanecarbonyl-pyridin-4-yl)-ureido]-5-(1-methyl-1*H*-tetrazol-5-ylsulfanyl)-thiazol-4-yl]-acetic acid
- [2-[3-(4-Cyclopentanecarbonyl-pyridin-3-yl)-ureido]-5-(1-methyl-1*H*-tetrazol-5-ylsulfanyl)-thiazol-4-yl]-acetic acid
- [2-[3-(3-Cyclopentanecarbonyl-pyridin-2-yl)-ureido]-5-(1-methyl-1*H*-tetrazol-5-ylsulfanyl)-thiazol-4-yl]-acetic acid
- 15 [2-[3-(2-Cyclopentanecarbonyl-4-methyl-phenyl)-ureido]-5-(1-methyl-1*H*-tetrazole-5-sulfonyl)-thiazol-4-yl]-acetic acid
- [2-[3-(2-Cyclopentanecarbonyl-phenyl)-ureido]-5-(1-methyl-1*H*-tetrazole-5-sulfonyl)-thiazol-4-yl]-acetic acid
- 20 [2-[3-(4-Chloro-2-cyclopentanecarbonyl-phenyl)-ureido]-5-(1-methyl-1*H*-tetrazole-5-sulfonyl)-thiazol-4-yl]-acetic acid
- [2-[3-(2-Cyclopentanecarbonyl-4-fluoro-phenyl)-ureido]-5-(1-methyl-1*H*-tetrazole-5-sulfonyl)-thiazol-4-yl]-acetic acid
- [2-[3-(2-Cyclopentanecarbonyl-4-trifluoromethyl-phenyl)-ureido]-5-(1-methyl-1*H*-tetrazole-5-sulfonyl)-thiazol-4-yl]-acetic acid
- 25 [2-[3-(2-Cyclopentanecarbonyl-4-methoxy-phenyl)-ureido]-5-(1-methyl-1*H*-tetrazole-5-sulfonyl)-thiazol-4-yl]-acetic acid
- [2-[3-(2-Cyclopentanecarbonyl-4-dimethylamino-phenyl)-ureido]-5-(1-methyl-1*H*-tetrazole-5-sulfonyl)-thiazol-4-yl]-acetic acid
- 30 [2-[3-(2-Cyclopentanecarbonyl-4-morpholin-4-yl-phenyl)-ureido]-5-(1-methyl-1*H*-tetrazole-5-sulfonyl)-thiazol-4-yl]-acetic acid
- [2-[3-(3-Cyclopentanecarbonyl-pyridin-4-yl)-ureido]-5-(1-methyl-1*H*-tetrazole-5-sulfonyl)-thiazol-4-yl]-acetic acid

- [2-[3-(4-Cyclopentanecarbonyl-pyridin-3-yl)-ureido]-5-(1-methyl-1*H*-tetrazole-5-sulfonyl)-thiazol-4-yl]-acetic acid
- [2-[3-(3-Cyclopentanecarbonyl-pyridin-2-yl)-ureido]-5-(1-methyl-1*H*-tetrazole-5-sulfonyl)-thiazol-4-yl]-acetic acid
- 5 [2-[3-(2-Cyclopentanecarbonyl-4-methyl-phenyl)-ureido]-5-(1*H*-tetrazole-5-sulfonyl)-thiazol-4-yl]-acetic acid
- [2-[3-(2-Cyclopentanecarbonyl-phenyl)-ureido]-5-(1*H*-tetrazole-5-sulfonyl)-thiazol-4-yl]-acetic acid
- [2-[3-(4-Chloro-2-cyclopentanecarbonyl-phenyl)-ureido]-5-(1*H*-tetrazole-5-sulfonyl)-thiazol-4-yl]-acetic acid
- 10 [2-[3-(2-Cyclopentanecarbonyl-4-fluoro-phenyl)-ureido]-5-(1*H*-tetrazole-5-sulfonyl)-thiazol-4-yl]-acetic acid
- [2-[3-(2-Cyclopentanecarbonyl-4-trifluoromethyl-phenyl)-ureido]-5-(1*H*-tetrazole-5-sulfonyl)-thiazol-4-yl]-acetic acid
- 15 [2-[3-(2-Cyclopentanecarbonyl-4-methoxy-phenyl)-ureido]-5-(1*H*-tetrazole-5-sulfonyl)-thiazol-4-yl]-acetic acid
- [2-[3-(2-Cyclopentanecarbonyl-4-dimethylamino-phenyl)-ureido]-5-(1*H*-tetrazole-5-sulfonyl)-thiazol-4-yl]-acetic acid
- [2-[3-(2-Cyclopentanecarbonyl-4-morpholin-4-yl-phenyl)-ureido]-5-(1*H*-tetrazole-5-sulfonyl)-thiazol-4-yl]-acetic acid
- 20 [2-[3-(3-Cyclopentanecarbonyl-pyridin-4-yl)-ureido]-5-(1*H*-tetrazole-5-sulfonyl)-thiazol-4-yl]-acetic acid
- [2-[3-(4-Cyclopentanecarbonyl-pyridin-3-yl)-ureido]-5-(1*H*-tetrazole-5-sulfonyl)-thiazol-4-yl]-acetic acid
- 25 [2-[3-(3-Cyclopentanecarbonyl-pyridin-2-yl)-ureido]-5-(1*H*-tetrazole-5-sulfonyl)-thiazol-4-yl]-acetic acid
- 1-(5-Chloro-4-phenylsulfanylmethyl-thiazol-2-yl)-3-(2-cyclopentanecarbonyl-4-methyl-phenyl)-urea
- 1-(5-Chloro-4-phenylsulfanylmethyl-thiazol-2-yl)-3-(2-cyclopentanecarbonyl-phenyl)-urea
- 30 1-(4-Chloro-2-cyclopentanecarbonyl-phenyl)-3-(5-chloro-4-phenylsulfanylmethyl-thiazol-2-yl)-urea
- 1-(5-Chloro-4-phenylsulfanylmethyl-thiazol-2-yl)-3-(2-cyclopentanecarbonyl-4-fluoro-phenyl)-urea
- 1-(5-Chloro-4-phenylsulfanylmethyl-thiazol-2-yl)-3-(2-cyclopentanecarbonyl-4-trifluoromethyl-phenyl)-urea
- 35

- 1-(5-Chloro-4-phenylsulfanylmethyl-thiazol-2-yl)-3-(2-cyclopentanecarbonyl-4-methoxy-phenyl)-urea
- 1-(5-Chloro-4-phenylsulfanylmethyl-thiazol-2-yl)-3-(2-cyclopentanecarbonyl-4-dimethylamino-phenyl)-urea
- 5 1-(5-Chloro-4-phenylsulfanylmethyl-thiazol-2-yl)-3-(2-cyclopentanecarbonyl-4-morpholin-4-yl-phenyl)-urea
- 1-(5-Chloro-4-phenylsulfanylmethyl-thiazol-2-yl)-3-(3-cyclopentanecarbonyl-pyridin-4-yl)-urea
- 1-(5-Chloro-4-phenylsulfanylmethyl-thiazol-2-yl)-3-(4-cyclopentanecarbonyl-pyridin-3-yl)-urea
- 1-(5-Chloro-4-phenylsulfanylmethyl-thiazol-2-yl)-3-(3-cyclopentanecarbonyl-pyridin-2-yl)-urea
- 10 1-(5-Bromo-4-phenylsulfanylmethyl-thiazol-2-yl)-3-(2-cyclopentanecarbonyl-4-methyl-phenyl)-urea
- 1-(5-Bromo-4-phenylsulfanylmethyl-thiazol-2-yl)-3-(2-cyclopentanecarbonyl-phenyl)-urea
- 1-(5-Bromo-4-phenylsulfanylmethyl-thiazol-2-yl)-3-(4-chloro-2-cyclopentanecarbonyl-phenyl)-urea
- 15 1-(5-Bromo-4-phenylsulfanylmethyl-thiazol-2-yl)-3-(2-cyclopentanecarbonyl-4-fluoro-phenyl)-urea
- 1-(5-Bromo-4-phenylsulfanylmethyl-thiazol-2-yl)-3-(2-cyclopentanecarbonyl-4-trifluoromethyl-phenyl)-urea
- 1-(5-Bromo-4-phenylsulfanylmethyl-thiazol-2-yl)-3-(2-cyclopentanecarbonyl-4-methoxy-phenyl)-urea
- 20 1-(5-Bromo-4-phenylsulfanylmethyl-thiazol-2-yl)-3-(2-cyclopentanecarbonyl-4-dimethylamino-phenyl)-urea
- 1-(5-Bromo-4-phenylsulfanylmethyl-thiazol-2-yl)-3-(2-cyclopentanecarbonyl-4-morpholin-4-yl-phenyl)-urea
- 25 1-(5-Bromo-4-phenylsulfanylmethyl-thiazol-2-yl)-3-(3-cyclopentanecarbonyl-pyridin-4-yl)-urea
- 1-(5-Bromo-4-phenylsulfanylmethyl-thiazol-2-yl)-3-(4-cyclopentanecarbonyl-pyridin-3-yl)-urea
- 1-(5-Bromo-4-phenylsulfanylmethyl-thiazol-2-yl)-3-(3-cyclopentanecarbonyl-pyridin-2-yl)-urea
- 1-(2-Cyclopentanecarbonyl-4-methyl-phenyl)-3-(5-methylsulfanyl-4-phenylsulfanylmethyl-thiazol-2-yl)-urea
- 30 1-(2-Cyclopentanecarbonyl-phenyl)-3-(5-methylsulfanyl-4-phenylsulfanylmethyl-thiazol-2-yl)-urea
- 1-(4-Chloro-2-cyclopentanecarbonyl-phenyl)-3-(5-methylsulfanyl-4-phenylsulfanylmethyl-thiazol-2-yl)-urea
- 1-(2-Cyclopentanecarbonyl-4-fluoro-phenyl)-3-(5-methylsulfanyl-4-phenylsulfanylmethyl-thiazol-2-yl)-urea
- 35

- 1-(2-Cyclopentanecarbonyl-4-trifluoromethyl-phenyl)-3-(5-methylsulfanyl-4-phenylsulfanylmethyl-thiazol-2-yl)-urea
- 1-(2-Cyclopentanecarbonyl-4-methoxy-phenyl)-3-(5-methylsulfanyl-4-phenylsulfanylmethyl-thiazol-2-yl)-urea
- 5 1-(2-Cyclopentanecarbonyl-4-dimethylamino-phenyl)-3-(5-methylsulfanyl-4-phenylsulfanylmethyl-thiazol-2-yl)-urea
- 1-(2-Cyclopentanecarbonyl-4-morpholin-4-yl-phenyl)-3-(5-methylsulfanyl-4-phenylsulfanylmethyl-thiazol-2-yl)-urea
- 1-(3-Cyclopentanecarbonyl-pyridin-4-yl)-3-(5-methylsulfanyl-4-phenylsulfanylmethyl-thiazol-2-yl)-urea
- 10 2-yl)-urea
- 1-(4-Cyclopentanecarbonyl-pyridin-3-yl)-3-(5-methylsulfanyl-4-phenylsulfanylmethyl-thiazol-2-yl)-urea
- 1-(3-Cyclopentanecarbonyl-pyridin-2-yl)-3-(5-methylsulfanyl-4-phenylsulfanylmethyl-thiazol-2-yl)-urea
- 15 1-(2-Cyclopentanecarbonyl-4-methyl-phenyl)-3-(5-methanesulfonyl-4-phenylsulfanylmethyl-thiazol-2-yl)-urea
- 1-(2-Cyclopentanecarbonyl-phenyl)-3-(5-methanesulfonyl-4-phenylsulfanylmethyl-thiazol-2-yl)-urea
- 1-(4-Chloro-2-cyclopentanecarbonyl-phenyl)-3-(5-methanesulfonyl-4-phenylsulfanylmethyl-thiazol-2-yl)-urea
- 20 1-(2-Cyclopentanecarbonyl-4-fluoro-phenyl)-3-(5-methanesulfonyl-4-phenylsulfanylmethyl-thiazol-2-yl)-urea
- 1-(2-Cyclopentanecarbonyl-4-trifluoromethyl-phenyl)-3-(5-methanesulfonyl-4-phenylsulfanylmethyl-thiazol-2-yl)-urea
- 25 1-(2-Cyclopentanecarbonyl-4-methoxy-phenyl)-3-(5-methanesulfonyl-4-phenylsulfanylmethyl-thiazol-2-yl)-urea
- 1-(2-Cyclopentanecarbonyl-4-dimethylamino-phenyl)-3-(5-methanesulfonyl-4-phenylsulfanylmethyl-thiazol-2-yl)-urea
- 1-(2-Cyclopentanecarbonyl-4-morpholin-4-yl-phenyl)-3-(5-methanesulfonyl-4-phenylsulfanylmethyl-thiazol-2-yl)-urea
- 30 phenylsulfanylmethyl-thiazol-2-yl)-urea
- 1-(3-Cyclopentanecarbonyl-pyridin-4-yl)-3-(5-methanesulfonyl-4-phenylsulfanylmethyl-thiazol-2-yl)-urea
- 1-(4-Cyclopentanecarbonyl-pyridin-3-yl)-3-(5-methanesulfonyl-4-phenylsulfanylmethyl-thiazol-2-yl)-urea

- 1-(3-Cyclopentanecarbonyl-pyridin-2-yl)-3-(5-methanesulfonyl-4-phenylsulfanylmethyl-thiazol-2-yl)-urea
- 1-(2-Cyclopentanecarbonyl-4-methyl-phenyl)-3-[4-phenylsulfanylmethyl-5-(pyridin-2-ylsulfanyl)-thiazol-2-yl]-urea
- 5 1-(2-Cyclopentanecarbonyl-phenyl)-3-[4-phenylsulfanylmethyl-5-(pyridin-2-ylsulfanyl)-thiazol-2-yl]-urea
- 1-(4-Chloro-2-cyclopentanecarbonyl-phenyl)-3-[4-phenylsulfanylmethyl-5-(pyridin-2-ylsulfanyl)-thiazol-2-yl]-urea
- 1-(2-Cyclopentanecarbonyl-4-fluoro-phenyl)-3-[4-phenylsulfanylmethyl-5-(pyridin-2-ylsulfanyl)-thiazol-2-yl]-urea
- 10 1-(2-Cyclopentanecarbonyl-4-trifluoromethyl-phenyl)-3-[4-phenylsulfanylmethyl-5-(pyridin-2-ylsulfanyl)-thiazol-2-yl]-urea
- 1-(2-Cyclopentanecarbonyl-4-methoxy-phenyl)-3-[4-phenylsulfanylmethyl-5-(pyridin-2-ylsulfanyl)-thiazol-2-yl]-urea
- 15 1-(2-Cyclopentanecarbonyl-4-dimethylamino-phenyl)-3-[4-phenylsulfanylmethyl-5-(pyridin-2-ylsulfanyl)-thiazol-2-yl]-urea
- 1-(2-Cyclopentanecarbonyl-4-morpholin-4-yl-phenyl)-3-[4-phenylsulfanylmethyl-5-(pyridin-2-ylsulfanyl)-thiazol-2-yl]-urea
- 1-(3-Cyclopentanecarbonyl-pyridin-4-yl)-3-[4-phenylsulfanylmethyl-5-(pyridin-2-ylsulfanyl)-thiazol-2-yl]-urea
- 20 1-(4-Cyclopentanecarbonyl-pyridin-3-yl)-3-[4-phenylsulfanylmethyl-5-(pyridin-2-ylsulfanyl)-thiazol-2-yl]-urea
- 1-(3-Cyclopentanecarbonyl-pyridin-2-yl)-3-[4-phenylsulfanylmethyl-5-(pyridin-2-ylsulfanyl)-thiazol-2-yl]-urea
- 25 1-(2-Cyclopentanecarbonyl-4-methyl-phenyl)-3-[4-phenylsulfanylmethyl-5-(pyridine-2-sulfonyl)-thiazol-2-yl]-urea
- 1-(2-Cyclopentanecarbonyl-phenyl)-3-[4-phenylsulfanylmethyl-5-(pyridine-2-sulfonyl)-thiazol-2-yl]-urea
- 1-(4-Chloro-2-cyclopentanecarbonyl-phenyl)-3-[4-phenylsulfanylmethyl-5-(pyridine-2-sulfonyl)-thiazol-2-yl]-urea
- 30 1-(2-Cyclopentanecarbonyl-4-fluoro-phenyl)-3-[4-phenylsulfanylmethyl-5-(pyridine-2-sulfonyl)-thiazol-2-yl]-urea
- 1-(2-Cyclopentanecarbonyl-4-trifluoromethyl-phenyl)-3-[4-phenylsulfanylmethyl-5-(pyridine-2-sulfonyl)-thiazol-2-yl]-urea

- 1-(2-Cyclopentanecarbonyl-4-methoxy-phenyl)-3-[4-phenylsulfanylmethyl-5-(pyridine-2-sulfonyl)-thiazol-2-yl]-urea
- 1-(2-Cyclopentanecarbonyl-4-dimethylamino-phenyl)-3-[4-phenylsulfanylmethyl-5-(pyridine-2-sulfonyl)-thiazol-2-yl]-urea
- 5 1-(2-Cyclopentanecarbonyl-4-morpholin-4-yl-phenyl)-3-[4-phenylsulfanylmethyl-5-(pyridine-2-sulfonyl)-thiazol-2-yl]-urea
- 1-(3-Cyclopentanecarbonyl-pyridin-4-yl)-3-[4-phenylsulfanylmethyl-5-(pyridine-2-sulfonyl)-thiazol-2-yl]-urea
- 1-(4-Cyclopentanecarbonyl-pyridin-3-yl)-3-[4-phenylsulfanylmethyl-5-(pyridine-2-sulfonyl)-thiazol-2-yl]-urea
- 10 1-(3-Cyclopentanecarbonyl-pyridin-2-yl)-3-[4-phenylsulfanylmethyl-5-(pyridine-2-sulfonyl)-thiazol-2-yl]-urea
- 1-(2-Cyclopentanecarbonyl-4-methyl-phenyl)-3-[5-(1*H*-imidazol-2-ylsulfanyl)-4-phenylsulfanylmethyl-thiazol-2-yl]-urea
- 15 1-(2-Cyclopentanecarbonyl-phenyl)-3-[5-(1*H*-imidazol-2-ylsulfanyl)-4-phenylsulfanylmethyl-thiazol-2-yl]-urea
- 1-(4-Chloro-2-cyclopentanecarbonyl-phenyl)-3-[5-(1*H*-imidazol-2-ylsulfanyl)-4-phenylsulfanylmethyl-thiazol-2-yl]-urea
- 1-(2-Cyclopentanecarbonyl-4-fluoro-phenyl)-3-[5-(1*H*-imidazol-2-ylsulfanyl)-4-phenylsulfanylmethyl-thiazol-2-yl]-urea
- 20 1-(2-Cyclopentanecarbonyl-4-trifluoromethyl-phenyl)-3-[5-(1*H*-imidazol-2-ylsulfanyl)-4-phenylsulfanylmethyl-thiazol-2-yl]-urea
- 1-(2-Cyclopentanecarbonyl-4-methoxy-phenyl)-3-[5-(1*H*-imidazol-2-ylsulfanyl)-4-phenylsulfanylmethyl-thiazol-2-yl]-urea
- 25 1-(2-Cyclopentanecarbonyl-4-dimethylamino-phenyl)-3-[5-(1*H*-imidazol-2-ylsulfanyl)-4-phenylsulfanylmethyl-thiazol-2-yl]-urea
- 1-(2-Cyclopentanecarbonyl-4-morpholin-4-yl-phenyl)-3-[5-(1*H*-imidazol-2-ylsulfanyl)-4-phenylsulfanylmethyl-thiazol-2-yl]-urea
- 1-(3-Cyclopentanecarbonyl-pyridin-4-yl)-3-[5-(1*H*-imidazol-2-ylsulfanyl)-4-phenylsulfanylmethyl-thiazol-2-yl]-urea
- 30 1-(4-Cyclopentanecarbonyl-pyridin-3-yl)-3-[5-(1*H*-imidazol-2-ylsulfanyl)-4-phenylsulfanylmethyl-thiazol-2-yl]-urea
- 1-(3-Cyclopentanecarbonyl-pyridin-2-yl)-3-[5-(1*H*-imidazol-2-ylsulfanyl)-4-phenylsulfanylmethyl-thiazol-2-yl]-urea

- 1-(2-Cyclopentanecarbonyl-4-methyl-phenyl)-3-[5-(1*H*-imidazole-2-sulfonyl)-4-phenylsulfanylmethyl-thiazol-2-yl]-urea
- 1-(2-Cyclopentanecarbonyl-phenyl)-3-[5-(1*H*-imidazole-2-sulfonyl)-4-phenylsulfanylmethyl-thiazol-2-yl]-urea
- 5 1-(4-Chloro-2-cyclopentanecarbonyl-phenyl)-3-[5-(1*H*-imidazole-2-sulfonyl)-4-phenylsulfanylmethyl-thiazol-2-yl]-urea
- 1-(2-Cyclopentanecarbonyl-4-fluoro-phenyl)-3-[5-(1*H*-imidazole-2-sulfonyl)-4-phenylsulfanylmethyl-thiazol-2-yl]-urea
- 1-(2-Cyclopentanecarbonyl-4-trifluoromethyl-phenyl)-3-[5-(1*H*-imidazole-2-sulfonyl)-4-phenylsulfanylmethyl-thiazol-2-yl]-urea
- 10 1-(2-Cyclopentanecarbonyl-4-methoxy-phenyl)-3-[5-(1*H*-imidazole-2-sulfonyl)-4-phenylsulfanylmethyl-thiazol-2-yl]-urea
- 1-(2-Cyclopentanecarbonyl-4-dimethylamino-phenyl)-3-[5-(1*H*-imidazole-2-sulfonyl)-4-phenylsulfanylmethyl-thiazol-2-yl]-urea
- 15 1-(2-Cyclopentanecarbonyl-4-morpholin-4-yl-phenyl)-3-[5-(1*H*-imidazole-2-sulfonyl)-4-phenylsulfanylmethyl-thiazol-2-yl]-urea
- 1-(3-Cyclopentanecarbonyl-pyridin-4-yl)-3-[5-(1*H*-imidazole-2-sulfonyl)-4-phenylsulfanylmethyl-thiazol-2-yl]-urea
- 1-(4-Cyclopentanecarbonyl-pyridin-3-yl)-3-[5-(1*H*-imidazole-2-sulfonyl)-4-phenylsulfanylmethyl-thiazol-2-yl]-urea
- 20 1-(3-Cyclopentanecarbonyl-pyridin-2-yl)-3-[5-(1*H*-imidazole-2-sulfonyl)-4-phenylsulfanylmethyl-thiazol-2-yl]-urea
- 1-(2-Cyclopentanecarbonyl-4-methyl-phenyl)-3-[5-(1-methyl-1*H*-tetrazol-5-ylsulfanyl)-4-phenylsulfanylmethyl-thiazol-2-yl]-urea
- 25 1-(2-Cyclopentanecarbonyl-phenyl)-3-[5-(1-methyl-1*H*-tetrazol-5-ylsulfanyl)-4-phenylsulfanylmethyl-thiazol-2-yl]-urea
- 1-(4-Chloro-2-cyclopentanecarbonyl-phenyl)-3-[5-(1-methyl-1*H*-tetrazol-5-ylsulfanyl)-4-phenylsulfanylmethyl-thiazol-2-yl]-urea
- 1-(2-Cyclopentanecarbonyl-4-fluoro-phenyl)-3-[5-(1-methyl-1*H*-tetrazol-5-ylsulfanyl)-4-phenylsulfanylmethyl-thiazol-2-yl]-urea
- 30 1-(2-Cyclopentanecarbonyl-4-trifluoromethyl-phenyl)-3-[5-(1-methyl-1*H*-tetrazol-5-ylsulfanyl)-4-phenylsulfanylmethyl-thiazol-2-yl]-urea
- 1-(2-Cyclopentanecarbonyl-4-methoxy-phenyl)-3-[5-(1-methyl-1*H*-tetrazol-5-ylsulfanyl)-4-phenylsulfanylmethyl-thiazol-2-yl]-urea

- 1-(2-Cyclopentanecarbonyl-4-dimethylamino-phenyl)-3-[5-(1-methyl-1*H*-tetrazol-5-ylsulfanyl)-4-phenylsulfanylmethyl-thiazol-2-yl]-urea
- 1-(2-Cyclopentanecarbonyl-4-morpholin-4-yl-phenyl)-3-[5-(1-methyl-1*H*-tetrazol-5-ylsulfanyl)-4-phenylsulfanylmethyl-thiazol-2-yl]-urea
- 5 1-(3-Cyclopentanecarbonyl-pyridin-4-yl)-3-[5-(1-methyl-1*H*-tetrazol-5-ylsulfanyl)-4-phenylsulfanylmethyl-thiazol-2-yl]-urea
- 1-(4-Cyclopentanecarbonyl-pyridin-3-yl)-3-[5-(1-methyl-1*H*-tetrazol-5-ylsulfanyl)-4-phenylsulfanylmethyl-thiazol-2-yl]-urea
- 1-(3-Cyclopentanecarbonyl-pyridin-2-yl)-3-[5-(1-methyl-1*H*-tetrazol-5-ylsulfanyl)-4-phenylsulfanylmethyl-thiazol-2-yl]-urea
- 10 1-(2-Cyclopentanecarbonyl-4-methyl-phenyl)-3-[5-(1-methyl-1*H*-tetrazole-5-sulfonyl)-4-phenylsulfanylmethyl-thiazol-2-yl]-urea
- 1-(2-Cyclopentanecarbonyl-phenyl)-3-[5-(1-methyl-1*H*-tetrazole-5-sulfonyl)-4-phenylsulfanylmethyl-thiazol-2-yl]-urea
- 15 1-(4-Chloro-2-cyclopentanecarbonyl-phenyl)-3-[5-(1-methyl-1*H*-tetrazole-5-sulfonyl)-4-phenylsulfanylmethyl-thiazol-2-yl]-urea
- 1-(2-Cyclopentanecarbonyl-4-fluoro-phenyl)-3-[5-(1-methyl-1*H*-tetrazole-5-sulfonyl)-4-phenylsulfanylmethyl-thiazol-2-yl]-urea
- 1-(2-Cyclopentanecarbonyl-4-trifluoromethyl-phenyl)-3-[5-(1-methyl-1*H*-tetrazole-5-sulfonyl)-4-phenylsulfanylmethyl-thiazol-2-yl]-urea
- 20 1-(2-Cyclopentanecarbonyl-4-methoxy-phenyl)-3-[5-(1-methyl-1*H*-tetrazole-5-sulfonyl)-4-phenylsulfanylmethyl-thiazol-2-yl]-urea
- 1-(2-Cyclopentanecarbonyl-4-dimethylamino-phenyl)-3-[5-(1-methyl-1*H*-tetrazole-5-sulfonyl)-4-phenylsulfanylmethyl-thiazol-2-yl]-urea
- 25 1-(2-Cyclopentanecarbonyl-4-morpholin-4-yl-phenyl)-3-[5-(1-methyl-1*H*-tetrazole-5-sulfonyl)-4-phenylsulfanylmethyl-thiazol-2-yl]-urea
- 1-(3-Cyclopentanecarbonyl-pyridin-4-yl)-3-[5-(1-methyl-1*H*-tetrazole-5-sulfonyl)-4-phenylsulfanylmethyl-thiazol-2-yl]-urea
- 1-(4-Cyclopentanecarbonyl-pyridin-3-yl)-3-[5-(1-methyl-1*H*-tetrazole-5-sulfonyl)-4-phenylsulfanylmethyl-thiazol-2-yl]-urea
- 30 1-(3-Cyclopentanecarbonyl-pyridin-2-yl)-3-[5-(1-methyl-1*H*-tetrazole-5-sulfonyl)-4-phenylsulfanylmethyl-thiazol-2-yl]-urea
- 1-(2-Cyclopentanecarbonyl-4-methyl-phenyl)-3-[4-phenylsulfanylmethyl-5-(1*H*-tetrazole-5-sulfonyl)-thiazol-2-yl]-urea

- 1-(2-Cyclopentanecarbonyl-phenyl)-3-[4-phenylsulfanylmethyl-5-(1*H*-tetrazole-5-sulfonyl)-thiazol-2-yl]-urea
- 1-(4-Chloro-2-cyclopentanecarbonyl-phenyl)-3-[4-phenylsulfanylmethyl-5-(1*H*-tetrazole-5-sulfonyl)-thiazol-2-yl]-urea
- 5 1-(2-Cyclopentanecarbonyl-4-fluoro-phenyl)-3-[4-phenylsulfanylmethyl-5-(1*H*-tetrazole-5-sulfonyl)-thiazol-2-yl]-urea
- 1-(2-Cyclopentanecarbonyl-4-trifluoromethyl-phenyl)-3-[4-phenylsulfanylmethyl-5-(1*H*-tetrazole-5-sulfonyl)-thiazol-2-yl]-urea
- 1-(2-Cyclopentanecarbonyl-4-methoxy-phenyl)-3-[4-phenylsulfanylmethyl-5-(1*H*-tetrazole-5-sulfonyl)-thiazol-2-yl]-urea
- 10 1-(2-Cyclopentanecarbonyl-4-dimethylamino-phenyl)-3-[4-phenylsulfanylmethyl-5-(1*H*-tetrazole-5-sulfonyl)-thiazol-2-yl]-urea
- 1-(2-Cyclopentanecarbonyl-4-morpholin-4-yl-phenyl)-3-[4-phenylsulfanylmethyl-5-(1*H*-tetrazole-5-sulfonyl)-thiazol-2-yl]-urea
- 15 1-(3-Cyclopentanecarbonyl-pyridin-4-yl)-3-[4-phenylsulfanylmethyl-5-(1*H*-tetrazole-5-sulfonyl)-thiazol-2-yl]-urea
- 1-(4-Cyclopentanecarbonyl-pyridin-3-yl)-3-[4-phenylsulfanylmethyl-5-(1*H*-tetrazole-5-sulfonyl)-thiazol-2-yl]-urea
- 1-(3-Cyclopentanecarbonyl-pyridin-2-yl)-3-[4-phenylsulfanylmethyl-5-(1*H*-tetrazole-5-sulfonyl)-thiazol-2-yl]-urea
- 20 *N*-(5-Chloro-2-[3-(2-cyclopentanecarbonyl-4-methyl-phenyl)-ureido]-thiazol-4-ylmethyl)-methanesulfonamide
- N*-(5-Chloro-2-[3-(2-cyclopentanecarbonyl-phenyl)-ureido]-thiazol-4-ylmethyl)-methanesulfonamide
- 25 *N*-(5-Chloro-2-[3-(4-chloro-2-cyclopentanecarbonyl-phenyl)-ureido]-thiazol-4-ylmethyl)-methanesulfonamide
- N*-(5-Chloro-2-[3-(2-cyclopentanecarbonyl-4-fluoro-phenyl)-ureido]-thiazol-4-ylmethyl)-methanesulfonamide
- N*-(5-Chloro-2-[3-(2-cyclopentanecarbonyl-4-trifluoromethyl-phenyl)-ureido]-thiazol-4-ylmethyl)-methanesulfonamide
- 30 *N*-(5-Chloro-2-[3-(2-cyclopentanecarbonyl-4-methoxy-phenyl)-ureido]-thiazol-4-ylmethyl)-methanesulfonamide
- N*-(5-Chloro-2-[3-(2-cyclopentanecarbonyl-4-dimethylamino-phenyl)-ureido]-thiazol-4-ylmethyl)-methanesulfonamide

- N*-(5-Chloro-2-[3-(2-cyclopentanecarbonyl-4-morpholin-4-yl-phenyl)-ureido]-thiazol-4-ylmethyl)-methanesulfonamide
- N*-(5-Chloro-2-[3-(3-cyclopentanecarbonyl-pyridin-4-yl)-ureido]-thiazol-4-ylmethyl)-methanesulfonamide
- 5 *N*-(5-Chloro-2-[3-(4-cyclopentanecarbonyl-pyridin-3-yl)-ureido]-thiazol-4-ylmethyl)-methanesulfonamide
- N*-(5-Chloro-2-[3-(3-cyclopentanecarbonyl-pyridin-2-yl)-ureido]-thiazol-4-ylmethyl)-methanesulfonamide
- N*-(5-Bromo-2-[3-(2-cyclopentanecarbonyl-4-methyl-phenyl)-ureido]-thiazol-4-ylmethyl)-methanesulfonamide
- 10 *N*-(5-Bromo-2-[3-(2-cyclopentanecarbonyl-phenyl)-ureido]-thiazol-4-ylmethyl)-methanesulfonamide
- N*-(5-Bromo-2-[3-(4-chloro-2-cyclopentanecarbonyl-phenyl)-ureido]-thiazol-4-ylmethyl)-methanesulfonamide
- 15 *N*-(5-Bromo-2-[3-(2-cyclopentanecarbonyl-4-fluoro-phenyl)-ureido]-thiazol-4-ylmethyl)-methanesulfonamide
- N*-(5-Bromo-2-[3-(2-cyclopentanecarbonyl-4-trifluoromethyl-phenyl)-ureido]-thiazol-4-ylmethyl)-methanesulfonamide
- N*-(5-Bromo-2-[3-(2-cyclopentanecarbonyl-4-methoxy-phenyl)-ureido]-thiazol-4-ylmethyl)-methanesulfonamide
- 20 *N*-(5-Bromo-2-[3-(2-cyclopentanecarbonyl-4-dimethylamino-phenyl)-ureido]-thiazol-4-ylmethyl)-methanesulfonamide
- N*-(5-Bromo-2-[3-(2-cyclopentanecarbonyl-4-morpholin-4-yl-phenyl)-ureido]-thiazol-4-ylmethyl)-methanesulfonamide
- 25 *N*-(5-Bromo-2-[3-(3-cyclopentanecarbonyl-pyridin-4-yl)-ureido]-thiazol-4-ylmethyl)-methanesulfonamide
- N*-(5-Bromo-2-[3-(4-cyclopentanecarbonyl-pyridin-3-yl)-ureido]-thiazol-4-ylmethyl)-methanesulfonamide
- N*-(5-Bromo-2-[3-(3-cyclopentanecarbonyl-pyridin-2-yl)-ureido]-thiazol-4-ylmethyl)-methanesulfonamide
- 30 *N*-(2-[3-(2-Cyclopentanecarbonyl-4-methyl-phenyl)-ureido]-5-methylsulfanyl-thiazol-4-ylmethyl)-methanesulfonamide
- N*-(2-[3-(2-Cyclopentanecarbonyl-phenyl)-ureido]-5-methylsulfanyl-thiazol-4-ylmethyl)-methanesulfonamide

- N*-{2-[3-(4-Chloro-2-cyclopentanecarbonyl-phenyl)-ureido]-5-methylsulfanyl-thiazol-4-ylmethyl}-methanesulfonamide
- N*-{2-[3-(2-Cyclopentanecarbonyl-4-fluoro-phenyl)-ureido]-5-methylsulfanyl-thiazol-4-ylmethyl}-methanesulfonamide
- 5 *N*-{2-[3-(2-Cyclopentanecarbonyl-4-trifluoromethyl-phenyl)-ureido]-5-methylsulfanyl-thiazol-4-ylmethyl}-methanesulfonamide
- N*-{2-[3-(2-Cyclopentanecarbonyl-4-methoxy-phenyl)-ureido]-5-methylsulfanyl-thiazol-4-ylmethyl}-methanesulfonamide
- N*-{2-[3-(2-Cyclopentanecarbonyl-4-dimethylamino-phenyl)-ureido]-5-methylsulfanyl-thiazol-10 4-ylmethyl}-methanesulfonamide
- N*-{2-[3-(2-Cyclopentanecarbonyl-4-morpholin-4-yl-phenyl)-ureido]-5-methylsulfanyl-thiazol-4-ylmethyl}-methanesulfonamide
- N*-{2-[3-(3-Cyclopentanecarbonyl-pyridin-4-yl)-ureido]-5-methylsulfanyl-thiazol-4-ylmethyl}-methanesulfonamide
- 15 *N*-{2-[3-(4-Cyclopentanecarbonyl-pyridin-3-yl)-ureido]-5-methylsulfanyl-thiazol-4-ylmethyl}-methanesulfonamide
- N*-{2-[3-(3-Cyclopentanecarbonyl-pyridin-2-yl)-ureido]-5-methylsulfanyl-thiazol-4-ylmethyl}-methanesulfonamide
- N*-{2-[3-(2-Cyclopentanecarbonyl-4-methyl-phenyl)-ureido]-5-methanesulfonyl-thiazol-4-ylmethyl}-methanesulfonamide
- 20 *N*-{2-[3-(2-Cyclopentanecarbonyl-phenyl)-ureido]-5-methanesulfonyl-thiazol-4-ylmethyl}-methanesulfonamide
- N*-{2-[3-(4-Chloro-2-cyclopentanecarbonyl-phenyl)-ureido]-5-methanesulfonyl-thiazol-4-ylmethyl}-methanesulfonamide
- 25 *N*-{2-[3-(2-Cyclopentanecarbonyl-4-fluoro-phenyl)-ureido]-5-methanesulfonyl-thiazol-4-ylmethyl}-methanesulfonamide
- N*-{2-[3-(2-Cyclopentanecarbonyl-4-trifluoromethyl-phenyl)-ureido]-5-methanesulfonyl-thiazol-4-ylmethyl}-methanesulfonamide
- N*-{2-[3-(2-Cyclopentanecarbonyl-4-methoxy-phenyl)-ureido]-5-methanesulfonyl-thiazol-4-ylmethyl}-methanesulfonamide
- 30 *N*-{2-[3-(2-Cyclopentanecarbonyl-4-dimethylamino-phenyl)-ureido]-5-methanesulfonyl-thiazol-4-ylmethyl}-methanesulfonamide
- N*-{2-[3-(2-Cyclopentanecarbonyl-4-morpholin-4-yl-phenyl)-ureido]-5-methanesulfonyl-thiazol-4-ylmethyl}-methanesulfonamide

- N*-[2-[3-(3-Cyclopentanecarbonyl-pyridin-4-yl)-ureido]-5-methanesulfonyl-thiazol-4-ylmethyl]-methanesulfonamide
- N*-[2-[3-(4-Cyclopentanecarbonyl-pyridin-3-yl)-ureido]-5-methanesulfonyl-thiazol-4-ylmethyl]-methanesulfonamide
- 5 *N*-[2-[3-(3-Cyclopentanecarbonyl-pyridin-2-yl)-ureido]-5-methanesulfonyl-thiazol-4-ylmethyl]-methanesulfonamide
- N*-[2-[3-(2-Cyclopentanecarbonyl-4-methyl-phenyl)-ureido]-5-(pyridin-2-ylsulfanyl)-thiazol-4-ylmethyl]-methanesulfonamide
- N*-[2-[3-(2-Cyclopentanecarbonyl-phenyl)-ureido]-5-(pyridin-2-ylsulfanyl)-thiazol-4-ylmethyl]-methanesulfonamide
- 10 *N*-[2-[3-(4-Chloro-2-cyclopentanecarbonyl-phenyl)-ureido]-5-(pyridin-2-ylsulfanyl)-thiazol-4-ylmethyl]-methanesulfonamide
- N*-[2-[3-(2-Cyclopentanecarbonyl-4-fluoro-phenyl)-ureido]-5-(pyridin-2-ylsulfanyl)-thiazol-4-ylmethyl]-methanesulfonamide
- 15 *N*-[2-[3-(2-Cyclopentanecarbonyl-4-trifluoromethyl-phenyl)-ureido]-5-(pyridin-2-ylsulfanyl)-thiazol-4-ylmethyl]-methanesulfonamide
- N*-[2-[3-(2-Cyclopentanecarbonyl-4-methoxy-phenyl)-ureido]-5-(pyridin-2-ylsulfanyl)-thiazol-4-ylmethyl]-methanesulfonamide
- N*-[2-[3-(2-Cyclopentanecarbonyl-4-dimethylamino-phenyl)-ureido]-5-(pyridin-2-ylsulfanyl)-thiazol-4-ylmethyl]-methanesulfonamide
- 20 *N*-[2-[3-(2-Cyclopentanecarbonyl-4-morpholin-4-yl-phenyl)-ureido]-5-(pyridin-2-ylsulfanyl)-thiazol-4-ylmethyl]-methanesulfonamide
- N*-[2-[3-(3-Cyclopentanecarbonyl-pyridin-4-yl)-ureido]-5-(pyridin-2-ylsulfanyl)-thiazol-4-ylmethyl]-methanesulfonamide
- 25 *N*-[2-[3-(4-Cyclopentanecarbonyl-pyridin-3-yl)-ureido]-5-(pyridin-2-ylsulfanyl)-thiazol-4-ylmethyl]-methanesulfonamide
- N*-[2-[3-(3-Cyclopentanecarbonyl-pyridin-2-yl)-ureido]-5-(pyridin-2-ylsulfanyl)-thiazol-4-ylmethyl]-methanesulfonamide
- N*-[2-[3-(2-Cyclopentanecarbonyl-4-methyl-phenyl)-ureido]-5-(pyridine-2-sulfonyl)-thiazol-4-ylmethyl]-methanesulfonamide
- 30 *N*-[2-[3-(2-Cyclopentanecarbonyl-phenyl)-ureido]-5-(pyridine-2-sulfonyl)-thiazol-4-ylmethyl]-methanesulfonamide
- N*-[2-[3-(4-Chloro-2-cyclopentanecarbonyl-phenyl)-ureido]-5-(pyridine-2-sulfonyl)-thiazol-4-ylmethyl]-methanesulfonamide

- N*-[2-[3-(2-Cyclopentanecarbonyl-4-fluoro-phenyl)-ureido]-5-(pyridine-2-sulfonyl)-thiazol-4-ylmethyl]-methanesulfonamide
- N*-[2-[3-(2-Cyclopentanecarbonyl-4-trifluoromethyl-phenyl)-ureido]-5-(pyridine-2-sulfonyl)-thiazol-4-ylmethyl]-methanesulfonamide
- 5 *N*-[2-[3-(2-Cyclopentanecarbonyl-4-methoxy-phenyl)-ureido]-5-(pyridine-2-sulfonyl)-thiazol-4-ylmethyl]-methanesulfonamide
- N*-[2-[3-(2-Cyclopentanecarbonyl-4-dimethylamino-phenyl)-ureido]-5-(pyridine-2-sulfonyl)-thiazol-4-ylmethyl]-methanesulfonamide
- N*-[2-[3-(2-Cyclopentanecarbonyl-4-morpholin-4-yl-phenyl)-ureido]-5-(pyridine-2-sulfonyl)-thiazol-4-ylmethyl]-methanesulfonamide
- 10 *N*-[2-[3-(3-Cyclopentanecarbonyl-pyridin-4-yl)-ureido]-5-(pyridine-2-sulfonyl)-thiazol-4-ylmethyl]-methanesulfonamide
- N*-[2-[3-(4-Cyclopentanecarbonyl-pyridin-3-yl)-ureido]-5-(pyridine-2-sulfonyl)-thiazol-4-ylmethyl]-methanesulfonamide
- 15 *N*-[2-[3-(3-Cyclopentanecarbonyl-pyridin-2-yl)-ureido]-5-(pyridine-2-sulfonyl)-thiazol-4-ylmethyl]-methanesulfonamide
- N*-[2-[3-(2-Cyclopentanecarbonyl-4-methyl-phenyl)-ureido]-5-(1*H*-imidazol-2-ylsulfanyl)-thiazol-4-ylmethyl]-methanesulfonamide
- N*-[2-[3-(2-Cyclopentanecarbonyl-phenyl)-ureido]-5-(1*H*-imidazol-2-ylsulfanyl)-thiazol-4-ylmethyl]-methanesulfonamide
- 20 *N*-[2-[3-(4-Chloro-2-cyclopentanecarbonyl-phenyl)-ureido]-5-(1*H*-imidazol-2-ylsulfanyl)-thiazol-4-ylmethyl]-methanesulfonamide
- N*-[2-[3-(2-Cyclopentanecarbonyl-4-fluoro-phenyl)-ureido]-5-(1*H*-imidazol-2-ylsulfanyl)-thiazol-4-ylmethyl]-methanesulfonamide
- 25 *N*-[2-[3-(2-Cyclopentanecarbonyl-4-trifluoromethyl-phenyl)-ureido]-5-(1*H*-imidazol-2-ylsulfanyl)-thiazol-4-ylmethyl]-methanesulfonamide
- N*-[2-[3-(2-Cyclopentanecarbonyl-4-methoxy-phenyl)-ureido]-5-(1*H*-imidazol-2-ylsulfanyl)-thiazol-4-ylmethyl]-methanesulfonamide
- N*-[2-[3-(2-Cyclopentanecarbonyl-4-dimethylamino-phenyl)-ureido]-5-(1*H*-imidazol-2-ylsulfanyl)-thiazol-4-ylmethyl]-methanesulfonamide
- 30 *N*-[2-[3-(2-Cyclopentanecarbonyl-4-morpholin-4-yl-phenyl)-ureido]-5-(1*H*-imidazol-2-ylsulfanyl)-thiazol-4-ylmethyl]-methanesulfonamide
- N*-[2-[3-(3-Cyclopentanecarbonyl-pyridin-4-yl)-ureido]-5-(1*H*-imidazol-2-ylsulfanyl)-thiazol-4-ylmethyl]-methanesulfonamide

- N*-[2-[3-(4-Cyclopentanecarbonyl-pyridin-3-yl)-ureido]-5-(1*H*-imidazol-2-ylsulfanyl)-thiazol-4-ylmethyl]-methanesulfonamide
- N*-[2-[3-(3-Cyclopentanecarbonyl-pyridin-2-yl)-ureido]-5-(1*H*-imidazol-2-ylsulfanyl)-thiazol-4-ylmethyl]-methanesulfonamide
- 5 *N*-[2-[3-(2-Cyclopentanecarbonyl-4-methyl-phenyl)-ureido]-5-(1*H*-imidazole-2-sulfonyl)-thiazol-4-ylmethyl]-methanesulfonamide
- N*-[2-[3-(2-Cyclopentanecarbonyl-phenyl)-ureido]-5-(1*H*-imidazole-2-sulfonyl)-thiazol-4-ylmethyl]-methanesulfonamide
- 10 *N*-[2-[3-(4-Chloro-2-cyclopentanecarbonyl-phenyl)-ureido]-5-(1*H*-imidazole-2-sulfonyl)-thiazol-4-ylmethyl]-methanesulfonamide
- N*-[2-[3-(2-Cyclopentanecarbonyl-4-fluoro-phenyl)-ureido]-5-(1*H*-imidazole-2-sulfonyl)-thiazol-4-ylmethyl]-methanesulfonamide
- N*-[2-[3-(2-Cyclopentanecarbonyl-4-trifluoromethyl-phenyl)-ureido]-5-(1*H*-imidazole-2-sulfonyl)-thiazol-4-ylmethyl]-methanesulfonamide
- 15 *N*-[2-[3-(2-Cyclopentanecarbonyl-4-methoxy-phenyl)-ureido]-5-(1*H*-imidazole-2-sulfonyl)-thiazol-4-ylmethyl]-methanesulfonamide
- N*-[2-[3-(2-Cyclopentanecarbonyl-4-dimethylamino-phenyl)-ureido]-5-(1*H*-imidazole-2-sulfonyl)-thiazol-4-ylmethyl]-methanesulfonamide
- N*-[2-[3-(2-Cyclopentanecarbonyl-4-morpholin-4-yl-phenyl)-ureido]-5-(1*H*-imidazole-2-sulfonyl)-thiazol-4-ylmethyl]-methanesulfonamide
- 20 *N*-[2-[3-(3-Cyclopentanecarbonyl-pyridin-4-yl)-ureido]-5-(1*H*-imidazole-2-sulfonyl)-thiazol-4-ylmethyl]-methanesulfonamide
- N*-[2-[3-(4-Cyclopentanecarbonyl-pyridin-3-yl)-ureido]-5-(1*H*-imidazole-2-sulfonyl)-thiazol-4-ylmethyl]-methanesulfonamide
- 25 *N*-[2-[3-(3-Cyclopentanecarbonyl-pyridin-2-yl)-ureido]-5-(1*H*-imidazole-2-sulfonyl)-thiazol-4-ylmethyl]-methanesulfonamide
- N*-[2-[3-(2-Cyclopentanecarbonyl-4-methyl-phenyl)-ureido]-5-(1-methyl-1*H*-tetrazol-5-ylsulfanyl)-thiazol-4-ylmethyl]-methanesulfonamide
- N*-[2-[3-(2-Cyclopentanecarbonyl-phenyl)-ureido]-5-(1-methyl-1*H*-tetrazol-5-ylsulfanyl)-thiazol-4-ylmethyl]-methanesulfonamide
- 30 *N*-[2-[3-(4-Chloro-2-cyclopentanecarbonyl-phenyl)-ureido]-5-(1-methyl-1*H*-tetrazol-5-ylsulfanyl)-thiazol-4-ylmethyl]-methanesulfonamide
- N*-[2-[3-(2-Cyclopentanecarbonyl-4-fluoro-phenyl)-ureido]-5-(1-methyl-1*H*-tetrazol-5-ylsulfanyl)-thiazol-4-ylmethyl]-methanesulfonamide

- N*-[2-[3-(2-Cyclopentanecarbonyl-4-trifluoromethyl-phenyl)-ureido]-5-(1-methyl-1*H*-tetrazol-5-ylsulfanyl)-thiazol-4-ylmethyl]-methanesulfonamide
- N*-[2-[3-(2-Cyclopentanecarbonyl-4-methoxy-phenyl)-ureido]-5-(1-methyl-1*H*-tetrazol-5-ylsulfanyl)-thiazol-4-ylmethyl]-methanesulfonamide
- 5 *N*-[2-[3-(2-Cyclopentanecarbonyl-4-dimethylamino-phenyl)-ureido]-5-(1-methyl-1*H*-tetrazol-5-ylsulfanyl)-thiazol-4-ylmethyl]-methanesulfonamide
- N*-[2-[3-(2-Cyclopentanecarbonyl-4-morpholin-4-yl-phenyl)-ureido]-5-(1-methyl-1*H*-tetrazol-5-ylsulfanyl)-thiazol-4-ylmethyl]-methanesulfonamide
- 10 *N*-[2-[3-(3-Cyclopentanecarbonyl-pyridin-4-yl)-ureido]-5-(1-methyl-1*H*-tetrazol-5-ylsulfanyl)-thiazol-4-ylmethyl]-methanesulfonamide
- N*-[2-[3-(4-Cyclopentanecarbonyl-pyridin-3-yl)-ureido]-5-(1-methyl-1*H*-tetrazol-5-ylsulfanyl)-thiazol-4-ylmethyl]-methanesulfonamide
- N*-[2-[3-(3-Cyclopentanecarbonyl-pyridin-2-yl)-ureido]-5-(1-methyl-1*H*-tetrazol-5-ylsulfanyl)-thiazol-4-ylmethyl]-methanesulfonamide
- 15 *N*-[2-[3-(2-Cyclopentanecarbonyl-4-methyl-phenyl)-ureido]-5-(1-methyl-1*H*-tetrazole-5-sulfonyl)-thiazol-4-ylmethyl]-methanesulfonamide
- N*-[2-[3-(2-Cyclopentanecarbonyl-phenyl)-ureido]-5-(1-methyl-1*H*-tetrazole-5-sulfonyl)-thiazol-4-ylmethyl]-methanesulfonamide
- 20 *N*-[2-[3-(4-Chloro-2-cyclopentanecarbonyl-phenyl)-ureido]-5-(1-methyl-1*H*-tetrazole-5-sulfonyl)-thiazol-4-ylmethyl]-methanesulfonamide
- N*-[2-[3-(2-Cyclopentanecarbonyl-4-fluoro-phenyl)-ureido]-5-(1-methyl-1*H*-tetrazole-5-sulfonyl)-thiazol-4-ylmethyl]-methanesulfonamide
- N*-[2-[3-(2-Cyclopentanecarbonyl-4-trifluoromethyl-phenyl)-ureido]-5-(1-methyl-1*H*-tetrazole-5-sulfonyl)-thiazol-4-ylmethyl]-methanesulfonamide
- 25 *N*-[2-[3-(2-Cyclopentanecarbonyl-4-methoxy-phenyl)-ureido]-5-(1-methyl-1*H*-tetrazole-5-sulfonyl)-thiazol-4-ylmethyl]-methanesulfonamide
- N*-[2-[3-(2-Cyclopentanecarbonyl-4-dimethylamino-phenyl)-ureido]-5-(1-methyl-1*H*-tetrazole-5-sulfonyl)-thiazol-4-ylmethyl]-methanesulfonamide
- N*-[2-[3-(2-Cyclopentanecarbonyl-4-morpholin-4-yl-phenyl)-ureido]-5-(1-methyl-1*H*-tetrazole-5-sulfonyl)-thiazol-4-ylmethyl]-methanesulfonamide
- 30 *N*-[2-[3-(3-Cyclopentanecarbonyl-pyridin-4-yl)-ureido]-5-(1-methyl-1*H*-tetrazole-5-sulfonyl)-thiazol-4-ylmethyl]-methanesulfonamide
- N*-[2-[3-(4-Cyclopentanecarbonyl-pyridin-3-yl)-ureido]-5-(1-methyl-1*H*-tetrazole-5-sulfonyl)-thiazol-4-ylmethyl]-methanesulfonamide

- N*-[2-[3-(3-Cyclopentanecarbonyl-pyridin-2-yl)-ureido]-5-(1-methyl-1*H*-tetrazole-5-sulfonyl)-thiazol-4-ylmethyl]-methanesulfonamide
- N*-[2-[3-(2-Cyclopentanecarbonyl-4-methyl-phenyl)-ureido]-5-(1*H*-tetrazole-5-sulfonyl)-thiazol-4-ylmethyl]-methanesulfonamide
- 5 *N*-[2-[3-(2-Cyclopentanecarbonyl-phenyl)-ureido]-5-(1*H*-tetrazole-5-sulfonyl)-thiazol-4-ylmethyl]-methanesulfonamide
- N*-[2-[3-(4-Chloro-2-cyclopentanecarbonyl-phenyl)-ureido]-5-(1*H*-tetrazole-5-sulfonyl)-thiazol-4-ylmethyl]-methanesulfonamide
- N*-[2-[3-(2-Cyclopentanecarbonyl-4-fluoro-phenyl)-ureido]-5-(1*H*-tetrazole-5-sulfonyl)-thiazol-4-ylmethyl]-methanesulfonamide
- 10 *N*-[2-[3-(2-Cyclopentanecarbonyl-4-trifluoromethyl-phenyl)-ureido]-5-(1*H*-tetrazole-5-sulfonyl)-thiazol-4-ylmethyl]-methanesulfonamide
- N*-[2-[3-(2-Cyclopentanecarbonyl-4-methoxy-phenyl)-ureido]-5-(1*H*-tetrazole-5-sulfonyl)-thiazol-4-ylmethyl]-methanesulfonamide
- 15 *N*-[2-[3-(2-Cyclopentanecarbonyl-4-dimethylamino-phenyl)-ureido]-5-(1*H*-tetrazole-5-sulfonyl)-thiazol-4-ylmethyl]-methanesulfonamide
- N*-[2-[3-(2-Cyclopentanecarbonyl-4-morpholin-4-yl-phenyl)-ureido]-5-(1*H*-tetrazole-5-sulfonyl)-thiazol-4-ylmethyl]-methanesulfonamide
- N*-[2-[3-(3-Cyclopentanecarbonyl-pyridin-4-yl)-ureido]-5-(1*H*-tetrazole-5-sulfonyl)-thiazol-4-ylmethyl]-methanesulfonamide
- 20 *N*-[2-[3-(4-Cyclopentanecarbonyl-pyridin-3-yl)-ureido]-5-(1*H*-tetrazole-5-sulfonyl)-thiazol-4-ylmethyl]-methanesulfonamide
- N*-[2-[3-(3-Cyclopentanecarbonyl-pyridin-2-yl)-ureido]-5-(1*H*-tetrazole-5-sulfonyl)-thiazol-4-ylmethyl]-methanesulfonamide
- 25 *N*-(2-cyclopentanecarbonyl-phenyl)-*N'*-thiazol-2-yl-oxalamide
- N*-(5-bromo-thiazol-2-yl)-*N'*-(2-cyclopentanecarbonyl-phenyl)-oxalamide
- N*-(5-chloro-thiazol-2-yl)-*N'*-(2-cyclopentanecarbonyl-phenyl)-oxalamide
- 4;*N*-(2-cyclopentanecarbonyl-phenyl)-*N'*-(5-methylsulfanyl-thiazol-2-yl)-oxalamide
- 30 *N*-(2-cyclopentanecarbonyl-phenyl)-*N'*-(5-methanesulfonyl-thiazol-2-yl)-oxalamide
- N*-(2-cyclopentanecarbonyl-phenyl)-*N'*-[5-(pyridin-2-ylsulfanyl)-thiazol-2-yl]-oxalamide
- N*-(2-cyclopentanecarbonyl-phenyl)-*N'*-[5-(1*H*-imidazol-2-ylsulfanyl)-thiazol-2-yl]-oxalamide
- {5-bromo-2-[(2-cyclopentanecarbonyl-phenylaminooxalyl)-amino]-thiazol-4-yl}-acetic acid
- {5-chloro-2-[(2-cyclopentanecarbonyl-phenylaminooxalyl)-amino]-thiazol-4-yl}-acetic acid

- {2-[(2-cyclopentanecarbonyl-phenylaminooxaly)-amino]-5-methylsulfanyl-thiazol-4-yl}-acetic acid
- {2-[(2-cyclopentanecarbonyl-phenylaminooxaly)-amino]-5-methanesulfonyl-thiazol-4-yl}-acetic acid
- 5 [2-[(2-cyclopentanecarbonyl-phenylaminooxaly)-amino]-5-(pyridin-2-ylsulfanyl)-thiazol-4-yl]-acetic acid
- [2-[(2-cyclopentanecarbonyl-phenylaminooxaly)-amino]-5-(1H-imidazol-2-ylsulfanyl)-thiazol-4-yl]-acetic acid
- N-(2-cyclopentanecarbonyl-phenyl)-N'-[4-(methanesulfonylamino-methyl)-thiazol-2-yl]-oxalamide
- 10 N-[5-bromo-4-(methanesulfonylamino-methyl)-thiazol-2-yl]-N'-(2-cyclopentanecarbonyl-phenyl)-oxalamide
- N-[5-chloro-4-(methanesulfonylamino-methyl)-thiazol-2-yl]-N'-(2-cyclopentanecarbonyl-phenyl)-oxalamide
- 15 N-(2-cyclopentanecarbonyl-phenyl)-N'-[4-(methanesulfonylamino-methyl)-5-methylsulfanyl-thiazol-2-yl]-oxalamide
- N-(2-cyclopentanecarbonyl-phenyl)-N'-[5-methanesulfonyl-4-(methanesulfonylamino-methyl)-thiazol-2-yl]-oxalamide
- N-(2-cyclopentanecarbonyl-phenyl)-N'-[4-(methanesulfonylamino-methyl)-5-(pyridin-2-ylsulfanyl)-thiazol-2-yl]-oxalamide
- 20 N-(2-cyclopentanecarbonyl-phenyl)-N'-[5-(1H-imidazol-2-ylsulfanyl)-4-(methanesulfonylamino-methyl)-thiazol-2-yl]-oxalamide
- N-(2-cyclopentanecarbonyl-phenyl)-N'-thiazol-2-yl-malonamide
- N-(5-bromo-thiazol-2-yl)-N'-(2-cyclopentanecarbonyl-phenyl)-malonamide
- 25 N-(5-chloro-thiazol-2-yl)-N'-(2-cyclopentanecarbonyl-phenyl)-malonamide
- N-(2-cyclopentanecarbonyl-phenyl)-N'-(5-methylsulfanyl-thiazol-2-yl)-malonamide
- N-(2-cyclopentanecarbonyl-phenyl)-N'-(5-methanesulfonyl-thiazol-2-yl)-malonamide
- N-(2-cyclopentanecarbonyl-phenyl)-N'-[5-(pyridin-2-ylsulfanyl)-thiazol-2-yl]-malonamide
- N-(2-cyclopentanecarbonyl-phenyl)-N'-[5-(1H-imidazol-2-ylsulfanyl)-thiazol-2-yl]-malonamide
- 30 {2-[2-(2-cyclopentanecarbonyl-phenylcarbamoyl)-acetylamino]-thiazol-4-yl}-acetic acid
- {5-bromo-2-[2-(2-cyclopentanecarbonyl-phenylcarbamoyl)-acetylamino]-thiazol-4-yl}-acetic acid
- {5-chloro-2-[2-(2-cyclopentanecarbonyl-phenylcarbamoyl)-acetylamino]-thiazol-4-yl}-acetic acid

- {2-[2-(2-cyclopentanecarbonyl-phenylcarbamoyl)-acetyl-amino]-5-methylsulfanyl-thiazol-4-yl}-acetic acid
- {2-[2-(2-cyclopentanecarbonyl-phenylcarbamoyl)-acetyl-amino]-5-methanesulfonyl-thiazol-4-yl}-acetic acid
- 5 {2-[2-(2-cyclopentanecarbonyl-phenylcarbamoyl)-acetyl-amino]-5-(pyridin-2-ylsulfanyl)-thiazol-4-yl}-acetic acid
- {2-[2-(2-cyclopentanecarbonyl-phenylcarbamoyl)-acetyl-amino]-5-(1H-imidazol-2-ylsulfanyl)-thiazol-4-yl}-acetic acid
- N-(2-cyclopentanecarbonyl-phenyl)-N'-[4-(methanesulfonylamino-methyl)-thiazol-2-yl]-malonamide
- 10 N-[5-bromo-4-(methanesulfonylamino-methyl)-thiazol-2-yl]-N'-(2-cyclopentanecarbonyl-phenyl)-malonamide
- N-[5-chloro-4-(methanesulfonylamino-methyl)-thiazol-2-yl]-N'-(2-cyclopentanecarbonyl-phenyl)-malonamide
- 15 N-(2-cyclopentanecarbonyl-phenyl)-N'-[4-(methanesulfonylamino-methyl)-5-methylsulfanyl-thiazol-2-yl]-malonamide
- N-(2-cyclopentanecarbonyl-phenyl)-N'-[5-methanesulfonyl-4-(methanesulfonylamino-methyl)-thiazol-2-yl]-malonamide
- N-(2-cyclopentanecarbonyl-phenyl)-N'-[4-(methanesulfonylamino-methyl)-5-(pyridin-2-ylsulfanyl)-thiazol-2-yl]-malonamide
- 20 N-(2-cyclopentanecarbonyl-phenyl)-N'-[5-(1H-imidazol-2-ylsulfanyl)-4-(methanesulfonylamino-methyl)-thiazol-2-yl]-malonamide
- N-(2-isobutoxy-phenyl)-N'-thiazol-2-yl-oxalamide
- N-(5-bromo-thiazol-2-yl)-N'-(2-isobutoxy-phenyl)-oxalamide
- 25 N-(5-chloro-thiazol-2-yl)-N'-(2-isobutoxy-phenyl)-oxalamide
- N-(2-isobutoxy-phenyl)-N'-(5-methylsulfanyl-thiazol-2-yl)-oxalamide
- N-(2-isobutoxy-phenyl)-N'-(5-methanesulfonyl-thiazol-2-yl)-oxalamide
- N-(2-isobutoxy-phenyl)-N'-[5-(pyridin-2-ylsulfanyl)-thiazol-2-yl]-oxalamide
- N-[5-(1H-imidazol-2-ylsulfanyl)-thiazol-2-yl]-N'-(2-isobutoxy-phenyl)-oxalamide
- 30 {2-[(2-isobutoxy-phenylamino)oxalyl]-amino}-thiazol-4-yl}-acetic acid
- {5-bromo-2-[(2-isobutoxy-phenylamino)oxalyl]-amino}-thiazol-4-yl}-acetic acid
- {5-chloro-2-[(2-isobutoxy-phenylamino)oxalyl]-amino}-thiazol-4-yl}-acetic acid
- {2-[(2-isobutoxy-phenylamino)oxalyl]-amino}-5-methylsulfanyl-thiazol-4-yl}-acetic acid
- {2-[(2-isobutoxy-phenylamino)oxalyl]-amino}-5-methanesulfonyl-thiazol-4-yl}-acetic acid
- 35 {2-[(2-isobutoxy-phenylamino)oxalyl]-amino}-5-(pyridin-2-ylsulfanyl)-thiazol-4-yl}-acetic acid

- {5-(1H-imidazol-2-ylsulfanyl)-2-[(2-isobutoxy-phenylaminooxalyl)-amino]-thiazol-4-yl}-acetic acid
- N-(2-isobutoxy-phenyl)-N'-[4-(methanesulfonylamino-methyl)-thiazol-2-yl]-oxalamide
- N-[5-bromo-4-(methanesulfonylamino-methyl)-thiazol-2-yl]-N'-(2-isobutoxy-phenyl)-oxalamide
- 5 N-[5-chloro-4-(methanesulfonylamino-methyl)-thiazol-2-yl]-N'-(2-isobutoxy-phenyl)-oxalamide
- N-(2-isobutoxy-phenyl)-N'-[4-(methanesulfonylamino-methyl)-5-methylsulfanyl-thiazol-2-yl]-oxalamide
- N-(2-isobutoxy-phenyl)-N'-[5-methanesulfonyl-4-(methanesulfonylamino-methyl)-thiazol-2-yl]-oxalamide
- 10 N-(2-isobutoxy-phenyl)-N'-[4-(methanesulfonylamino-methyl)-5-(pyridin-2-ylsulfanyl)-thiazol-2-yl]-oxalamide
- N-[5-(1H-imidazol-2-ylsulfanyl)-4-(methanesulfonylamino-methyl)-thiazol-2-yl]-N'-(2-isobutoxy-phenyl)-oxalamide
- N-(2-isobutoxy-phenyl)-N'-thiazol-2-yl-malonamide
- 15 N-(5-bromo-thiazol-2-yl)-N'-(2-isobutoxy-phenyl)-malonamide
- N-(5-chloro-thiazol-2-yl)-N'-(2-isobutoxy-phenyl)-malonamide
- N-(2-isobutoxy-phenyl)-N'-(5-methylsulfanyl-thiazol-2-yl)-malonamide
- N-(2-isobutoxy-phenyl)-N'-(5-methanesulfonyl-thiazol-2-yl)-malonamide
- N-(2-isobutoxy-phenyl)-N'-[5-(pyridin-2-ylsulfanyl)-thiazol-2-yl]-malonamide
- 20 N-[5-(1H-imidazol-2-ylsulfanyl)-thiazol-2-yl]-N'-(2-isobutoxy-phenyl)-malonamide
- {2-[2-(2-isobutoxy-phenylcarbonyl)-acetylamino]-thiazol-4-yl}-acetic acid
- {5-bromo-2-[2-(2-isobutoxy-phenylcarbonyl)-acetylamino]-thiazol-4-yl}-acetic acid
- {5-chloro-2-[2-(2-isobutoxy-phenylcarbonyl)-acetylamino]-thiazol-4-yl}-acetic acid
- {2-[2-(2-isobutoxy-phenylcarbonyl)-acetylamino]-5-methylsulfanyl-thiazol-4-yl}-acetic acid
- 25 {2-[2-(2-isobutoxy-phenylcarbonyl)-acetylamino]-5-methanesulfonyl-thiazol-4-yl}-acetic acid
- [2-[2-(2-isobutoxy-phenylcarbonyl)-acetylamino]-5-(pyridin-2-ylsulfanyl)-thiazol-4-yl]-acetic acid
- {5-(1H-imidazol-2-ylsulfanyl)-2-[2-(2-isobutoxy-phenylcarbonyl)-acetylamino]-thiazol-4-yl}-acetic acid
- 30 N-(2-isobutoxy-phenyl)-N'-[4-(methanesulfonylamino-methyl)-thiazol-2-yl]-malonamide
- N-[5-bromo-4-(methanesulfonylamino-methyl)-thiazol-2-yl]-N'-(2-isobutoxy-phenyl)-malonamide
- N-[5-chloro-4-(methanesulfonylamino-methyl)-thiazol-2-yl]-N'-(2-isobutoxy-phenyl)-malonamide

- N-(2-isobutoxy-phenyl)-N'-[4-(methanesulfonylamino-methyl)-5-methylsulfanyl-thiazol-2-yl]-malonamide
- N-(2-isobutoxy-phenyl)-N'-[5-methanesulfonyl-4-(methanesulfonylamino-methyl)-thiazol-2-yl]-malonamide
- 5 N-(2-isobutoxy-phenyl)-N'-[4-(methanesulfonylamino-methyl)-5-(pyridin-2-ylsulfanyl)-thiazol-2-yl]-malonamide
- N-[5-(1H-imidazol-2-ylsulfanyl)-4-(methanesulfonylamino-methyl)-thiazol-2-yl]-N'-(2-isobutoxy-phenyl)-malonamide
- N-[2-(2-methoxy-phenoxy)-phenyl]-N'-thiazol-2-yl-oxalamide
- 10 N-(5-bromo-thiazol-2-yl)-N'-[2-(2-methoxy-phenoxy)-phenyl]-oxalamide
- N-(5-chloro-thiazol-2-yl)-N'-[2-(2-methoxy-phenoxy)-phenyl]-oxalamide
- N-[2-(2-methoxy-phenoxy)-phenyl]-N'-(5-methylsulfanyl-thiazol-2-yl)-oxalamide
- N-(5-methanesulfonyl-thiazol-2-yl)-N'-[2-(2-methoxy-phenoxy)-phenyl]-oxalamide
- N-[2-(2-methoxy-phenoxy)-phenyl]-N'-[5-(pyridin-2-ylsulfanyl)-thiazol-2-yl]-oxalamide
- 15 N-[5-(1H-imidazol-2-ylsulfanyl)-thiazol-2-yl]-N'-[2-(2-methoxy-phenoxy)-phenyl]-oxalamide
- (2-[[2-(2-methoxy-phenoxy)-phenylaminooxalyl]-amino]-thiazol-4-yl)-acetic acid
- (5-bromo-2-[[2-(2-methoxy-phenoxy)-phenylaminooxalyl]-amino]-thiazol-4-yl)-acetic acid
- (5-chloro-2-[[2-(2-methoxy-phenoxy)-phenylaminooxalyl]-amino]-thiazol-4-yl)-acetic acid
- (2-[[2-(2-methoxy-phenoxy)-phenylaminooxalyl]-amino]-5-methylsulfanyl-thiazol-4-yl)-acetic acid
- 20 acid
- (5-methanesulfonyl-2-[[2-(2-methoxy-phenoxy)-phenylaminooxalyl]-amino]-thiazol-4-yl)-acetic acid
- [2-[[2-(2-methoxy-phenoxy)-phenylaminooxalyl]-amino]-5-(pyridin-2-ylsulfanyl)-thiazol-4-yl]-acetic acid
- 25 (5-(1H-imidazol-2-ylsulfanyl)-2-[[2-(2-methoxy-phenoxy)-phenylaminooxalyl]-amino]-thiazol-4-yl)-acetic acid
- N-[4-(methanesulfonylamino-methyl)-thiazol-2-yl]-N'-[2-(2-methoxy-phenoxy)-phenyl]-oxalamide
- N-[5-bromo-4-(methanesulfonylamino-methyl)-thiazol-2-yl]-N'-[2-(2-methoxy-phenoxy)-phenyl]-oxalamide
- 30 N-[5-chloro-4-(methanesulfonylamino-methyl)-thiazol-2-yl]-N'-[2-(2-methoxy-phenoxy)-phenyl]-oxalamide
- N-[4-(methanesulfonylamino-methyl)-5-methylsulfanyl-thiazol-2-yl]-N'-[2-(2-methoxy-phenoxy)-phenyl]-oxalamide

- N-[5-methanesulfonyl-4-(methanesulfonylamino-methyl)-thiazol-2-yl]-N'-[2-(2-methoxy-phenoxy)-phenyl]-oxalamide
- N-[4-(methanesulfonylamino-methyl)-5-(pyridin-2-ylsulfanyl)-thiazol-2-yl]-N'-[2-(2-methoxy-phenoxy)-phenyl]-oxalamide
- 5 N-[5-(1H-imidazol-2-ylsulfanyl)-4-(methanesulfonylamino-methyl)-thiazol-2-yl]-N'-[2-(2-methoxy-phenoxy)-phenyl]-oxalamide
- N-[2-(2-methoxy-phenoxy)-phenyl]-N'-thiazol-2-yl-malonamide
- N-(5-bromo-thiazol-2-yl)-N'-[2-(2-methoxy-phenoxy)-phenyl]-malonamide
- N-(5-chloro-thiazol-2-yl)-N'-[2-(2-methoxy-phenoxy)-phenyl]-malonamide
- 10 N-[2-(2-methoxy-phenoxy)-phenyl]-N'-(5-methylsulfanyl-thiazol-2-yl)-malonamide
- N-(5-methanesulfonyl-thiazol-2-yl)-N'-[2-(2-methoxy-phenoxy)-phenyl]-malonamide
- N-[2-(2-methoxy-phenoxy)-phenyl]-N'-[5-(pyridin-2-ylsulfanyl)-thiazol-2-yl]-malonamide
- N-[5-(1H-imidazol-2-ylsulfanyl)-thiazol-2-yl]-N'-[2-(2-methoxy-phenoxy)-phenyl]-malonamide
- (2-{2-[2-(2-methoxy-phenoxy)-phenyl]carbamoyl}-acetyl-amino)-thiazol-4-yl)-acetic acid
- 15 (5-bromo-2-{2-[2-(2-methoxy-phenoxy)-phenyl]carbamoyl}-acetyl-amino)-thiazol-4-yl)-acetic acid
- (5-chloro-2-{2-[2-(2-methoxy-phenoxy)-phenyl]carbamoyl}-acetyl-amino)-thiazol-4-yl)-acetic acid
- (2-{2-[2-(2-methoxy-phenoxy)-phenyl]carbamoyl}-acetyl-amino)-5-methylsulfanyl-thiazol-4-yl)-acetic acid
- 20 (5-methanesulfonyl-2-{2-[2-(2-methoxy-phenoxy)-phenyl]carbamoyl}-acetyl-amino)-thiazol-4-yl)-acetic acid
- [2-{2-[2-(2-methoxy-phenoxy)-phenyl]carbamoyl}-acetyl-amino)-5-(pyridin-2-ylsulfanyl)-thiazol-4-yl)-acetic acid
- 25 (5-(1H-imidazol-2-ylsulfanyl)-2-{2-[2-(2-methoxy-phenoxy)-phenyl]carbamoyl}-acetyl-amino)-thiazol-4-yl)-acetic acid
- N[4-(methanesulfonylamino-methyl)-thiazol-2-yl]-N'-[2-(2-methoxy-phenoxy)-phenyl]-malonamide
- N-[5-bromo-4-(methanesulfonylamino-methyl)-thiazol-2-yl]-N'-[2-(2-methoxy-phenoxy)-phenyl]-malonamide
- 30 N-[5-chloro-4-(methanesulfonylamino-methyl)-thiazol-2-yl]-N'-[2-(2-methoxy-phenoxy)-phenyl]-malonamide
- N-[4-(methanesulfonylamino-methyl)-5-methylsulfanyl-thiazol-2-yl]-N'-[2-(2-methoxy-phenoxy)-phenyl]-malonamide

- N-[5-methanesulfonyl-4-(methanesulfonylamino-methyl)-thiazol-2-yl]-N'-[2-(2-methoxy-phenoxy)-phenyl]-malonamide
- N-[4-(methanesulfonylamino-methyl)-5-(pyridin-2-ylsulfanyl)-thiazol-2-yl]-N'-[2-(2-methoxy-phenoxy)-phenyl]-malonamide
- 5 N-[5-(1H-imidazol-2-ylsulfanyl)-4-(methanesulfonylamino-methyl)-thiazol-2-yl]-N'-[2-(2-methoxy-phenoxy)-phenyl]-malonamide
- N-(2-cyclopentanecarbonyl-4-methyl-phenyl)-N'-thiazol-2-yl-oxalamide
- N-(5-bromo-thiazol-2-yl)-N'-(2-cyclopentanecarbonyl-4-methyl-phenyl)-oxalamide
- N-(5-chloro-thiazol-2-yl)-N'-(2-cyclopentanecarbonyl-4-methyl-phenyl)-oxalamide
- 10 N-(2-cyclopentanecarbonyl-4-methyl-phenyl)-N'-(5-methylsulfanyl-thiazol-2-yl)-oxalamide
- N-(2-cyclopentanecarbonyl-4-methyl-phenyl)-N'-(5-methanesulfonyl-thiazol-2-yl)-oxalamide
- N-(2-cyclopentanecarbonyl-4-methyl-phenyl)-N'-[5-(pyridin-2-ylsulfanyl)-thiazol-2-yl]-oxalamide
- N-(2-cyclopentanecarbonyl-4-methyl-phenyl)-N'-[5-(1H-imidazol-2-ylsulfanyl)-thiazol-2-yl]-oxalamide
- 15 {2-[(2-cyclopentanecarbonyl-4-methyl-phenylaminooxalyl)-amino]-thiazol-4-yl}-acetic acid
- {5-bromo-2-[(2-cyclopentanecarbonyl-4-methyl-phenylaminooxalyl)-amino]-thiazol-4-yl}-acetic acid
- {5-chloro-2-[(2-cyclopentanecarbonyl-4-methyl-phenylaminooxalyl)-amino]-thiazol-4-yl}-acetic acid
- 20 {2-[(2-cyclopentanecarbonyl-4-methyl-phenylaminooxalyl)-amino]-5-methylsulfanyl-thiazol-4-yl}-acetic acid
- {2-[(2-cyclopentanecarbonyl-4-methyl-phenylaminooxalyl)-amino]-5-methanesulfonyl-thiazol-4-yl}-acetic acid
- 25 [2-[(2-cyclopentanecarbonyl-4-methyl-phenylaminooxalyl)-amino]-5-(pyridin-2-ylsulfanyl)-thiazol-4-yl]-acetic acid
- [2-[(2-cyclopentanecarbonyl-4-methyl-phenylaminooxalyl)-amino]-5-(1H-imidazol-2-ylsulfanyl)-thiazol-4-yl]-acetic acid
- N-(2-cyclopentanecarbonyl-4-methyl-phenyl)-N'-[4-(methanesulfonylamino-methyl)-thiazol-2-yl]-oxalamide
- 30 N-[5-bromo-4-(methanesulfonylamino-methyl)-thiazol-2-yl]-N'-(2-cyclopentanecarbonyl-4-methyl-phenyl)-oxalamide
- N-[5-chloro-4-(methanesulfonylamino-methyl)-thiazol-2-yl]-N'-(2-cyclopentanecarbonyl-4-methyl-phenyl)-oxalamide

- N-(2-cyclopentanecarbonyl-4-methyl-phenyl)-N'-[4-(methanesulfonylamino-methyl)-5-methylsulfanyl-thiazol-2-yl]-oxalamide
- N-(2-cyclopentanecarbonyl-4-methyl-phenyl)-N'-[5-methanesulfonyl-4-(methanesulfonylamino-methyl)-thiazol-2-yl]-oxalamide
- 5 N-(2-cyclopentanecarbonyl-4-methyl-phenyl)-N'-[4-(methanesulfonylamino-methyl)-5-(pyridin-2-ylsulfanyl)-thiazol-2-yl]-oxalamide
- N-(2-cyclopentanecarbonyl-4-methyl-phenyl)-N'-[5-(1H-imidazol-2-ylsulfanyl)-4-(methanesulfonylamino-methyl)-thiazol-2-yl]-oxalamide
- N-(2-cyclopentanecarbonyl-4-methyl-phenyl)-N'-thiazol-2-yl-malonamide
- 10 N-(5-bromo-thiazol-2-yl)-N'-(2-cyclopentanecarbonyl-4-methyl-phenyl)-malonamide
- N-(5-chloro-thiazol-2-yl)-N'-(2-cyclopentanecarbonyl-4-methyl-phenyl)-malonamide
- N-(2-cyclopentanecarbonyl-4-methyl-phenyl)-N'-(5-methylsulfanyl-thiazol-2-yl)-malonamide
- N-(2-cyclopentanecarbonyl-4-methyl-phenyl)-N'-(5-methanesulfonyl-thiazol-2-yl)-malonamide
- N-(2-cyclopentanecarbonyl-4-methyl-phenyl)-N'-[5-(pyridin-2-ylsulfanyl)-thiazol-2-yl]-
- 15 malonamide
- N-(2-cyclopentanecarbonyl-4-methyl-phenyl)-N'-[5-(1H-imidazol-2-ylsulfanyl)-thiazol-2-yl]-malonamide
- {2-[2-(2-cyclopentanecarbonyl-4-methyl-phenylcarbamoyl)-acetylamino]-thiazol-4-yl}-acetic acid
- 20 {5-bromo-2-[2-(2-cyclopentanecarbonyl-4-methyl-phenylcarbamoyl)-acetylamino]-thiazol-4-yl}-acetic acid
- {5-chloro-2-[2-(2-cyclopentanecarbonyl-4-methyl-phenylcarbamoyl)-acetylamino]-thiazol-4-yl}-acetic acid
- {2-[2-(2-cyclopentanecarbonyl-4-methyl-phenylcarbamoyl)-acetylamino]-5-methylsulfanyl-thiazol-4-yl}-acetic acid
- 25 {2-[2-(2-cyclopentanecarbonyl-4-methyl-phenylcarbamoyl)-acetylamino]-5-methanesulfonyl-thiazol-4-yl}-acetic acid
- [2-[2-(2-cyclopentanecarbonyl-4-methyl-phenylcarbamoyl)-acetylamino]-5-(pyridin-2-ylsulfanyl)-thiazol-4-yl]-acetic acid
- 30 [2-[2-(2-cyclopentanecarbonyl-4-methyl-phenylcarbamoyl)-acetylamino]-5-(1H-imidazol-2-ylsulfanyl)-thiazol-4-yl]-acetic acid
- N-(2-cyclopentanecarbonyl-4-methyl-phenyl)-N'-[4-(methanesulfonylamino-methyl)-thiazol-2-yl]-malonamide
- N-[5-bromo-4-(methanesulfonylamino-methyl)-thiazol-2-yl]-N'-(2-cyclopentanecarbonyl-4-
- 35 methyl-phenyl)-malonamide

- N-[5-chloro-4-(methanesulfonylamino-methyl)-thiazol-2-yl]-N'-(2-cyclopentanecarbonyl-4-methyl-phenyl)-malonamide
165;165;N-(2-cyclopentanecarbonyl-4-methyl-phenyl)-N'-[4-(methanesulfonylamino-methyl)-5-methylsulfanyl-thiazol-2-yl]-malonamide
- 5 N-(2-cyclopentanecarbonyl-4-methyl-phenyl)-N'-[5-methanesulfonyl-4-(methanesulfonylamino-methyl)-thiazol-2-yl]-malonamide
N-(2-cyclopentanecarbonyl-4-methyl-phenyl)-N'-[4-(methanesulfonylamino-methyl)-5-(pyridin-2-ylsulfanyl)-thiazol-2-yl]-malonamide
N-(2-cyclopentanecarbonyl-4-methyl-phenyl)-N'-[5-(1H-imidazol-2-ylsulfanyl)-4-(methanesulfonylamino-methyl)-thiazol-2-yl]-malonamide
- 10 N-(2-isobutoxy-4-methyl-phenyl)-N'-thiazol-2-yl-oxalamide
N-(5-bromo-thiazol-2-yl)-N'-(2-isobutoxy-4-methyl-phenyl)-oxalamide
N-(5-chloro-thiazol-2-yl)-N'-(2-isobutoxy-4-methyl-phenyl)-oxalamide
N-(2-isobutoxy-4-methyl-phenyl)-N'-(5-methylsulfanyl-thiazol-2-yl)-oxalamide
- 15 N-(2-isobutoxy-4-methyl-phenyl)-N'-(5-methanesulfonyl-thiazol-2-yl)-oxalamide
N-(2-isobutoxy-4-methyl-phenyl)-N'-[5-(pyridin-2-ylsulfanyl)-thiazol-2-yl]-oxalamide
N-[5-(1H-imidazol-2-ylsulfanyl)-thiazol-2-yl]-N'-(2-isobutoxy-4-methyl-phenyl)-oxalamide
2-[(2-isobutoxy-4-methyl-phenylaminooxalyl)-amino]-thiazol-4-yl)-acetic acid
5-bromo-2-[(2-isobutoxy-4-methyl-phenylaminooxalyl)-amino]-thiazol-4-yl)-acetic acid
- 20 5-chloro-2-[(2-isobutoxy-4-methyl-phenylaminooxalyl)-amino]-thiazol-4-yl)-acetic acid
2-[(2-isobutoxy-4-methyl-phenylaminooxalyl)-amino]-5-methylsulfanyl-thiazol-4-yl)-acetic acid
2-[(2-isobutoxy-4-methyl-phenylaminooxalyl)-amino]-5-methanesulfonyl-thiazol-4-yl)-acetic acid
- 25 2-[(2-isobutoxy-4-methyl-phenylaminooxalyl)-amino]-5-(pyridin-2-ylsulfanyl)-thiazol-4-yl)-acetic acid
5-(1H-imidazol-2-ylsulfanyl)-2-[(2-isobutoxy-4-methyl-phenylaminooxalyl)-amino]-thiazol-4-yl)-acetic acid
N-(2-isobutoxy-4-methyl-phenyl)-N'-[4-(methanesulfonylamino-methyl)-thiazol-2-yl]-oxalamide
- 30 N-[5-bromo-4-(methanesulfonylamino-methyl)-thiazol-2-yl]-N'-(2-isobutoxy-4-methyl-phenyl)-oxalamide
N-[5-chloro-4-(methanesulfonylamino-methyl)-thiazol-2-yl]-N'-(2-isobutoxy-4-methyl-phenyl)-oxalamide

- N-(2-isobutoxy-4-methyl-phenyl)-N'-[4-(methanesulfonylamino-methyl)-5-methylsulfanyl-thiazol-2-yl]-oxalamide
- N-(2-isobutoxy-4-methyl-phenyl)-N'-[5-methanesulfonyl-4-(methanesulfonylamino-methyl)-thiazol-2-yl]-oxalamide
- 5 N-(2-isobutoxy-4-methyl-phenyl)-N'-[4-(methanesulfonylamino-methyl)-5-(pyridin-2-ylsulfanyl)-thiazol-2-yl]-oxalamide
- N-[5-(1H-imidazol-2-ylsulfanyl)-4-(methanesulfonylamino-methyl)-thiazol-2-yl]-N'-(2-isobutoxy-4-methyl-phenyl)-oxalamide
- N-(2-isobutoxy-4-methyl-phenyl)-N'-thiazol-2-yl-malonamide
- 10 N-(5-bromo-thiazol-2-yl)-N'-(2-isobutoxy-4-methyl-phenyl)-malonamide
- N-(5-chloro-thiazol-2-yl)-N'-(2-isobutoxy-4-methyl-phenyl)-malonamide
- N-(2-isobutoxy-4-methyl-phenyl)-N'-(5-methylsulfanyl-thiazol-2-yl)-malonamide
- N-(2-isobutoxy-4-methyl-phenyl)-N'-(5-methanesulfonyl-thiazol-2-yl)-malonamide
- N-(2-isobutoxy-4-methyl-phenyl)-N'-[5-(pyridin-2-ylsulfanyl)-thiazol-2-yl]-malonamide
- 15 N-[5-(1H-imidazol-2-ylsulfanyl)-thiazol-2-yl]-N'-(2-isobutoxy-4-methyl-phenyl)-malonamide
- {2-[2-(2-isobutoxy-4-methyl-phenylcarbamoyl)-acetylamino]-thiazol-4-yl}-acetic acid
- {5-bromo-2-[2-(2-isobutoxy-4-methyl-phenylcarbamoyl)-acetylamino]-thiazol-4-yl}-acetic acid
- {5-chloro-2-[2-(2-isobutoxy-4-methyl-phenylcarbamoyl)-acetylamino]-thiazol-4-yl}-acetic acid
- {2-[2-(2-isobutoxy-4-methyl-phenylcarbamoyl)-acetylamino]-5-methylsulfanyl-thiazol-4-yl}-
- 20 acetic acid
- {2-[2-(2-isobutoxy-4-methyl-phenylcarbamoyl)-acetylamino]-5-methanesulfonyl-thiazol-4-yl}-acetic acid
- [2-[2-(2-isobutoxy-4-methyl-phenylcarbamoyl)-acetylamino]-5-(pyridin-2-ylsulfanyl)-thiazol-4-yl]-acetic acid
- 25 {5-(1H-imidazol-2-ylsulfanyl)-2-[2-(2-isobutoxy-4-methyl-phenylcarbamoyl)-acetylamino]-thiazol-4-yl}-acetic acid
- N-(2-isobutoxy-4-methyl-phenyl)-N'-[4-(methanesulfonylamino-methyl)-thiazol-2-yl]-malonamide
- N-[5-bromo-4-(methanesulfonylamino-methyl)-thiazol-2-yl]-N'-(2-isobutoxy-4-methyl-phenyl)-malonamide
- 30 malonamide
- N-[5-chloro-4-(methanesulfonylamino-methyl)-thiazol-2-yl]-N'-(2-isobutoxy-4-methyl-phenyl)-malonamide
- N-(2-isobutoxy-4-methyl-phenyl)-N'-[4-(methanesulfonylamino-methyl)-5-methylsulfanyl-thiazol-2-yl]-malonamide

- N-(2-isobutoxy-4-methyl-phenyl)-N'-[5-methanesulfonyl-4-(methanesulfonylamino-methyl)-thiazol-2-yl]-malonamide
- N-(2-isobutoxy-4-methyl-phenyl)-N'-[4-(methanesulfonylamino-methyl)-5-(pyridin-2-ylsulfanyl)-thiazol-2-yl]-malonamide
- 5 N-[5-(1H-imidazol-2-ylsulfanyl)-4-(methanesulfonylamino-methyl)-thiazol-2-yl]-N'-(2-isobutoxy-4-methyl-phenyl)-malonamide
- N-[2-(2-methoxy-phenoxy)-4-methyl-phenyl]-N'-thiazol-2-yl-oxalamide
- N-(5-bromo-thiazol-2-yl)-N'-[2-(2-methoxy-phenoxy)-4-methyl-phenyl]-oxalamide
- N-(5-chloro-thiazol-2-yl)-N'-[2-(2-methoxy-phenoxy)-4-methyl-phenyl]-oxalamide
- 10 N-[N-(5-methanesulfonyl-thiazol-N-[N-[5-(1H-imidazol-((5-bromo-(5-chloro-((5-methanesulfonyl-((5-(1H-imidazol-N-[4-(methanesulfonylamino-methyl)-thiazol-N-[5-bromo-4-(methanesulfonylamino-methyl)-thiazol-N-[5-chloro-4-(methanesulfonylamino-methyl)-thiazol-N-[4-(methanesulfonylamino-methyl)-5-methylsulfanyl-thiazol-N-[5-methanesulfonyl-4-(methanesulfonylamino-methyl)-thiazol-N-[4-(methanesulfonylamino-methyl)-5-(pyridin-N-[5-
- 15 (1H-imidazol-N-[N-(5-bromo-thiazol-N-(5-chloro-thiazol-N-[N-(5-methanesulfonyl-thiazol-N-[N-[5-(1H-imidazol-((5-bromo-(5-chloro-((5-methanesulfonyl-((5-(1H-imidazol-N-[4-(methanesulfonylamino-methyl)-thiazol-N-[5-bromo-4-(methanesulfonylamino-methyl)-thiazol-N-[5-chloro-4-(methanesulfonylamino-methyl)-thiazol-N-[4-(methanesulfonylamino-methyl)-5-methylsulfanyl-thiazol-N-[5-methanesulfonyl-4-(methanesulfonylamino-methyl)-thiazol-N-[4-(methanesulfonylamino-methyl)-5-(pyridin-N-[5-(1H-imidazol-N-(N-(5-bromo-thiazol-N-(5-chloro-thiazol-N-(N-(N-(N-(((5-bromo-{5-chloro-{[[[N-(N-[5-bromo-4-(methanesulfonylamino-methyl)-thiazol-N-[5-chloro-4-(methanesulfonylamino-methyl)-thiazol-N-(N-(N-(N-(N-(5-bromo-thiazol-N-(5-chloro-thiazol-N-(N-(N-(N-(((5-bromo-{5-chloro-{[[[N-(N-[5-bromo-4-(methanesulfonylamino-methyl)-thiazol-N-[5-chloro-4-(methanesulfonylamino-methyl)-thiazol-N-(N-(N-(N-(N-(4-fluoro-N-(5-bromo-thiazol-N-(5-chloro-thiazol-N-(4-fluoro-N-(4-fluoro-2-isobutoxy-phenyl)-N'-(5-methanesulfonyl-thiazol-2-yl)-oxalamide
- 20 N-(4-fluoro-2-isobutoxy-phenyl)-N'-[5-(pyridin-2-ylsulfanyl)-thiazol-2-yl]-oxalamide
- N-(4-fluoro-2-isobutoxy-phenyl)-N'-[5-(1H-imidazol-2-ylsulfanyl)-thiazol-2-yl]-oxalamide
- {2-[(4-fluoro-2-isobutoxy-phenylamino)oxalyl]-amino]-thiazol-4-yl}-acetic acid
- 25 {5-bromo-2-[(4-fluoro-2-isobutoxy-phenylamino)oxalyl]-amino]-thiazol-4-yl}-acetic acid
- {5-chloro-2-[(4-fluoro-2-isobutoxy-phenylamino)oxalyl]-amino]-thiazol-4-yl}-acetic acid
- {2-[(4-fluoro-2-isobutoxy-phenylamino)oxalyl]-amino]-5-methylsulfanyl-thiazol-4-yl}-acetic acid
- {2-[(4-fluoro-2-isobutoxy-phenylamino)oxalyl]-amino]-5-methanesulfonyl-thiazol-4-yl}-acetic acid
- 30 acid

- [2-[(4-fluoro-2-isobutoxy-phenylamino)oxalyl]-amino]-5-(pyridin-2-ylsulfanyl)-thiazol-4-yl]-acetic acid
- [2-[(4-fluoro-2-isobutoxy-phenylamino)oxalyl]-amino]-5-(1H-imidazol-2-ylsulfanyl)-thiazol-4-yl]-acetic acid
- 5 N-(4-fluoro-2-isobutoxy-phenyl)-N'-[4-(methanesulfonylamino-methyl)-thiazol-2-yl]-oxalamide
N-[5-bromo-4-(methanesulfonylamino-methyl)-thiazol-2-yl]-N'-(4-fluoro-2-isobutoxy-phenyl)-oxalamide
N-[5-chloro-4-(methanesulfonylamino-methyl)-thiazol-2-yl]-N'-(4-fluoro-2-isobutoxy-phenyl)-oxalamide
- 10 N-(4-fluoro-2-isobutoxy-phenyl)-N'-[4-(methanesulfonylamino-methyl)-5-methylsulfanyl-thiazol-2-yl]-oxalamide
N-(4-fluoro-2-isobutoxy-phenyl)-N'-[5-methanesulfonyl-4-(methanesulfonylamino-methyl)-thiazol-2-yl]-oxalamide
N-(4-fluoro-2-isobutoxy-phenyl)-N'-[4-(methanesulfonylamino-methyl)-5-(pyridin-2-ylsulfanyl)-thiazol-2-yl]-oxalamide
- 15 N-(4-fluoro-2-isobutoxy-phenyl)-N'-[5-(1H-imidazol-2-ylsulfanyl)-4-(methanesulfonylamino-methyl)-thiazol-2-yl]-oxalamide
N-(4-fluoro-2-isobutoxy-phenyl)-N'-thiazol-2-yl-malonamide
N-(5-bromo-thiazol-2-yl)-N'-(4-fluoro-2-isobutoxy-phenyl)-malonamide
- 20 N-(5-chloro-thiazol-2-yl)-N'-(4-fluoro-2-isobutoxy-phenyl)-malonamide
N-(4-fluoro-2-isobutoxy-phenyl)-N'-(5-methylsulfanyl-thiazol-2-yl)-malonamide
N-(4-fluoro-2-isobutoxy-phenyl)-N'-(5-methanesulfonyl-thiazol-2-yl)-malonamide
N-(4-fluoro-2-isobutoxy-phenyl)-N'-[5-(pyridin-2-ylsulfanyl)-thiazol-2-yl]-malonamide
N-(4-fluoro-2-isobutoxy-phenyl)-N'-[5-(1H-imidazol-2-ylsulfanyl)-thiazol-2-yl]-malonamide
- 25 {2-[2-(4-fluoro-2-isobutoxy-phenylcarbonyl)-acetyl]amino}-thiazol-4-yl]-acetic acid
{5-bromo-2-[2-(4-fluoro-2-isobutoxy-phenylcarbonyl)-acetyl]amino}-thiazol-4-yl]-acetic acid
{5-chloro-2-[2-(4-fluoro-2-isobutoxy-phenylcarbonyl)-acetyl]amino}-thiazol-4-yl]-acetic acid
{2-[2-(4-fluoro-2-isobutoxy-phenylcarbonyl)-acetyl]amino}-5-methylsulfanyl-thiazol-4-yl]-acetic acid
- 30 {2-[2-(4-fluoro-2-isobutoxy-phenylcarbonyl)-acetyl]amino}-5-methanesulfonyl-thiazol-4-yl]-acetic acid
{2-[2-(4-fluoro-2-isobutoxy-phenylcarbonyl)-acetyl]amino}-5-(pyridin-2-ylsulfanyl)-thiazol-4-yl]-acetic acid
{2-[2-(4-fluoro-2-isobutoxy-phenylcarbonyl)-acetyl]amino}-5-(1H-imidazol-2-ylsulfanyl)-thiazol-4-yl]-acetic acid
- 35

- N-(4-fluoro-2-isobutoxy-phenyl)-N'-[4-(methanesulfonylamino-methyl)-thiazol-2-yl]-malonamide
- N-[5-bromo-4-(methanesulfonylamino-methyl)-thiazol-2-yl]-N'-(4-fluoro-2-isobutoxy-phenyl)-malonamide
- 5 N-[5-chloro-4-(methanesulfonylamino-methyl)-thiazol-2-yl]-N'-(4-fluoro-2-isobutoxy-phenyl)-malonamide
- N-(4-fluoro-2-isobutoxy-phenyl)-N'-[4-(methanesulfonylamino-methyl)-5-methylsulfanyl-thiazol-2-yl]-malonamide
- N-(4-fluoro-2-isobutoxy-phenyl)-N'-[5-methanesulfonyl-4-(methanesulfonylamino-methyl)-thiazol-2-yl]-malonamide
- 10 N-(4-fluoro-2-isobutoxy-phenyl)-N'-[4-(methanesulfonylamino-methyl)-5-(pyridin-2-ylsulfanyl)-thiazol-2-yl]-malonamide
- N-(4-fluoro-2-isobutoxy-phenyl)-N'-[5-(1H-imidazol-2-ylsulfanyl)-4-(methanesulfonylamino-methyl)-thiazol-2-yl]-malonamide
- 15 N-[4-fluoro-2-(2-methoxy-phenoxy)-phenyl]-N'-thiazol-2-yl-oxalamide
- N-(5-bromo-thiazol-2-yl)-N'-[4-fluoro-2-(2-methoxy-phenoxy)-phenyl]-oxalamide
- N-(5-chloro-thiazol-2-yl)-N'-[4-fluoro-2-(2-methoxy-phenoxy)-phenyl]-oxalamide
- N-[4-fluoro-2-(2-methoxy-phenoxy)-phenyl]-N'-(5-methylsulfanyl-thiazol-2-yl)-oxalamide
- N-[4-fluoro-2-(2-methoxy-phenoxy)-phenyl]-N'-(5-methanesulfonyl-thiazol-2-yl)-oxalamide
- 20 N-[4-fluoro-2-(2-methoxy-phenoxy)-phenyl]-N'-[5-(pyridin-2-ylsulfanyl)-thiazol-2-yl]-oxalamide
- N-[4-fluoro-2-(2-methoxy-phenoxy)-phenyl]-N'-[5-(1H-imidazol-2-ylsulfanyl)-thiazol-2-yl]-oxalamide
- (2-[[4-fluoro-2-(2-methoxy-phenoxy)-phenylaminooxalyl]-amino]-thiazol-4-yl)-acetic acid
- (5-bromo-2-[[4-fluoro-2-(2-methoxy-phenoxy)-phenylaminooxalyl]-amino]-thiazol-4-yl)-acetic acid
- 25 acid
- (5-chloro-2-[[4-fluoro-2-(2-methoxy-phenoxy)-phenylaminooxalyl]-amino]-thiazol-4-yl)-acetic acid
- (2-[[4-fluoro-2-(2-methoxy-phenoxy)-phenylaminooxalyl]-amino]-5-methylsulfanyl-thiazol-4-yl)-acetic acid
- 30 (2-[[4-fluoro-2-(2-methoxy-phenoxy)-phenylaminooxalyl]-amino]-5-methanesulfonyl-thiazol-4-yl)-acetic acid
- [2-[[4-fluoro-2-(2-methoxy-phenoxy)-phenylaminooxalyl]-amino]-5-(pyridin-2-ylsulfanyl)-thiazol-4-yl]-acetic acid
- [2-[[4-fluoro-2-(2-methoxy-phenoxy)-phenylaminooxalyl]-amino]-5-(1H-imidazol-2-ylsulfanyl)-thiazol-4-yl]-acetic acid
- 35 thiazol-4-yl]-acetic acid

- N-[4-fluoro-2-(2-methoxy-phenoxy)-phenyl]-N'-[4-(methanesulfonylamino-methyl)-thiazol-2-yl]-oxalamide
- N-[5-bromo-4-(methanesulfonylamino-methyl)-thiazol-2-yl]-N'-[4-fluoro-2-(2-methoxy-phenoxy)-phenyl]-oxalamide
- 5 N-[5-chloro-4-(methanesulfonylamino-methyl)-thiazol-2-yl]-N'-[4-fluoro-2-(2-methoxy-phenoxy)-phenyl]-oxalamide
- N-[4-fluoro-2-(2-methoxy-phenoxy)-phenyl]-N'-[4-(methanesulfonylamino-methyl)-5-methylsulfanyl-thiazol-2-yl]-oxalamide
- N-[4-fluoro-2-(2-methoxy-phenoxy)-phenyl]-N'-[5-methanesulfonyl-4-(methanesulfonylamino-methyl)-thiazol-2-yl]-oxalamide
- 10 N-[4-fluoro-2-(2-methoxy-phenoxy)-phenyl]-N'-[4-(methanesulfonylamino-methyl)-5-(pyridin-2-ylsulfanyl)-thiazol-2-yl]-oxalamide
- N-[4-fluoro-2-(2-methoxy-phenoxy)-phenyl]-N'-[5-(1H-imidazol-2-ylsulfanyl)-4-(methanesulfonylamino-methyl)-thiazol-2-yl]-oxalamide
- 15 N-[4-fluoro-2-(2-methoxy-phenoxy)-phenyl]-N'-thiazol-2-yl-malonamide
- N-(5-bromo-thiazol-2-yl)-N'-[4-fluoro-2-(2-methoxy-phenoxy)-phenyl]-malonamide
- N-(5-chloro-thiazol-2-yl)-N'-[4-fluoro-2-(2-methoxy-phenoxy)-phenyl]-malonamide
- N-[4-fluoro-2-(2-methoxy-phenoxy)-phenyl]-N'-(5-methylsulfanyl-thiazol-2-yl)-malonamide
- N-[4-fluoro-2-(2-methoxy-phenoxy)-phenyl]-N'-(5-methanesulfonyl-thiazol-2-yl)-malonamide
- 20 N-[4-fluoro-2-(2-methoxy-phenoxy)-phenyl]-N'-[5-(pyridin-2-ylsulfanyl)-thiazol-2-yl]-malonamide
- N-[4-fluoro-2-(2-methoxy-phenoxy)-phenyl]-N'-[5-(1H-imidazol-2-ylsulfanyl)-thiazol-2-yl]-malonamide
- (2-{2-[4-fluoro-2-(2-methoxy-phenoxy)-phenylcarbamoyl]-acetylamino}-thiazol-4-yl)-acetic acid
- 25 acid
- (5-bromo-2-{2-[4-fluoro-2-(2-methoxy-phenoxy)-phenylcarbamoyl]-acetylamino}-thiazol-4-yl)-acetic acid
- (5-chloro-2-{2-[4-fluoro-2-(2-methoxy-phenoxy)-phenylcarbamoyl]-acetylamino}-thiazol-4-yl)-acetic acid
- 30 (2-{2-[4-fluoro-2-(2-methoxy-phenoxy)-phenylcarbamoyl]-acetylamino}-5-methylsulfanyl-thiazol-4-yl)-acetic acid
- (2-{2-[4-fluoro-2-(2-methoxy-phenoxy)-phenylcarbamoyl]-acetylamino}-5-methanesulfonyl-thiazol-4-yl)-acetic acid
- [2-{2-[4-fluoro-2-(2-methoxy-phenoxy)-phenylcarbamoyl]-acetylamino}-5-(pyridin-2-ylsulfanyl)-thiazol-4-yl]-acetic acid
- 35

- [2-{2-[4-fluoro-2-(2-methoxy-phenoxy)-phenylcarbamoyl]-acetylamino}-5-(1H-imidazol-2-ylsulfanyl)-thiazol-4-yl]-acetic acid
- N-[4-fluoro-2-(2-methoxy-phenoxy)-phenyl]-N'-[4-(methanesulfonylamino-methyl)-thiazol-2-yl]-malonamide
- 5 N-[5-bromo-4-(methanesulfonylamino-methyl)-thiazol-2-yl]-N'-[4-fluoro-2-(2-methoxy-phenoxy)-phenyl]-malonamide
- N-[5-chloro-4-(methanesulfonylamino-methyl)-thiazol-2-yl]-N'-[4-fluoro-2-(2-methoxy-phenoxy)-phenyl]-malonamide
- N-[4-fluoro-2-(2-methoxy-phenoxy)-phenyl]-N'-[4-(methanesulfonylamino-methyl)-5-methylsulfanyl-thiazol-2-yl]-malonamide
- 10 N-[4-fluoro-2-(2-methoxy-phenoxy)-phenyl]-N'-[5-methanesulfonyl-4-(methanesulfonylamino-methyl)-thiazol-2-yl]-malonamide
- N-[4-fluoro-2-(2-methoxy-phenoxy)-phenyl]-N'-[4-(methanesulfonylamino-methyl)-5-(pyridin-2-ylsulfanyl)-thiazol-2-yl]-malonamide
- 15 N-[4-fluoro-2-(2-methoxy-phenoxy)-phenyl]-N'-[5-(1H-imidazol-2-ylsulfanyl)-4-(methanesulfonylamino-methyl)-thiazol-2-yl]-malonamide
- N-(2-cyclopentanecarbonyl-4-methoxy-phenyl)-N'-thiazol-2-yl-oxalamide
- N-(5-bromo-thiazol-2-yl)-N'-(2-cyclopentanecarbonyl-4-methoxy-phenyl)-oxalamide
- N-(5-chloro-thiazol-2-yl)-N'-(2-cyclopentanecarbonyl-4-methoxy-phenyl)-oxalamide
- 20 N-(2-cyclopentanecarbonyl-4-methoxy-phenyl)-N'-(5-methylsulfanyl-thiazol-2-yl)-oxalamide
- N-(2-cyclopentanecarbonyl-4-methoxy-phenyl)-N'-(5-methanesulfonyl-thiazol-2-yl)-oxalamide
- N-(2-cyclopentanecarbonyl-4-methoxy-phenyl)-N'-[5-(pyridin-2-ylsulfanyl)-thiazol-2-yl]-oxalamide
- N-(2-cyclopentanecarbonyl-4-methoxy-phenyl)-N'-[5-(1H-imidazol-2-ylsulfanyl)-thiazol-2-yl]-oxalamide
- 25 {2-[(2-cyclopentanecarbonyl-4-methoxy-phenylaminooxalyl)-amino]-thiazol-4-yl}-acetic acid
- {5-bromo-2-[(2-cyclopentanecarbonyl-4-methoxy-phenylaminooxalyl)-amino]-thiazol-4-yl}-acetic acid
- {5-chloro-2-[(2-cyclopentanecarbonyl-4-methoxy-phenylaminooxalyl)-amino]-thiazol-4-yl}-acetic acid
- 30 {2-[(2-cyclopentanecarbonyl-4-methoxy-phenylaminooxalyl)-amino]-5-methylsulfanyl-thiazol-4-yl}-acetic acid
- {2-[(2-cyclopentanecarbonyl-4-methoxy-phenylaminooxalyl)-amino]-5-methanesulfonyl-thiazol-4-yl}-acetic acid

- [2-[(2-cyclopentanecarbonyl-4-methoxy-phenylaminooxalyl)-amino]-5-(pyridin-2-ylsulfanyl)-thiazol-4-yl]-acetic acid
- [2-[(2-cyclopentanecarbonyl-4-methoxy-phenylaminooxalyl)-amino]-5-(1H-imidazol-2-ylsulfanyl)-thiazol-4-yl]-acetic acid
- 5 N-(2-cyclopentanecarbonyl-4-methoxy-phenyl)-N'-[4-(methanesulfonylamino-methyl)-thiazol-2-yl]-oxalamide
- N-[5-bromo-4-(methanesulfonylamino-methyl)-thiazol-2-yl]-N'-(2-cyclopentanecarbonyl-4-methoxy-phenyl)-oxalamide
- N-[5-chloro-4-(methanesulfonylamino-methyl)-thiazol-2-yl]-N'-(2-cyclopentanecarbonyl-4-methoxy-phenyl)-oxalamide
- 10 N-(2-cyclopentanecarbonyl-4-methoxy-phenyl)-N'-[4-(methanesulfonylamino-methyl)-5-methylsulfanyl-thiazol-2-yl]-oxalamide
- N-(2-cyclopentanecarbonyl-4-methoxy-phenyl)-N'-[5-methanesulfonyl-4-(methanesulfonylamino-methyl)-thiazol-2-yl]-oxalamide
- 15 N-(2-cyclopentanecarbonyl-4-methoxy-phenyl)-N'-[N-(2-cyclopentanecarbonyl-N-(N-[N-[4-(methanesulfonylamino-methyl)-N-[N-[4-(methanesulfonylamino-methyl)-N-[5-(1H-imidazol-2-ylsulfanyl)-4-(methanesulfonylamino-methyl)-thiazol-2-yl]-N'-[4-methoxy-2-(2-methoxy-phenoxy)-phenyl]-malonamide
- 1-(2-Cyclopentanecarbonyl-4-methyl-phenyl)-3-(5-morpholin-4-ylmethyl-thiazol-2-yl)-urea
- 20 1-(2-Cyclopentanecarbonyl-4-methyl-phenyl)-3-(5-dimethylaminomethyl-thiazol-2-yl)-urea
- 1-(2-Cyclopentanecarbonyl-4-methyl-phenyl)-3-(5-piperazin-1-ylmethyl-thiazol-2-yl)-urea
- 1-(2-Cyclopentanecarbonyl-4-methyl-phenyl)-3-[5-(4-methyl-piperazin-1-ylmethyl)-thiazol-2-yl]-urea
- 1-[5-(4-Acetyl-piperazin-1-ylmethyl)-thiazol-2-yl]-3-(2-cyclopentanecarbonyl-4-methyl-phenyl)-urea
- 25 1-(2-Cyclopentanecarbonyl-4-methyl-phenyl)-3-[5-(4-methanesulfonyl-piperazin-1-ylmethyl)-thiazol-2-yl]-urea
- (4-{2-[3-(2-Cyclopentanecarbonyl-4-methyl-phenyl)-ureido]-thiazol-5-ylmethyl}-piperazin-1-yl)-acetic acid
- 30 2-{2-[3-(2-Cyclopentanecarbonyl-4-methyl-phenyl)-ureido]-thiazol-5-ylmethylsulfanyl}-1H-imidazole-4-carboxylic acid
- 1-(2-Cyclopentanecarbonyl-4-methyl-phenyl)-3-[5-(1H-imidazol-2-ylsulfanylmethyl)-thiazol-2-yl]-urea
- 1-(2-Cyclopentanecarbonyl-4-methyl-phenyl)-3-[5-(1-methyl-1H-imidazol-2-ylsulfanylmethyl)-thiazol-2-yl]-urea
- 35

- 1-(2-Cyclopentanecarbonyl-4-methyl-phenyl)-3-[5-(4-methyl-4H-[1,2,4]triazol-3-ylsulfanylmethyl)-thiazol-2-yl]-urea
- 1-(2-Cyclopentanecarbonyl-4-methyl-phenyl)-3-[5-(4H-[1,2,4]triazol-3-ylsulfanylmethyl)-thiazol-2-yl]-urea
- 5 1-(2-Cyclopentanecarbonyl-4-methyl-phenyl)-3-[5-(1-methyl-1H-tetrazol-5-ylsulfanylmethyl)-thiazol-2-yl]-urea
- (5-{2-[3-(2-Cyclopentanecarbonyl-4-methyl-phenyl)-ureido]-thiazol-5-ylmethylsulfanyl}-tetrazol-1-yl)-acetic acid
- 1-(2-Cyclopentanecarbonyl-4-methyl-phenyl)-3-[5-[1-(2-dimethylamino-ethyl)-1H-tetrazol-5-ylsulfanylmethyl]-thiazol-2-yl]-urea
- 10 (5-{2-[3-(2-Cyclopentanecarbonyl-4-methyl-phenyl)-ureido]-thiazol-5-ylmethanesulfonyl}-tetrazol-1-yl)-acetic acid
- 1-(2-Cyclopentanecarbonyl-4-methyl-phenyl)-3-[5-[1-(2-dimethylamino-ethyl)-1H-tetrazole-5-sulfonylmethyl]-thiazol-2-yl]-urea
- 15 1-(2-Cyclopentanecarbonyl-4-methyl-phenyl)-3-[5-(1-methyl-1H-tetrazole-5-sulfonylmethyl)-thiazol-2-yl]-urea
- 1-(2-Cyclopentanecarbonyl-4-methyl-phenyl)-3-[5-(4H-[1,2,4]triazole-3-sulfonylmethyl)-thiazol-2-yl]-urea
- 1-(2-Cyclopentanecarbonyl-4-methyl-phenyl)-3-[5-(4-methyl-4H-[1,2,4]triazole-3-sulfonylmethyl)-thiazol-2-yl]-urea
- 20 1-(2-Cyclopentanecarbonyl-4-methyl-phenyl)-3-[5-(1-methyl-1H-imidazole-2-sulfonylmethyl)-thiazol-2-yl]-urea
- 1-(2-Cyclopentanecarbonyl-4-methyl-phenyl)-3-[5-(1H-imidazole-2-sulfonylmethyl)-thiazol-2-yl]-urea
- 25 1-(2-Cyclopentanecarbonyl-4-methyl-phenyl)-3-[5-(pyridine-2-sulfonylmethyl)-thiazol-2-yl]-urea
- 1-(2-Cyclopentanecarbonyl-4-methyl-phenyl)-3-[5-(pyridin-2-ylsulfanylmethyl)-thiazol-2-yl]-urea
- 1-(2-Cyclopentanecarbonyl-4-methyl-phenyl)-3-[5-(pyridin-3-ylsulfanylmethyl)-thiazol-2-yl]-urea
- 30 urea
- 1-(2-Cyclopentanecarbonyl-4-methyl-phenyl)-3-[5-(pyridine-3-sulfonylmethyl)-thiazol-2-yl]-urea
- urea
- 1-(2-Cyclopentanecarbonyl-4-methyl-phenyl)-3-[5-(pyridin-4-ylsulfanylmethyl)-thiazol-2-yl]-urea
- urea

- 1-(2-Cyclopentanecarbonyl-4-methyl-phenyl)-3-[5-(pyridine-4-sulfonylmethyl)-thiazol-2-yl]-urea
- 1-(2-Cyclopentanecarbonyl-4-methyl-phenyl)-3-[5-(pyridin-4-yloxymethyl)-thiazol-2-yl]-urea
- 1-(2-Cyclopentanecarbonyl-4-methyl-phenyl)-3-[5-(pyridin-3-yloxymethyl)-thiazol-2-yl]-urea
- 5 1-(2-Cyclopentanecarbonyl-4-methyl-phenyl)-3-[5-(pyridin-2-yloxymethyl)-thiazol-2-yl]-urea
- (4-{2-[3-(2-Cyclopentanecarbonyl-4-methyl-phenyl)-ureido]-thiazol-5-ylmethoxy}-phenyl)-acetic acid
- (4-{2-[3-(2-Cyclopentanecarbonyl-4-methyl-phenyl)-ureido]-thiazol-5-ylmethylsulfanyl}-phenyl)-acetic acid
- 10 (4-{2-[3-(2-Cyclopentanecarbonyl-4-methyl-phenyl)-ureido]-thiazol-5-ylmethanesulfonyl}-phenyl)-acetic acid
- 3-(4-{2-[3-(2-Cyclopentanecarbonyl-4-methyl-phenyl)-ureido]-thiazol-5-ylmethoxy}-phenyl)-propionic acid
- 3-(4-{2-[3-(2-Cyclopentanecarbonyl-4-methyl-phenyl)-ureido]-thiazol-5-ylmethylsulfanyl}-phenyl)-propionic acid
- 15 (4-{2-[3-(2-Cyclopentanecarbonyl-4-methyl-phenyl)-ureido]-thiazol-5-ylmethanesulfonyl}-phenyl)-propionic acid
- 4-{2-[3-(2-Cyclopentanecarbonyl-4-methyl-phenyl)-ureido]-thiazol-5-ylmethylsulfanyl}-benzoic acid
- 20 4-{2-[3-(2-Cyclopentanecarbonyl-4-methyl-phenyl)-ureido]-thiazol-5-ylmethanesulfonyl}-benzoic acid
- 4-{2-[3-(2-Cyclopentanecarbonyl-4-methyl-phenyl)-ureido]-thiazol-5-ylmethoxy}-benzoic acid
- N-{2-[3-(2-Cyclopentanecarbonyl-4-methyl-phenyl)-ureido]-thiazol-5-ylmethyl}-acetamide
- N-{2-[3-(2-Cyclopentanecarbonyl-4-methyl-phenyl)-ureido]-thiazol-5-ylmethyl}-methanesulfonamide
- 25 1-(2-Cyclopentanecarbonyl-4-methyl-phenyl)-3-[5-(pyridin-2-ylaminomethyl)-thiazol-2-yl]-urea
- 2-{2-[3-(2-Cyclopentanecarbonyl-4-methyl-phenyl)-ureido]-thiazol-5-ylmethanesulfonyl}-1H-imidazole-4-carboxylic acid
- 2-(2-{2-[3-(2-Cyclopentanecarbonyl-4-methyl-phenyl)-ureido]-thiazol-5-yl}-ethyl)-1H-imidazole-4-carboxylic acid
- 30 2-(2-{2-[3-(2-Cyclopentanecarbonyl-4-methyl-phenyl)-ureido]-thiazol-5-yl}-ethyl)-oxazole-4-carboxylic acid
- 2-(2-{2-[3-(2-Cyclopentanecarbonyl-4-methyl-phenyl)-ureido]-thiazol-5-yl}-ethyl)-1-methyl-1H-imidazole-4-carboxylic acid
- 35 2-[3-(2-Cyclopentanecarbonyl-4-methyl-phenyl)-ureido]-thiazole-5-carboxylic acid

- 2-[3-(2-Cyclopentanecarbonyl-4-methyl-phenyl)-ureido]-thiazole-5-carboxylic acid amide
2-[3-(2-Cyclopentanecarbonyl-4-methyl-phenyl)-ureido]-thiazole-5-carboxylic acid
methanamide
(2-[3-(2-Cyclopentanecarbonyl-4-methyl-phenyl)-ureido]-thiazole-5-carbonyl)-amino)-acetic
5 acid
(2-[3-(2-Cyclopentanecarbonyl-4-methyl-phenyl)-ureido]-thiazole-5-carbonyl)-
methanesulfonyl-amino)-acetic acid
N-(2-[3-(2-Cyclopentanecarbonyl-4-methyl-phenyl)-ureido]-thiazole-5-carbonyl)-
methanesulfonamide
10 {2-[3-(2-Cyclopentanecarbonyl-4-methyl-phenyl)-ureido]-thiazol-5-yl}-acetic acid
2-{2-[3-(2-Cyclopentanecarbonyl-4-methyl-phenyl)-ureido]-thiazol-5-yl}-acetamide
2-{2-[3-(2-Cyclopentanecarbonyl-4-methyl-phenyl)-ureido]-thiazol-5-yl}-N-methyl-acetamide
(2-{2-[3-(2-Cyclopentanecarbonyl-4-methyl-phenyl)-ureido]-thiazol-5-yl}-acetyl-amino)-acetic
acid
15 N-(2-{2-[3-(2-Cyclopentanecarbonyl-4-methyl-phenyl)-ureido]-thiazol-5-yl}-acetyl)-
methanesulfonamide
N-(2-{2-[3-(2-Cyclopentanecarbonyl-4-methyl-phenyl)-ureido]-thiazol-5-yl}-ethyl)-
methanesulfonamide
N-(2-{2-[3-(2-Cyclopentanecarbonyl-4-methyl-phenyl)-ureido]-thiazol-5-yl}-ethyl)-acetamide
20 1-(2-Cyclopentanecarbonyl-4-methyl-phenyl)-3-(5-guanidinomethyl-thiazol-2-yl)-urea
1-(2-Cyclopentanecarbonyl-4-methyl-phenyl)-3-(5-ureidomethyl-thiazol-2-yl)-urea
1-(2-Cyclopentanecarbonyl-4-methyl-phenyl)-3-{5-[3-(2-dimethylamino-ethyl)-ureidomethyl]-
thiazol-2-yl}-urea
{2-[3-(2-Cyclopentanecarbonyl-4-methyl-phenyl)-ureido]-thiazol-5-ylmethyl}-carbamic acid 2-
25 dimethylamino-ethyl ester
{2-[3-(2-Cyclopentanecarbonyl-4-methyl-phenyl)-ureido]-thiazol-5-ylmethyl}-carbamic acid
methyl ester
(2-{2-[3-(2-Cyclopentanecarbonyl-4-methyl-phenyl)-ureido]-thiazol-5-yl}-ethyl)-carbamic acid
methyl ester
30 1-(2-Cyclopentanecarbonyl-4-methyl-phenyl)-3-[5-(2-guanidino-ethyl)-thiazol-2-yl]-urea
1-(2-Cyclopentanecarbonyl-4-methyl-phenyl)-3-[5-(2-ureido-ethyl)-thiazol-2-yl]-urea
[3-(2-{2-[3-(2-Cyclopentanecarbonyl-4-methyl-phenyl)-ureido]-thiazol-5-yl}-ethyl)-ureido]-
acetic acid
1-(2-Cyclopentanecarbonyl-4-methyl-phenyl)-3-(5-{2-[3-(2-dimethylamino-ethyl)-ureido]-
35 ethyl}-thiazol-2-yl)-urea

- (2-{2-[3-(2-Cyclopentanecarbonyl-4-methyl-phenyl)-ureido]-thiazol-5-yl}-ethyl)-carbamic acid
2-dimethylamino-ethyl ester
- 1-(2-Cyclopentanecarbonyl-4-methyl-phenyl)-3-{5-[2-(1H-imidazol-2-ylsulfanyl)-ethyl]-thiazol-
2-yl}-urea
- 5 1-(2-Cyclopentanecarbonyl-4-methyl-phenyl)-3-{5-[2-(1H-imidazole-2-sulfonyl)-ethyl]-thiazol-
2-yl}-urea
- 1-(2-Cyclopentanecarbonyl-4-methyl-phenyl)-3-{5-[2-(1-methyl-1H-imidazol-2-ylsulfanyl)-
ethyl]-thiazol-2-yl}-urea
- 1-(2-Cyclopentanecarbonyl-4-methyl-phenyl)-3-{5-[2-(1-methyl-1H-imidazole-2-sulfonyl)-
ethyl]-thiazol-2-yl}-urea
- 10 2-(2-{2-[3-(2-Cyclopentanecarbonyl-4-methyl-phenyl)-ureido]-thiazol-5-yl}-ethylsulfanyl)-1H-
imidazole-4-carboxylic acid
- 6-{2-[3-(2-Cyclopentanecarbonyl-4-methyl-phenyl)-ureido]-thiazol-5-ylmethylsulfanyl}-
nicotinic acid
- 15 6-{2-[3-(2-Cyclopentanecarbonyl-4-methyl-phenyl)-ureido]-thiazol-5-ylmethanesulfonyl}-
nicotinic acid
- 2-(2-{2-[3-(2-Cyclopentanecarbonyl-4-methyl-phenyl)-ureido]-thiazol-5-yl}-ethylsulfanyl)-1-
methyl-1H-imidazole-4-carboxylic acid
- [4-(2-{2-[3-(2-Cyclopentanecarbonyl-4-methyl-phenyl)-ureido]-thiazol-5-yl}-ethylsulfanyl)-
phenyl]-acetic acid
- 20 [4-(2-{2-[3-(2-Cyclopentanecarbonyl-4-methyl-phenyl)-ureido]-thiazol-5-yl}-ethanesulfonyl)-
phenyl]-acetic acid
- 1-(2-Cyclopentanecarbonyl-4-methyl-phenyl)-3-{5-[2-(pyridin-2-ylsulfanyl)-ethyl]-thiazol-2-yl}-
urea
- 25 1-(2-Cyclopentanecarbonyl-4-methyl-phenyl)-3-{5-[2-(pyridine-2-sulfonyl)-ethyl]-thiazol-2-yl}-
urea
- 6-(2-{2-[3-(2-Cyclopentanecarbonyl-4-methyl-phenyl)-ureido]-thiazol-5-yl}-ethylsulfanyl)-
nicotinic acid
- 6-(2-{2-[3-(2-Cyclopentanecarbonyl-4-methyl-phenyl)-ureido]-thiazol-5-yl}-ethanesulfonyl)-
nicotinic acid
- 30 1-(2-Cyclopentanecarbonyl-4-methyl-phenyl)-3-(5-[1,2,4]oxadiazol-5-ylmethyl-thiazol-2-yl)-
urea
- 1-(2-Cyclopentanecarbonyl-4-methyl-phenyl)-3-[5-(2-[1,2,4]oxadiazol-5-yl-ethyl)-thiazol-2-yl]-
urea
- 35 4-{2-[3-(2-Cyclopentanecarbonyl-4-methyl-phenyl)-ureido]-thiazol-5-yl}-butyric acid

- 1-(2-Isobutoxy-4-methyl-phenyl)-3-(5-piperidin-1-ylmethyl-thiazol-2-yl)-urea
1-(2-Isobutoxy-4-methyl-phenyl)-3-(5-morpholin-4-ylmethyl-thiazol-2-yl)-urea
1-(5-Dimethylaminomethyl-thiazol-2-yl)-3-(2-isobutoxy-4-methyl-phenyl)-urea
1-(2-Isobutoxy-4-methyl-phenyl)-3-(5-piperazin-1-ylmethyl-thiazol-2-yl)-urea
5 1-(2-Isobutoxy-4-methyl-phenyl)-3-[5-(4-methyl-piperazin-1-ylmethyl)-thiazol-2-yl]-urea
1-[5-(4-Acetyl-piperazin-1-ylmethyl)-thiazol-2-yl]-3-(2-isobutoxy-4-methyl-phenyl)-urea
1-(2-Isobutoxy-4-methyl-phenyl)-3-[5-(4-methanesulfonyl-piperazin-1-ylmethyl)-thiazol-2-yl]-
urea
(4-{2-[3-(2-Isobutoxy-4-methyl-phenyl)-ureido]-thiazol-5-ylmethyl}-piperazin-1-yl)-acetic acid
10 2-{2-[3-(2-Isobutoxy-4-methyl-phenyl)-ureido]-thiazol-5-ylmethysulfanyl}-1H-imidazole-4-
carboxylic acid
1-[5-(1H-Imidazol-2-ylsulfanylmethyl)-thiazol-2-yl]-3-(2-isobutoxy-4-methyl-phenyl)-urea
1-(2-Isobutoxy-4-methyl-phenyl)-3-[5-(1-methyl-1H-imidazol-2-ylsulfanylmethyl)-thiazol-2-yl]-
urea
15 1-(2-Isobutoxy-4-methyl-phenyl)-3-[5-(4-methyl-4H-[1,2,4]triazol-3-ylsulfanylmethyl)-thiazol-2-
yl]-urea
1-(2-Isobutoxy-4-methyl-phenyl)-3-[5-(4H-[1,2,4]triazol-3-ylsulfanylmethyl)-thiazol-2-yl]-urea
1-(2-Isobutoxy-4-methyl-phenyl)-3-[5-(1-methyl-1H-tetrazol-5-ylsulfanylmethyl)-thiazol-2-yl]-
urea
20 (5-{2-[3-(2-Isobutoxy-4-methyl-phenyl)-ureido]-thiazol-5-ylmethysulfanyl}-tetrazol-1-yl)-acetic
acid
1-{5-[1-(2-Dimethylamino-ethyl)-1H-tetrazol-5-ylsulfanylmethyl]-thiazol-2-yl}-3-(2-isobutoxy-4-
methyl-phenyl)-urea
(5-{2-[3-(2-Isobutoxy-4-methyl-phenyl)-ureido]-thiazol-5-ylmethanesulfonyl}-tetrazol-1-yl)-
25 acetic acid
1-{5-[1-(2-Dimethylamino-ethyl)-1H-tetrazole-5-sulfonylmethyl]-thiazol-2-yl}-3-(2-isobutoxy-4-
methyl-phenyl)-urea
1-(2-Isobutoxy-4-methyl-phenyl)-3-[5-(1-methyl-1H-tetrazole-5-sulfonylmethyl)-thiazol-2-yl]-
urea
30 1-(2-Isobutoxy-4-methyl-phenyl)-3-[5-(4H-[1,2,4]triazole-3-sulfonylmethyl)-thiazol-2-yl]-urea
1-(2-Isobutoxy-4-methyl-phenyl)-3-[5-(4-methyl-4H-[1,2,4]triazole-3-sulfonylmethyl)-thiazol-2-
yl]-urea
1-(2-Isobutoxy-4-methyl-phenyl)-3-[5-(1-methyl-1H-imidazole-2-sulfonylmethyl)-thiazol-2-yl]-
urea
35 1-[5-(1H-Imidazole-2-sulfonylmethyl)-thiazol-2-yl]-3-(2-isobutoxy-4-methyl-phenyl)-urea

- 1-(2-Isobutoxy-4-methyl-phenyl)-3-[5-(pyridine-2-sulfonylmethyl)-thiazol-2-yl]-urea
1-(2-Isobutoxy-4-methyl-phenyl)-3-[5-(pyridin-2-ylsulfanylmethyl)-thiazol-2-yl]-urea
1-(2-Isobutoxy-4-methyl-phenyl)-3-[5-(pyridin-3-ylsulfanylmethyl)-thiazol-2-yl]-urea
1-(2-Isobutoxy-4-methyl-phenyl)-3-[5-(pyridine-3-sulfonylmethyl)-thiazol-2-yl]-urea
5 1-(2-Isobutoxy-4-methyl-phenyl)-3-[5-(pyridin-4-ylsulfanylmethyl)-thiazol-2-yl]-urea
1-(2-Isobutoxy-4-methyl-phenyl)-3-[5-(pyridine-4-sulfonylmethyl)-thiazol-2-yl]-urea
1-(2-Isobutoxy-4-methyl-phenyl)-3-[5-(pyridin-4-yloxymethyl)-thiazol-2-yl]-urea
1-(2-Isobutoxy-4-methyl-phenyl)-3-[5-(pyridin-3-yloxymethyl)-thiazol-2-yl]-urea
1-(2-Isobutoxy-4-methyl-phenyl)-3-[5-(pyridin-2-yloxymethyl)-thiazol-2-yl]-urea
10 (4-{2-[3-(2-Isobutoxy-4-methyl-phenyl)-ureido]-thiazol-5-ylmethoxy}-phenyl)-acetic acid
(4-{2-[3-(2-Isobutoxy-4-methyl-phenyl)-ureido]-thiazol-5-ylmethylsulfanyl}-phenyl)-acetic acid
(4-{2-[3-(2-Isobutoxy-4-methyl-phenyl)-ureido]-thiazol-5-ylmethanesulfonyl}-phenyl)-acetic acid
3-(4-{2-[3-(2-Isobutoxy-4-methyl-phenyl)-ureido]-thiazol-5-ylmethoxy}-phenyl)-propionic acid
15 3-(4-{2-[3-(2-Isobutoxy-4-methyl-phenyl)-ureido]-thiazol-5-ylmethylsulfanyl}-phenyl)-propionic acid
3-(4-{2-[3-(2-Isobutoxy-4-methyl-phenyl)-ureido]-thiazol-5-ylmethanesulfonyl}-phenyl)-propionic acid
4-{2-[3-(2-Isobutoxy-4-methyl-phenyl)-ureido]-thiazol-5-ylmethylsulfanyl}-benzoic acid
20 4-{2-[3-(2-Isobutoxy-4-methyl-phenyl)-ureido]-thiazol-5-ylmethanesulfonyl}-benzoic acid
4-{2-[3-(2-Isobutoxy-4-methyl-phenyl)-ureido]-thiazol-5-ylmethoxy}-benzoic acid
N-{2-[3-(2-Isobutoxy-4-methyl-phenyl)-ureido]-thiazol-5-ylmethyl}-acetamide
N-{2-[3-(2-Isobutoxy-4-methyl-phenyl)-ureido]-thiazol-5-ylmethyl}-methanesulfonamide
1-(2-Isobutoxy-4-methyl-phenyl)-3-[5-(pyridin-2-ylaminomethyl)-thiazol-2-yl]-urea
25 2-{2-[3-(2-Isobutoxy-4-methyl-phenyl)-ureido]-thiazol-5-ylmethanesulfonyl}-1H-imidazole-4-carboxylic acid
2-(2-{2-[3-(2-Isobutoxy-4-methyl-phenyl)-ureido]-thiazol-5-yl}-ethyl)-1H-imidazole-4-carboxylic acid
2-(2-{2-[3-(2-Isobutoxy-4-methyl-phenyl)-ureido]-thiazol-5-yl}-ethyl)-oxazole-4-carboxylic acid
30 2-(2-{2-[3-(2-Isobutoxy-4-methyl-phenyl)-ureido]-thiazol-5-yl}-ethyl)-1-methyl-1H-imidazole-4-carboxylic acid
2-[3-(2-Isobutoxy-4-methyl-phenyl)-ureido]-thiazole-5-carboxylic acid
2-[3-(2-Isobutoxy-4-methyl-phenyl)-ureido]-thiazole-5-carboxylic acid amide
2-[3-(2-Isobutoxy-4-methyl-phenyl)-ureido]-thiazole-5-carboxylic acid methylamide
35 ((2-[3-(2-Isobutoxy-4-methyl-phenyl)-ureido]-thiazole-5-carbonyl)-amino)-acetic acid

- ({2-[3-(2-Isobutoxy-4-methyl-phenyl)-ureido]-thiazole-5-carbonyl}-methanesulfonyl-amino)-acetic acid
N-{2-[3-(2-Isobutoxy-4-methyl-phenyl)-ureido]-thiazole-5-carbonyl}-methanesulfonamide
{2-[3-(2-Isobutoxy-4-methyl-phenyl)-ureido]-thiazol-5-yl}-acetic acid
5 2-{2-[3-(2-Isobutoxy-4-methyl-phenyl)-ureido]-thiazol-5-yl}-acetamide
2-{2-[3-(2-Isobutoxy-4-methyl-phenyl)-ureido]-thiazol-5-yl}-N-methyl-acetamide
(2-{2-[3-(2-Isobutoxy-4-methyl-phenyl)-ureido]-thiazol-5-yl}-acetyl-amino)-acetic acid
N-(2-{2-[3-(2-Isobutoxy-4-methyl-phenyl)-ureido]-thiazol-5-yl}-acetyl)-methanesulfonamide
N-(2-{2-[3-(2-Isobutoxy-4-methyl-phenyl)-ureido]-thiazol-5-yl}-ethyl)-methanesulfonamide
10 N-(2-{2-[3-(2-Isobutoxy-4-methyl-phenyl)-ureido]-thiazol-5-yl}-ethyl)-acetamide
1-(5-Guanidinomethyl-thiazol-2-yl)-3-(2-isobutoxy-4-methyl-phenyl)-urea
1-(2-Isobutoxy-4-methyl-phenyl)-3-(5-ureidomethyl-thiazol-2-yl)-urea
1-{5-[3-(2-Dimethylamino-ethyl)-ureidomethyl]-thiazol-2-yl}-3-(2-isobutoxy-4-methyl-phenyl)-urea
15 {2-[3-(2-Isobutoxy-4-methyl-phenyl)-ureido]-thiazol-5-ylmethyl}-carbamic acid 2-dimethylamino-ethyl ester
{2-[3-(2-Isobutoxy-4-methyl-phenyl)-ureido]-thiazol-5-ylmethyl}-carbamic acid methyl ester
(2-{2-[3-(2-Isobutoxy-4-methyl-phenyl)-ureido]-thiazol-5-yl}-ethyl)-carbamic acid methyl ester
1-[5-(2-Guanidino-ethyl)-thiazol-2-yl]-3-(2-isobutoxy-4-methyl-phenyl)-urea
20 1-(2-Isobutoxy-4-methyl-phenyl)-3-[5-(2-ureido-ethyl)-thiazol-2-yl]-urea
[3-(2-{2-[3-(2-Isobutoxy-4-methyl-phenyl)-ureido]-thiazol-5-yl}-ethyl)-ureido]-acetic acid
1-[5-(2-[3-(2-Dimethylamino-ethyl)-ureido]-ethyl)-thiazol-2-yl]-3-(2-isobutoxy-4-methyl-phenyl)-urea
(2-{2-[3-(2-Isobutoxy-4-methyl-phenyl)-ureido]-thiazol-5-yl}-ethyl)-carbamic acid 2-
25 dimethylamino-ethyl ester
1-[5-[2-(1H-Imidazol-2-ylsulfanyl)-ethyl]-thiazol-2-yl]-3-(2-isobutoxy-4-methyl-phenyl)-urea
1-[5-[2-(1H-Imidazole-2-sulfonyl)-ethyl]-thiazol-2-yl]-3-(2-isobutoxy-4-methyl-phenyl)-urea
1-(2-Isobutoxy-4-methyl-phenyl)-3-[5-[2-(1-methyl-1H-imidazol-2-ylsulfanyl)-ethyl]-thiazol-2-yl]-urea
30 1-(2-Isobutoxy-4-methyl-phenyl)-3-[5-[2-(1-methyl-1H-imidazole-2-sulfonyl)-ethyl]-thiazol-2-yl]-urea
2-(2-{2-[3-(2-Isobutoxy-4-methyl-phenyl)-ureido]-thiazol-5-yl}-ethylsulfanyl)-1H-imidazole-4-carboxylic acid
6-[2-[3-(2-Isobutoxy-4-methyl-phenyl)-ureido]-thiazol-5-ylmethylsulfanyl]-nicotinic acid
35 6-[2-[3-(2-Isobutoxy-4-methyl-phenyl)-ureido]-thiazol-5-ylmethanesulfonyl]-nicotinic acid

- 2-(2-{2-[3-(2-Isobutoxy-4-methyl-phenyl)-ureido]-thiazol-5-yl}-ethylsulfanyl)-1-methyl-1H-imidazole-4-carboxylic acid
- [4-(2-{2-[3-(2-Isobutoxy-4-methyl-phenyl)-ureido]-thiazol-5-yl}-ethylsulfanyl)-phenyl]-acetic acid
- 5 [4-(2-{2-[3-(2-Isobutoxy-4-methyl-phenyl)-ureido]-thiazol-5-yl}-ethanesulfonyl)-phenyl]-acetic acid
- 1-(2-Isobutoxy-4-methyl-phenyl)-3-{5-[2-(pyridin-2-ylsulfanyl)-ethyl]-thiazol-2-yl}-urea
- 1-(2-Isobutoxy-4-methyl-phenyl)-3-{5-[2-(pyridine-2-sulfonyl)-ethyl]-thiazol-2-yl}-urea
- 6-(2-{2-[3-(2-Isobutoxy-4-methyl-phenyl)-ureido]-thiazol-5-yl}-ethylsulfanyl)-nicotinic acid
- 10 6-(2-{2-[3-(2-Isobutoxy-4-methyl-phenyl)-ureido]-thiazol-5-yl}-ethanesulfonyl)-nicotinic acid
- 1-(2-Isobutoxy-4-methyl-phenyl)-3-(5-[1,2,4]oxadiazol-5-ylmethyl-thiazol-2-yl)-urea
- 1-(2-Isobutoxy-4-methyl-phenyl)-3-[5-(2-[1,2,4]oxadiazol-5-yl-ethyl)-thiazol-2-yl]-urea
- 4-(2-[3-(2-Isobutoxy-4-methyl-phenyl)-ureido]-thiazol-5-yl)-butyric acid
- 1-[5-(1H-Imidazol-2-ylsulfanyl)-thiazol-2-yl]-3-(2-isobutoxy-4-methyl-phenyl)-urea
- 15 1-(2-Isobutoxy-4-methyl-phenyl)-3-[5-(1-methyl-1H-imidazol-2-ylsulfanyl)-thiazol-2-yl]-urea
- 2-{2-[3-(2-Isobutoxy-4-methyl-phenyl)-ureido]-thiazol-5-ylsulfanyl}-1-methyl-1H-imidazole-4-carboxylic acid
- 2-{2-[3-(2-Isobutoxy-4-methyl-phenyl)-ureido]-thiazole-5-sulfonyl}-1-methyl-1H-imidazole-4-carboxylic acid
- 20 2-{2-[3-(2-Isobutoxy-4-methyl-phenyl)-ureido]-thiazole-5-sulfonyl}-1H-imidazole-4-carboxylic acid
- 1-[5-(1H-Imidazole-2-sulfonyl)-thiazol-2-yl]-3-(2-isobutoxy-4-methyl-phenyl)-urea
- 1-(2-Isobutoxy-4-methyl-phenyl)-3-[5-(pyridine-2-sulfonyl)-thiazol-2-yl]-urea
- 1-(2-Isobutoxy-4-methyl-phenyl)-3-[5-(pyridine-3-sulfonyl)-thiazol-2-yl]-urea
- 25 1-(2-Isobutoxy-4-methyl-phenyl)-3-[5-(pyridine-4-sulfonyl)-thiazol-2-yl]-urea
- 1-(2-Isobutoxy-4-methyl-phenyl)-3-[5-(pyridin-4-ylsulfanyl)-thiazol-2-yl]-urea
- 1-(2-Isobutoxy-4-methyl-phenyl)-3-[5-(pyridin-3-ylsulfanyl)-thiazol-2-yl]-urea
- 1-(2-Isobutoxy-4-methyl-phenyl)-3-[5-(pyridin-2-ylsulfanyl)-thiazol-2-yl]-urea
- 6-(2-[3-(2-Isobutoxy-4-methyl-phenyl)-ureido]-thiazol-5-ylsulfanyl)-nicotinic acid
- 30 6-(2-[3-(2-Isobutoxy-4-methyl-phenyl)-ureido]-thiazole-5-sulfonyl)-nicotinic acid
- 1-(2-Isobutoxy-4-methyl-phenyl)-3-(5-methanesulfonyl-thiazol-2-yl)-urea
- 3-(2-[3-(2-Isobutoxy-4-methyl-phenyl)-ureido]-thiazole-5-sulfonyl)-propionic acid
- 3-(2-[3-(2-Isobutoxy-4-methyl-phenyl)-ureido]-thiazol-5-ylsulfanyl)-propionic acid
- 1-(5-Bromo-thiazol-2-yl)-3-(2-isobutoxy-4-methyl-phenyl)-urea
- 35 {5-Chloro-2-[3-(2-isobutoxy-4-methyl-phenyl)-ureido]-thiazol-4-yl}-acetic acid

- {5-Bromo-2-[3-(2-isobutoxy-4-methyl-phenyl)-ureido]-thiazol-4-yl}-acetic acid
1-[2-(2-Methoxy-benzoyl)-4-methyl-phenyl]-3-(5-piperidin-1-ylmethyl-thiazol-2-yl)-urea
1-[2-(2-Methoxy-benzoyl)-4-methyl-phenyl]-3-(5-morpholin-4-ylmethyl-thiazol-2-yl)-urea
1-(5-Dimethylaminomethyl-thiazol-2-yl)-3-[2-(2-methoxy-benzoyl)-4-methyl-phenyl]-urea
5 1-[2-(2-Methoxy-benzoyl)-4-methyl-phenyl]-3-(5-piperazin-1-ylmethyl-thiazol-2-yl)-urea
1-[2-(2-Methoxy-benzoyl)-4-methyl-phenyl]-3-[5-(4-methyl-piperazin-1-ylmethyl)-thiazol-2-yl]-
urea
1-[5-(4-Acetyl-piperazin-1-ylmethyl)-thiazol-2-yl]-3-[2-(2-methoxy-benzoyl)-4-methyl-phenyl]-
urea
10 1-[5-(4-Methanesulfonyl-piperazin-1-ylmethyl)-thiazol-2-yl]-3-[2-(2-methoxy-benzoyl)-4-
methyl-phenyl]-urea
[4-(2-{3-[2-(2-Methoxy-benzoyl)-4-methyl-phenyl]-ureido)-thiazol-5-ylmethyl]-piperazin-1-yl]-
acetic acid
2-(2-{3-[2-(2-Methoxy-benzoyl)-4-methyl-phenyl]-ureido)-thiazol-5-ylmethylsulfanyl)-1H-
15 imidazole-4-carboxylic acid
1-[5-(1H-Imidazol-2-ylsulfanylmethyl)-thiazol-2-yl]-3-[2-(2-methoxy-benzoyl)-4-methyl-
phenyl]-urea
1-[2-(2-Methoxy-benzoyl)-4-methyl-phenyl]-3-[5-(1-methyl-1H-imidazol-2-ylsulfanylmethyl)-
thiazol-2-yl]-urea
20 1-[2-(2-Methoxy-benzoyl)-4-methyl-phenyl]-3-[5-(4-methyl-4H-[1,2,4]triazol-3-
ylsulfanylmethyl)-thiazol-2-yl]-urea
1-[2-(2-Methoxy-benzoyl)-4-methyl-phenyl]-3-[5-(4H-[1,2,4]triazol-3-ylsulfanylmethyl)-thiazol-
2-yl]-urea
1-[2-(2-Methoxy-benzoyl)-4-methyl-phenyl]-3-[5-(1-methyl-1H-tetrazol-5-ylsulfanylmethyl)-
25 thiazol-2-yl]-urea
[5-(2-{3-[2-(2-Methoxy-benzoyl)-4-methyl-phenyl]-ureido)-thiazol-5-ylmethylsulfanyl]-tetrazol-
1-yl]-acetic acid
1-[5-[1-(2-Dimethylamino-ethyl)-1H-tetrazol-5-ylsulfanylmethyl]-thiazol-2-yl]-3-[2-(2-methoxy-
benzoyl)-4-methyl-phenyl]-urea
30 [5-(2-{3-[2-(2-Methoxy-benzoyl)-4-methyl-phenyl]-ureido)-thiazol-5-ylmethanesulfonyl]-
tetrazol-1-yl]-acetic acid
1-[5-[1-(2-Dimethylamino-ethyl)-1H-tetrazole-5-sulfonylmethyl]-thiazol-2-yl]-3-[2-(2-methoxy-
benzoyl)-4-methyl-phenyl]-urea
1-[2-(2-Methoxy-benzoyl)-4-methyl-phenyl]-3-[5-(1-methyl-1H-tetrazole-5-sulfonylmethyl)-
35 thiazol-2-yl]-urea

- 1-[2-(2-Methoxy-benzoyl)-4-methyl-phenyl]-3-[5-(4H-[1,2,4]triazole-3-sulfonylmethyl)-thiazol-2-yl]-urea
- 1-[2-(2-Methoxy-benzoyl)-4-methyl-phenyl]-3-[5-(4-methyl-4H-[1,2,4]triazole-3-sulfonylmethyl)-thiazol-2-yl]-urea
- 5 1-[2-(2-Methoxy-benzoyl)-4-methyl-phenyl]-3-[5-(1-methyl-1H-imidazole-2-sulfonylmethyl)-thiazol-2-yl]-urea
- 1-[5-(1H-Imidazole-2-sulfonylmethyl)-thiazol-2-yl]-3-[2-(2-methoxy-benzoyl)-4-methyl-phenyl]-urea
- 1-[2-(2-Methoxy-benzoyl)-4-methyl-phenyl]-3-[5-(pyridine-2-sulfonylmethyl)-thiazol-2-yl]-urea
- 10 1-[2-(2-Methoxy-benzoyl)-4-methyl-phenyl]-3-[5-(pyridin-2-ylsulfanylmethyl)-thiazol-2-yl]-urea
- 1-[2-(2-Methoxy-benzoyl)-4-methyl-phenyl]-3-[5-(pyridin-3-ylsulfanylmethyl)-thiazol-2-yl]-urea
- 1-[2-(2-Methoxy-benzoyl)-4-methyl-phenyl]-3-[5-(pyridine-3-sulfonylmethyl)-thiazol-2-yl]-urea
- 1-[2-(2-Methoxy-benzoyl)-4-methyl-phenyl]-3-[5-(pyridin-4-ylsulfanylmethyl)-thiazol-2-yl]-urea
- 1-[2-(2-Methoxy-benzoyl)-4-methyl-phenyl]-3-[5-(pyridine-4-sulfonylmethyl)-thiazol-2-yl]-urea
- 15 1-[2-(2-Methoxy-benzoyl)-4-methyl-phenyl]-3-[5-(pyridin-4-yloxymethyl)-thiazol-2-yl]-urea
- 1-[2-(2-Methoxy-benzoyl)-4-methyl-phenyl]-3-[5-(pyridin-3-yloxymethyl)-thiazol-2-yl]-urea
- 1-[2-(2-Methoxy-benzoyl)-4-methyl-phenyl]-3-[5-(pyridin-2-yloxymethyl)-thiazol-2-yl]-urea
- [4-(2-{3-[2-(2-Methoxy-benzoyl)-4-methyl-phenyl]-ureido)-thiazol-5-ylmethoxy)-phenyl]-acetic acid
- 20 [4-(2-{3-[2-(2-Methoxy-benzoyl)-4-methyl-phenyl]-ureido)-thiazol-5-ylmethylsulfanyl)-phenyl]-acetic acid
- [4-(2-{3-[2-(2-Methoxy-benzoyl)-4-methyl-phenyl]-ureido)-thiazol-5-ylmethanesulfonyl)-phenyl]-acetic acid
- 3-[4-(2-{3-[2-(2-Methoxy-benzoyl)-4-methyl-phenyl]-ureido)-thiazol-5-ylmethoxy)-phenyl]-propionic acid
- 25 3-[4-(2-{3-[2-(2-Methoxy-benzoyl)-4-methyl-phenyl]-ureido)-thiazol-5-ylmethylsulfanyl)-phenyl]-propionic acid
- 3-[4-(2-{3-[2-(2-Methoxy-benzoyl)-4-methyl-phenyl]-ureido)-thiazol-5-ylmethanesulfonyl)-phenyl]-propionic acid
- 30 4-(2-{3-[2-(2-Methoxy-benzoyl)-4-methyl-phenyl]-ureido)-thiazol-5-ylmethylsulfanyl)-benzoic acid
- 4-(2-{3-[2-(2-Methoxy-benzoyl)-4-methyl-phenyl]-ureido)-thiazol-5-ylmethanesulfonyl)-benzoic acid
- 4-(2-{3-[2-(2-Methoxy-benzoyl)-4-methyl-phenyl]-ureido)-thiazol-5-ylmethoxy)-benzoic acid
- 35 N-(2-{3-[2-(2-Methoxy-benzoyl)-4-methyl-phenyl]-ureido)-thiazol-5-ylmethyl)-acetamide

- N-(2-{3-[2-(2-Methoxy-benzoyl)-4-methyl-phenyl]-ureido}-thiazol-5-ylmethyl)-methanesulfonamide
- 1-[2-(2-Methoxy-benzoyl)-4-methyl-phenyl]-3-[5-(pyridin-2-ylaminomethyl)-thiazol-2-yl]-urea
- 2-(2-{3-[2-(2-Methoxy-benzoyl)-4-methyl-phenyl]-ureido}-thiazol-5-ylmethanesulfonyl)-1H-
5 imidazole-4-carboxylic acid
- 2-[2-(2-{3-[2-(2-Methoxy-benzoyl)-4-methyl-phenyl]-ureido}-thiazol-5-yl)-ethyl]-1H-imidazole-4-carboxylic acid
- 2-[2-(2-{3-[2-(2-Methoxy-benzoyl)-4-methyl-phenyl]-ureido}-thiazol-5-yl)-ethyl]-oxazole-4-carboxylic acid
- 10 2-[2-(2-{3-[2-(2-Methoxy-benzoyl)-4-methyl-phenyl]-ureido}-thiazol-5-yl)-ethyl]-1-methyl-1H-imidazole-4-carboxylic acid
- 2-{3-[2-(2-Methoxy-benzoyl)-4-methyl-phenyl]-ureido}-thiazole-5-carboxylic acid
- 2-{3-[2-(2-Methoxy-benzoyl)-4-methyl-phenyl]-ureido}-thiazole-5-carboxylic acid amide
- 2-{3-[2-(2-Methoxy-benzoyl)-4-methyl-phenyl]-ureido}-thiazole-5-carboxylic acid methylamide
- 15 [(2-{3-[2-(2-Methoxy-benzoyl)-4-methyl-phenyl]-ureido}-thiazole-5-carbonyl)-amino]-acetic acid
- [Methanesulfonyl-(2-{3-[2-(2-methoxy-benzoyl)-4-methyl-phenyl]-ureido}-thiazole-5-carbonyl)-amino]-acetic acid
- N-(2-{3-[2-(2-Methoxy-benzoyl)-4-methyl-phenyl]-ureido}-thiazole-5-carbonyl)-
20 methanesulfonamide
- (2-{3-[2-(2-Methoxy-benzoyl)-4-methyl-phenyl]-ureido}-thiazol-5-yl)-acetic acid
- 2-(2-{3-[2-(2-Methoxy-benzoyl)-4-methyl-phenyl]-ureido}-thiazol-5-yl)-acetamide
- 2-(2-{3-[2-(2-Methoxy-benzoyl)-4-methyl-phenyl]-ureido}-thiazol-5-yl)-N-methyl-acetamide
- [2-(2-{3-[2-(2-Methoxy-benzoyl)-4-methyl-phenyl]-ureido}-thiazol-5-yl)-acetylamino]-acetic
25 acid
- N-[2-(2-{3-[2-(2-Methoxy-benzoyl)-4-methyl-phenyl]-ureido}-thiazol-5-yl)-acetyl]-methanesulfonamide
- N-[2-(2-{3-[2-(2-Methoxy-benzoyl)-4-methyl-phenyl]-ureido}-thiazol-5-yl)-ethyl]-methanesulfonamide
- 30 N-[2-(2-{3-[2-(2-Methoxy-benzoyl)-4-methyl-phenyl]-ureido}-thiazol-5-yl)-ethyl]-acetamide
- 1-(5-Guanidinomethyl-thiazol-2-yl)-3-[2-(2-methoxy-benzoyl)-4-methyl-phenyl]-urea
- 1-[2-(2-Methoxy-benzoyl)-4-methyl-phenyl]-3-(5-ureidomethyl-thiazol-2-yl)-urea
- 1-[5-[3-(2-Dimethylamino-ethyl)-ureidomethyl]-thiazol-2-yl]-3-[2-(2-methoxy-benzoyl)-4-methyl-phenyl]-urea

- (2-{3-[2-(2-Methoxy-benzoyl)-4-methyl-phenyl]-ureido}-thiazol-5-ylmethyl)-carbamic acid 2-dimethylamino-ethyl ester
- (2-{3-[2-(2-Methoxy-benzoyl)-4-methyl-phenyl]-ureido}-thiazol-5-ylmethyl)-carbamic acid methyl ester
- 5 [2-(2-{3-[2-(2-Methoxy-benzoyl)-4-methyl-phenyl]-ureido}-thiazol-5-yl)-ethyl]-carbamic acid methyl ester
- 1-[5-(2-Guanidino-ethyl)-thiazol-2-yl]-3-[2-(2-methoxy-benzoyl)-4-methyl-phenyl]-urea
- 1-[2-(2-Methoxy-benzoyl)-4-methyl-phenyl]-3-[5-(2-ureido-ethyl)-thiazol-2-yl]-urea
- {3-[2-(2-{3-[2-(2-Methoxy-benzoyl)-4-methyl-phenyl]-ureido}-thiazol-5-yl)-ethyl]-ureido}-acetic acid
- 10 acid
- 1-(5-{2-[3-(2-Dimethylamino-ethyl)-ureido]-ethyl}-thiazol-2-yl)-3-[2-(2-methoxy-benzoyl)-4-methyl-phenyl]-urea
- [2-(2-{3-[2-(2-Methoxy-benzoyl)-4-methyl-phenyl]-ureido}-thiazol-5-yl)-ethyl]-carbamic acid 2-dimethylamino-ethyl ester
- 15 1-{5-[2-(1H-Imidazol-2-ylsulfanyl)-ethyl]-thiazol-2-yl}-3-[2-(2-methoxy-benzoyl)-4-methyl-phenyl]-urea
- 1-{5-[2-(1H-Imidazole-2-sulfonyl)-ethyl]-thiazol-2-yl}-3-[2-(2-methoxy-benzoyl)-4-methyl-phenyl]-urea
- 1-[2-(2-Methoxy-benzoyl)-4-methyl-phenyl]-3-[5-[2-(1-methyl-1H-imidazol-2-ylsulfanyl)-ethyl]-thiazol-2-yl]-urea
- 20 thiazol-2-yl)-urea
- 1-[2-(2-Methoxy-benzoyl)-4-methyl-phenyl]-3-[5-[2-(1-methyl-1H-imidazole-2-sulfonyl)-ethyl]-thiazol-2-yl]-urea
- 2-[2-(2-{3-[2-(2-Methoxy-benzoyl)-4-methyl-phenyl]-ureido}-thiazol-5-yl)-ethylsulfanyl]-1H-imidazole-4-carboxylic acid
- 25 6-(2-{3-[2-(2-Methoxy-benzoyl)-4-methyl-phenyl]-ureido}-thiazol-5-ylmethylsulfanyl)-nicotinic acid
- 6-(2-{3-[2-(2-Methoxy-benzoyl)-4-methyl-phenyl]-ureido}-thiazol-5-ylmethanesulfonyl)-nicotinic acid
- 2-[2-(2-{3-[2-(2-Methoxy-benzoyl)-4-methyl-phenyl]-ureido}-thiazol-5-yl)-ethylsulfanyl]-1-methyl-1H-imidazole-4-carboxylic acid
- 30 methyl-1H-imidazole-4-carboxylic acid
- {4-[2-(2-{3-[2-(2-Methoxy-benzoyl)-4-methyl-phenyl]-ureido}-thiazol-5-yl)-ethylsulfanyl]-phenyl}-acetic acid
- {4-[2-(2-{3-[2-(2-Methoxy-benzoyl)-4-methyl-phenyl]-ureido}-thiazol-5-yl)-ethanesulfonyl]-phenyl}-acetic acid

- 1-[2-(2-Methoxy-benzoyl)-4-methyl-phenyl]-3-{5-[2-(pyridin-2-ylsulfanyl)-ethyl]-thiazol-2-yl}-
urea
- 1-[2-(2-Methoxy-benzoyl)-4-methyl-phenyl]-3-{5-[2-(pyridine-2-sulfonyl)-ethyl]-thiazol-2-yl}-
urea
- 5 6-[2-(2-{3-[2-(2-Methoxy-benzoyl)-4-methyl-phenyl]-ureido}-thiazol-5-yl)-ethylsulfanyl]-
nicotinic acid
- 6-[2-(2-{3-[2-(2-Methoxy-benzoyl)-4-methyl-phenyl]-ureido}-thiazol-5-yl)-ethanesulfonyl]-
nicotinic acid
- 1-[2-(2-Methoxy-benzoyl)-4-methyl-phenyl]-3-(5-[1,2,4]oxadiazol-5-ylmethyl-thiazol-2-yl)-urea
- 10 1-[2-(2-Methoxy-benzoyl)-4-methyl-phenyl]-3-[5-(2-[1,2,4]oxadiazol-5-yl-ethyl)-thiazol-2-yl]-
urea
- 4-(2-{3-[2-(2-Methoxy-benzoyl)-4-methyl-phenyl]-ureido}-thiazol-5-yl)-butyric acid
- 1-[5-(1H-Imidazol-2-ylsulfanyl)-thiazol-2-yl]-3-[2-(2-methoxy-benzoyl)-4-methyl-phenyl]-urea
- 1-[2-(2-Methoxy-benzoyl)-4-methyl-phenyl]-3-[5-(1-methyl-1H-imidazol-2-ylsulfanyl)-thiazol-2-
15 yl]-urea
- 2-(2-{3-[2-(2-Methoxy-benzoyl)-4-methyl-phenyl]-ureido}-thiazol-5-ylsulfanyl)-1-methyl-1H-
imidazole-4-carboxylic acid
- 2-(2-{3-[2-(2-Methoxy-benzoyl)-4-methyl-phenyl]-ureido}-thiazole-5-sulfonyl)-1-methyl-1H-
imidazole-4-carboxylic acid
- 20 2-(2-{3-[2-(2-Methoxy-benzoyl)-4-methyl-phenyl]-ureido}-thiazole-5-sulfonyl)-1H-imidazole-4-
carboxylic acid
- 1-[5-(1H-Imidazole-2-sulfonyl)-thiazol-2-yl]-3-[2-(2-methoxy-benzoyl)-4-methyl-phenyl]-urea
- 1-[2-(2-Methoxy-benzoyl)-4-methyl-phenyl]-3-[5-(pyridine-2-sulfonyl)-thiazol-2-yl]-urea
- 1-[2-(2-Methoxy-benzoyl)-4-methyl-phenyl]-3-[5-(pyridine-3-sulfonyl)-thiazol-2-yl]-urea
- 25 1-[2-(2-Methoxy-benzoyl)-4-methyl-phenyl]-3-[5-(pyridine-4-sulfonyl)-thiazol-2-yl]-urea
- 1-[2-(2-Methoxy-benzoyl)-4-methyl-phenyl]-3-[5-(pyridin-4-ylsulfanyl)-thiazol-2-yl]-urea
- 1-[2-(2-Methoxy-benzoyl)-4-methyl-phenyl]-3-[5-(pyridin-3-ylsulfanyl)-thiazol-2-yl]-urea
- 1-[2-(2-Methoxy-benzoyl)-4-methyl-phenyl]-3-[5-(pyridin-2-ylsulfanyl)-thiazol-2-yl]-urea
- 6-(2-{3-[2-(2-Methoxy-benzoyl)-4-methyl-phenyl]-ureido}-thiazol-5-ylsulfanyl)-nicotinic acid
- 30 6-(2-{3-[2-(2-Methoxy-benzoyl)-4-methyl-phenyl]-ureido}-thiazole-5-sulfonyl)-nicotinic acid
- 1-(5-Methanesulfonyl-thiazol-2-yl)-3-[2-(2-methoxy-benzoyl)-4-methyl-phenyl]-urea
- 3-(2-{3-[2-(2-Methoxy-benzoyl)-4-methyl-phenyl]-ureido}-thiazole-5-sulfonyl)-propionic acid
- 3-(2-{3-[2-(2-Methoxy-benzoyl)-4-methyl-phenyl]-ureido}-thiazol-5-ylsulfanyl)-propionic acid
- 1-(5-Bromo-thiazol-2-yl)-3-[2-(2-methoxy-benzoyl)-4-methyl-phenyl]-urea
- 35 (5-Chloro-2-{3-[2-(2-methoxy-benzoyl)-4-methyl-phenyl]-ureido}-thiazol-4-yl)-acetic acid

- (5-Bromo-2-{3-[2-(2-methoxy-benzoyl)-4-methyl-phenyl]-ureido}-thiazol-4-yl)-acetic acid
1-(5-Chloro-thiazol-2-yl)-3-[2-(2-methoxy-benzoyl)-4-methyl-phenyl]-urea
1-(5-Chloro-thiazol-2-yl)-3-(2-isobutoxy-4-methyl-phenyl)-urea
1-(4-Methyl-2-piperidin-1-yl-phenyl)-3-(5-piperidin-1-ylmethyl-thiazol-2-yl)-urea
5 1-(4-Methyl-2-piperidin-1-yl-phenyl)-3-(5-morpholin-4-ylmethyl-thiazol-2-yl)-urea
1-(5-Dimethylaminomethyl-thiazol-2-yl)-3-(4-methyl-2-piperidin-1-yl-phenyl)-urea
1-(4-Methyl-2-piperidin-1-yl-phenyl)-3-(5-piperazin-1-ylmethyl-thiazol-2-yl)-urea
1-[5-(4-Methyl-piperazin-1-ylmethyl)-thiazol-2-yl]-3-(4-methyl-2-piperidin-1-yl-phenyl)-urea
1-[5-(4-Acetyl-piperazin-1-ylmethyl)-thiazol-2-yl]-3-(4-methyl-2-piperidin-1-yl-phenyl)-urea
10 1-[5-(4-Methanesulfonyl-piperazin-1-ylmethyl)-thiazol-2-yl]-3-(4-methyl-2-piperidin-1-yl-phenyl)-urea
(4-{2-[3-(4-Methyl-2-piperidin-1-yl-phenyl)-ureido]-thiazol-5-ylmethyl}-piperazin-1-yl)-acetic acid
2-[2-[3-(4-Methyl-2-piperidin-1-yl-phenyl)-ureido]-thiazol-5-ylmethylsulfanyl]-1H-imidazole-4-
15 carboxylic acid
1-[5-(1H-Imidazol-2-ylsulfanylmethyl)-thiazol-2-yl]-3-(4-methyl-2-piperidin-1-yl-phenyl)-urea
1-[5-(1-Methyl-1H-imidazol-2-ylsulfanylmethyl)-thiazol-2-yl]-3-(4-methyl-2-piperidin-1-yl-phenyl)-urea
1-(4-Methyl-2-piperidin-1-yl-phenyl)-3-[5-(4-methyl-4H-[1,2,4]triazol-3-ylsulfanylmethyl)-
20 thiazol-2-yl]-urea
1-(4-Methyl-2-piperidin-1-yl-phenyl)-3-[5-(4H-[1,2,4]triazol-3-ylsulfanylmethyl)-thiazol-2-yl]-urea
1-(4-Methyl-2-piperidin-1-yl-phenyl)-3-[5-(1-methyl-1H-tetrazol-5-ylsulfanylmethyl)-thiazol-2-yl]-urea
25 (5-{2-[3-(4-Methyl-2-piperidin-1-yl-phenyl)-ureido]-thiazol-5-ylmethylsulfanyl}-tetrazol-1-yl)-acetic acid
1-[5-{1-(2-Dimethylamino-ethyl)-1H-tetrazol-5-ylsulfanylmethyl]-thiazol-2-yl]-3-(4-methyl-2-piperidin-1-yl-phenyl)-urea
(5-{2-[3-(4-Methyl-2-piperidin-1-yl-phenyl)-ureido]-thiazol-5-ylmethanesulfonyl}-tetrazol-1-yl)-
30 acetic acid
1-[5-{1-(2-Dimethylamino-ethyl)-1H-tetrazole-5-sulfonylmethyl]-thiazol-2-yl]-3-(4-methyl-2-piperidin-1-yl-phenyl)-urea
1-(4-Methyl-2-piperidin-1-yl-phenyl)-3-[5-(1-methyl-1H-tetrazole-5-sulfonylmethyl)-thiazol-2-yl]-urea

- 1-(4-Methyl-2-piperidin-1-yl-phenyl)-3-[5-(4H-[1,2,4]triazole-3-sulfonylmethyl)-thiazol-2-yl]-urea
- 1-(4-Methyl-2-piperidin-1-yl-phenyl)-3-[5-(4-methyl-4H-[1,2,4]triazole-3-sulfonylmethyl)-thiazol-2-yl]-urea
- 5 1-[5-(1-Methyl-1H-imidazole-2-sulfonylmethyl)-thiazol-2-yl]-3-(4-methyl-2-piperidin-1-yl-phenyl)-urea
- 1-[5-(1H-imidazole-2-sulfonylmethyl)-thiazol-2-yl]-3-(4-methyl-2-piperidin-1-yl-phenyl)-urea
- 1-(4-Methyl-2-piperidin-1-yl-phenyl)-3-[5-(pyridine-2-sulfonylmethyl)-thiazol-2-yl]-urea
- 1-(4-Methyl-2-piperidin-1-yl-phenyl)-3-[5-(pyridin-2-ylsulfanylmethyl)-thiazol-2-yl]-urea
- 10 1-(4-Methyl-2-piperidin-1-yl-phenyl)-3-[5-(pyridin-3-ylsulfanylmethyl)-thiazol-2-yl]-urea
- 1-(4-Methyl-2-piperidin-1-yl-phenyl)-3-[5-(pyridine-3-sulfonylmethyl)-thiazol-2-yl]-urea
- 1-(4-Methyl-2-piperidin-1-yl-phenyl)-3-[5-(pyridin-4-ylsulfanylmethyl)-thiazol-2-yl]-urea
- 1-(4-Methyl-2-piperidin-1-yl-phenyl)-3-[5-(pyridine-4-sulfonylmethyl)-thiazol-2-yl]-urea
- 1-(4-Methyl-2-piperidin-1-yl-phenyl)-3-[5-(pyridin-4-yloxymethyl)-thiazol-2-yl]-urea
- 15 1-(4-Methyl-2-piperidin-1-yl-phenyl)-3-[5-(pyridin-3-yloxymethyl)-thiazol-2-yl]-urea
- 1-(4-Methyl-2-piperidin-1-yl-phenyl)-3-[5-(pyridin-2-yloxymethyl)-thiazol-2-yl]-urea
- (4-{2-[3-(4-Methyl-2-piperidin-1-yl-phenyl)-ureido]-thiazol-5-ylmethoxy}-phenyl)-acetic acid
- (4-{2-[3-(4-Methyl-2-piperidin-1-yl-phenyl)-ureido]-thiazol-5-ylmethylsulfanyl}-phenyl)-acetic acid
- 20 (4-{2-[3-(4-Methyl-2-piperidin-1-yl-phenyl)-ureido]-thiazol-5-ylmethanesulfonyl}-phenyl)-acetic acid
- 3-(4-{2-[3-(4-Methyl-2-piperidin-1-yl-phenyl)-ureido]-thiazol-5-ylmethoxy}-phenyl)-propionic acid
- 3-(4-{2-[3-(4-Methyl-2-piperidin-1-yl-phenyl)-ureido]-thiazol-5-ylmethylsulfanyl}-phenyl)-propionic acid
- 25 3-(4-{2-[3-(4-Methyl-2-piperidin-1-yl-phenyl)-ureido]-thiazol-5-ylmethanesulfonyl}-phenyl)-propionic acid
- 4-{2-[3-(4-Methyl-2-piperidin-1-yl-phenyl)-ureido]-thiazol-5-ylmethylsulfanyl}-benzoic acid
- 4-{2-[3-(4-Methyl-2-piperidin-1-yl-phenyl)-ureido]-thiazol-5-ylmethanesulfonyl}-benzoic acid
- 30 4-{2-[3-(4-Methyl-2-piperidin-1-yl-phenyl)-ureido]-thiazol-5-ylmethoxy}-benzoic acid
- N-{2-[3-(4-Methyl-2-piperidin-1-yl-phenyl)-ureido]-thiazol-5-ylmethyl}-acetamide
- N-{2-[3-(4-Methyl-2-piperidin-1-yl-phenyl)-ureido]-thiazol-5-ylmethyl}-methanesulfonamide
- 1-(4-Methyl-2-piperidin-1-yl-phenyl)-3-[5-(pyridin-2-ylaminomethyl)-thiazol-2-yl]-urea
- 2-{2-[3-(4-Methyl-2-piperidin-1-yl-phenyl)-ureido]-thiazol-5-ylmethanesulfonyl}-1H-imidazole-
- 35 4-carboxylic acid

- 2-(2-{2-[3-(4-Methyl-2-piperidin-1-yl-phenyl)-ureido]-thiazol-5-yl}-ethyl)-1H-imidazole-4-carboxylic acid
- 2-(2-{2-[3-(4-Methyl-2-piperidin-1-yl-phenyl)-ureido]-thiazol-5-yl}-ethyl)-oxazole-4-carboxylic acid
- 5 1-Methyl-2-(2-{2-[3-(4-methyl-2-piperidin-1-yl-phenyl)-ureido]-thiazol-5-yl}-ethyl)-1H-imidazole-4-carboxylic acid
- 2-[3-(4-Methyl-2-piperidin-1-yl-phenyl)-ureido]-thiazole-5-carboxylic acid
- 2-[3-(4-Methyl-2-piperidin-1-yl-phenyl)-ureido]-thiazole-5-carboxylic acid amide
- 2-[3-(4-Methyl-2-piperidin-1-yl-phenyl)-ureido]-thiazole-5-carboxylic acid methylamide
- 10 ({2-[3-(4-Methyl-2-piperidin-1-yl-phenyl)-ureido]-thiazole-5-carbonyl}-amino)-acetic acid
- (Methanesulfonyl-{2-[3-(4-methyl-2-piperidin-1-yl-phenyl)-ureido]-thiazole-5-carbonyl}-amino)-acetic acid
- N-{2-[3-(4-Methyl-2-piperidin-1-yl-phenyl)-ureido]-thiazole-5-carbonyl}-methanesulfonamide
- {2-[3-(4-Methyl-2-piperidin-1-yl-phenyl)-ureido]-thiazol-5-yl}-acetic acid
- 15 2-{2-[3-(4-Methyl-2-piperidin-1-yl-phenyl)-ureido]-thiazol-5-yl}-acetamide
- N-Methyl-2-{2-[3-(4-methyl-2-piperidin-1-yl-phenyl)-ureido]-thiazol-5-yl}-acetamide
- (2-{2-[3-(4-Methyl-2-piperidin-1-yl-phenyl)-ureido]-thiazol-5-yl}-acetyl-amino)-acetic acid
- N-(2-{2-[3-(4-Methyl-2-piperidin-1-yl-phenyl)-ureido]-thiazol-5-yl}-acetyl)-methanesulfonamide
- N-(2-{2-[3-(4-Methyl-2-piperidin-1-yl-phenyl)-ureido]-thiazol-5-yl}-ethyl)-methanesulfonamide
- 20 N-(2-{2-[3-(4-Methyl-2-piperidin-1-yl-phenyl)-ureido]-thiazol-5-yl}-ethyl)-acetamide
- 1-(5-Guanidinomethyl-thiazol-2-yl)-3-(4-methyl-2-piperidin-1-yl-phenyl)-urea
- 1-(4-Methyl-2-piperidin-1-yl-phenyl)-3-(5-ureidomethyl-thiazol-2-yl)-urea
- 1-{5-[3-(2-Dimethylamino-ethyl)-ureidomethyl]-thiazol-2-yl}-3-(4-methyl-2-piperidin-1-yl-phenyl)-urea
- 25 {2-[3-(4-Methyl-2-piperidin-1-yl-phenyl)-ureido]-thiazol-5-ylmethyl}-carbamic acid 2-dimethylamino-ethyl ester
- {2-[3-(4-Methyl-2-piperidin-1-yl-phenyl)-ureido]-thiazol-5-ylmethyl}-carbamic acid methyl ester
- (2-{2-[3-(4-Methyl-2-piperidin-1-yl-phenyl)-ureido]-thiazol-5-yl}-ethyl)-carbamic acid methyl ester
- 30 1-[5-(2-Guanidino-ethyl)-thiazol-2-yl]-3-(4-methyl-2-piperidin-1-yl-phenyl)-urea
- 1-(4-Methyl-2-piperidin-1-yl-phenyl)-3-[5-(2-ureido-ethyl)-thiazol-2-yl]-urea
- [3-(2-{2-[3-(4-Methyl-2-piperidin-1-yl-phenyl)-ureido]-thiazol-5-yl}-ethyl)-ureido]-acetic acid
- 1-(5-{2-[3-(2-Dimethylamino-ethyl)-ureido]-ethyl}-thiazol-2-yl)-3-(4-methyl-2-piperidin-1-yl-phenyl)-urea

- (2-{2-[3-(4-Methyl-2-piperidin-1-yl-phenyl)-ureido]-thiazol-5-yl}-ethyl)-carbamic acid 2-dimethylamino-ethyl ester
- 1-{5-[2-(1H-Imidazol-2-ylsulfanyl)-ethyl]-thiazol-2-yl}-3-(4-methyl-2-piperidin-1-yl-phenyl)-urea
- 1-{5-[2-(1H-Imidazole-2-sulfonyl)-ethyl]-thiazol-2-yl}-3-(4-methyl-2-piperidin-1-yl-phenyl)-urea
- 5 1-{5-[2-(1-Methyl-1H-imidazol-2-ylsulfanyl)-ethyl]-thiazol-2-yl}-3-(4-methyl-2-piperidin-1-yl-phenyl)-urea
- 1-{5-[2-(1-Methyl-1H-imidazole-2-sulfonyl)-ethyl]-thiazol-2-yl}-3-(4-methyl-2-piperidin-1-yl-phenyl)-urea
- 2-(2-{2-[3-(4-Methyl-2-piperidin-1-yl-phenyl)-ureido]-thiazol-5-yl}-ethylsulfanyl)-1H-imidazole-
- 10 4-carboxylic acid
- 6-{2-[3-(4-Methyl-2-piperidin-1-yl-phenyl)-ureido]-thiazol-5-ylmethylsulfanyl}-nicotinic acid
- 6-{2-[3-(4-Methyl-2-piperidin-1-yl-phenyl)-ureido]-thiazol-5-ylmethanesulfonyl}-nicotinic acid
- 1-Methyl-2-(2-{2-[3-(4-methyl-2-piperidin-1-yl-phenyl)-ureido]-thiazol-5-yl}-ethylsulfanyl)-1H-imidazole-4-carboxylic acid
- 15 [4-(2-{2-[3-(4-Methyl-2-piperidin-1-yl-phenyl)-ureido]-thiazol-5-yl}-ethylsulfanyl)-phenyl]-acetic acid
- [4-(2-{2-[3-(4-Methyl-2-piperidin-1-yl-phenyl)-ureido]-thiazol-5-yl}-ethanesulfonyl)-phenyl]-acetic acid
- 1-(4-Methyl-2-piperidin-1-yl-phenyl)-3-{5-[2-(pyridin-2-ylsulfanyl)-ethyl]-thiazol-2-yl}-urea
- 20 1-(4-Methyl-2-piperidin-1-yl-phenyl)-3-{5-[2-(pyridine-2-sulfonyl)-ethyl]-thiazol-2-yl}-urea
- 6-(2-{2-[3-(4-Methyl-2-piperidin-1-yl-phenyl)-ureido]-thiazol-5-yl}-ethylsulfanyl)-nicotinic acid
- 6-(2-{2-[3-(4-Methyl-2-piperidin-1-yl-phenyl)-ureido]-thiazol-5-yl}-ethanesulfonyl)-nicotinic acid
- 1-(4-Methyl-2-piperidin-1-yl-phenyl)-3-(5-[1,2,4]oxadiazol-5-ylmethyl-thiazol-2-yl)-urea
- 25 1-(4-Methyl-2-piperidin-1-yl-phenyl)-3-[5-(2-[1,2,4]oxadiazol-5-yl-ethyl)-thiazol-2-yl]-urea
- 4-{2-[3-(4-Methyl-2-piperidin-1-yl-phenyl)-ureido]-thiazol-5-yl}-butyric acid
- 1-[5-(1H-Imidazol-2-ylsulfanyl)-thiazol-2-yl]-3-(4-methyl-2-piperidin-1-yl-phenyl)-urea
- 1-[5-(1-Methyl-1H-imidazol-2-ylsulfanyl)-thiazol-2-yl]-3-(4-methyl-2-piperidin-1-yl-phenyl)-urea
- 1-Methyl-2-{2-[3-(4-methyl-2-piperidin-1-yl-phenyl)-ureido]-thiazol-5-ylsulfanyl}-1H-imidazole-
- 30 4-carboxylic acid
- 1-Methyl-2-{2-[3-(4-methyl-2-piperidin-1-yl-phenyl)-ureido]-thiazole-5-sulfonyl}-1H-imidazole-4-carboxylic acid
- 2-{2-[3-(4-Methyl-2-piperidin-1-yl-phenyl)-ureido]-thiazole-5-sulfonyl}-1H-imidazole-4-carboxylic acid
- 35 1-[5-(1H-Imidazole-2-sulfonyl)-thiazol-2-yl]-3-(4-methyl-2-piperidin-1-yl-phenyl)-urea

- 1-(4-Methyl-2-piperidin-1-yl-phenyl)-3-[5-(pyridine-2-sulfonyl)-thiazol-2-yl]-urea
1-(4-Methyl-2-piperidin-1-yl-phenyl)-3-[5-(pyridine-3-sulfonyl)-thiazol-2-yl]-urea
1-(4-Methyl-2-piperidin-1-yl-phenyl)-3-[5-(pyridine-4-sulfonyl)-thiazol-2-yl]-urea
1-(4-Methyl-2-piperidin-1-yl-phenyl)-3-[5-(pyridin-4-ylsulfanyl)-thiazol-2-yl]-urea
5 1-(4-Methyl-2-piperidin-1-yl-phenyl)-3-[5-(pyridin-3-ylsulfanyl)-thiazol-2-yl]-urea
1-(4-Methyl-2-piperidin-1-yl-phenyl)-3-[5-(pyridin-2-ylsulfanyl)-thiazol-2-yl]-urea
6-{2-[3-(4-Methyl-2-piperidin-1-yl-phenyl)-ureido]-thiazol-5-ylsulfanyl}-nicotinic acid
6-{2-[3-(4-Methyl-2-piperidin-1-yl-phenyl)-ureido]-thiazole-5-sulfonyl}-nicotinic acid
1-(5-Methanesulfonyl-thiazol-2-yl)-3-(4-methyl-2-piperidin-1-yl-phenyl)-urea
10 3-{2-[3-(4-Methyl-2-piperidin-1-yl-phenyl)-ureido]-thiazole-5-sulfonyl}-propionic acid
3-{2-[3-(4-Methyl-2-piperidin-1-yl-phenyl)-ureido]-thiazol-5-ylsulfanyl}-propionic acid
1-(5-Bromo-thiazol-2-yl)-3-(4-methyl-2-piperidin-1-yl-phenyl)-urea
{5-Chloro-2-[3-(4-methyl-2-piperidin-1-yl-phenyl)-ureido]-thiazol-4-yl}-acetic acid
{5-Bromo-2-[3-(4-methyl-2-piperidin-1-yl-phenyl)-ureido]-thiazol-4-yl}-acetic acid
15 N-{2-[3-(4-Methyl-2-piperidin-1-yl-phenyl)-ureido]-thiazol-4-ylmethyl}-methanesulfonamide
N-{2-[2-[3-(4-Methyl-2-piperidin-1-yl-phenyl)-ureido]-thiazol-4-yl]-acetyl}-methanesulfonamide
1-(4-Dimethylaminomethyl-thiazol-2-yl)-3-(4-methyl-2-piperidin-1-yl-phenyl)-urea
1-(4-Methyl-2-piperidin-1-yl-phenyl)-3-(4-piperazin-1-ylmethyl-thiazol-2-yl)-urea
1-[4-(4-Methanesulfonyl-piperazin-1-ylmethyl)-thiazol-2-yl]-3-(4-methyl-2-piperidin-1-yl-
20 phenyl)-urea
1-(4-Methyl-2-piperidin-1-yl-phenyl)-3-(4-morpholin-4-ylmethyl-thiazol-2-yl)-urea
2-[3-(4-Methyl-2-piperidin-1-yl-phenyl)-ureido]-thiazole-4-carboxylic acid methylamide
1-(4-Methyl-2-piperidin-1-yl-phenyl)-3-[4-(morpholine-4-carbonyl)-thiazol-2-yl]-urea
{2-[3-(4-Methyl-2-piperidin-1-yl-phenyl)-ureido]-thiazol-4-ylmethyl}-carbamic acid methyl ester
25 1-(4-Methyl-2-piperidin-1-yl-phenyl)-3-(4-ureidomethyl-thiazol-2-yl)-urea
{2-[3-(4-Methyl-2-piperidin-1-yl-phenyl)-ureido]-thiazol-4-ylmethyl}-carbamic acid 2-
dimethylamino-ethyl ester
1-(4-Guanidinomethyl-thiazol-2-yl)-3-(4-methyl-2-piperidin-1-yl-phenyl)-urea
(3-{2-[3-(4-Methyl-2-piperidin-1-yl-phenyl)-ureido]-thiazol-4-ylmethyl}-ureido)-acetic acid
30 1-[4-[3-(2-Dimethylamino-ethyl)-ureidomethyl]-thiazol-2-yl]-3-(4-methyl-2-piperidin-1-yl-
phenyl)-urea
N-{2-[3-(2-Isobutoxy-4-methyl-phenyl)-ureido]-thiazol-4-ylmethyl}-methanesulfonamide
N-{2-[2-[3-(2-Isobutoxy-4-methyl-phenyl)-ureido]-thiazol-4-yl]-acetyl}-methanesulfonamide
1-(4-Dimethylaminomethyl-thiazol-2-yl)-3-(2-isobutoxy-4-methyl-phenyl)-urea
35 1-(2-Isobutoxy-4-methyl-phenyl)-3-(4-piperazin-1-ylmethyl-thiazol-2-yl)-urea

- 1-(2-Isobutoxy-4-methyl-phenyl)-3-[4-(4-methanesulfonyl-piperazin-1-ylmethyl)-thiazol-2-yl]-urea
- 1-(2-Isobutoxy-4-methyl-phenyl)-3-(4-morpholin-4-ylmethyl-thiazol-2-yl)-urea
- 2-[3-(2-Isobutoxy-4-methyl-phenyl)-ureido]-thiazole-4-carboxylic acid methylamide
- 5 1-(2-Isobutoxy-4-methyl-phenyl)-3-[4-(morpholine-4-carbonyl)-thiazol-2-yl]-urea
{2-[3-(2-Isobutoxy-4-methyl-phenyl)-ureido]-thiazol-4-ylmethyl}-carbamic acid methyl ester
1-(2-Isobutoxy-4-methyl-phenyl)-3-(4-ureidomethyl-thiazol-2-yl)-urea
{2-[3-(2-Isobutoxy-4-methyl-phenyl)-ureido]-thiazol-4-ylmethyl}-carbamic acid 2-dimethylamino-ethyl ester
- 10 1-(4-Guanidinomethyl-thiazol-2-yl)-3-(2-isobutoxy-4-methyl-phenyl)-urea
(3-{2-[3-(2-Isobutoxy-4-methyl-phenyl)-ureido]-thiazol-4-ylmethyl}-ureido)-acetic acid
1-[4-[3-(2-Dimethylamino-ethyl)-ureidomethyl]-thiazol-2-yl]-3-(2-isobutoxy-4-methyl-phenyl)-urea
N-(2-[3-[2-(2-Methoxy-benzoyl)-4-methyl-phenyl]-ureido]-thiazol-4-ylmethyl)-methanesulfonamide
- 15 N-[2-(2-[3-[2-(2-Methoxy-benzoyl)-4-methyl-phenyl]-ureido]-thiazol-4-yl)-acetyl]-methanesulfonamide
1-(4-Dimethylaminomethyl-thiazol-2-yl)-3-[2-(2-methoxy-benzoyl)-4-methyl-phenyl]-urea
1-[2-(2-Methoxy-benzoyl)-4-methyl-phenyl]-3-(4-piperazin-1-ylmethyl-thiazol-2-yl)-urea
- 20 1-[4-(4-Methanesulfonyl-piperazin-1-ylmethyl)-thiazol-2-yl]-3-[2-(2-methoxy-benzoyl)-4-methyl-phenyl]-urea
1-[2-(2-Methoxy-benzoyl)-4-methyl-phenyl]-3-(4-morpholin-4-ylmethyl-thiazol-2-yl)-urea
2-[3-[2-(2-Methoxy-benzoyl)-4-methyl-phenyl]-ureido]-thiazole-4-carboxylic acid methylamide
1-[2-(2-Methoxy-benzoyl)-4-methyl-phenyl]-3-[4-(morpholine-4-carbonyl)-thiazol-2-yl]-urea
- 25 (2-[3-[2-(2-Methoxy-benzoyl)-4-methyl-phenyl]-ureido]-thiazol-4-ylmethyl)-carbamic acid methyl ester
1-[2-(2-Methoxy-benzoyl)-4-methyl-phenyl]-3-(4-ureidomethyl-thiazol-2-yl)-urea
(2-[3-[2-(2-Methoxy-benzoyl)-4-methyl-phenyl]-ureido]-thiazol-4-ylmethyl)-carbamic acid 2-dimethylamino-ethyl ester
- 30 1-(4-Guanidinomethyl-thiazol-2-yl)-3-[2-(2-methoxy-benzoyl)-4-methyl-phenyl]-urea
[3-(2-[3-[2-(2-Methoxy-benzoyl)-4-methyl-phenyl]-ureido]-thiazol-4-ylmethyl)-ureido]-acetic acid
1-[4-[3-(2-Dimethylamino-ethyl)-ureidomethyl]-thiazol-2-yl]-3-[2-(2-methoxy-benzoyl)-4-methyl-phenyl]-urea

- 2-({2-[3-(2-Cyclopentanecarbonyl-4-methyl-phenyl)-ureido]-thiazole-5-carbonyl}-amino)-1H-imidazole-4-carboxylic acid
- 2-({2-[3-(2-Cyclopentanecarbonyl-4-methyl-phenyl)-ureido]-thiazole-5-carbonyl}-amino)-oxazole-4-carboxylic acid
- 5 2-({2-[3-(2-Cyclopentanecarbonyl-4-methyl-phenyl)-ureido]-thiazole-5-carbonyl}-amino)-1-methyl-1H-imidazole-4-carboxylic acid
- 2-({2-[3-(2-Cyclopentanecarbonyl-4-methyl-phenyl)-ureido]-thiazol-5-ylmethyl}-amino)-1H-imidazole-4-carboxylic acid
- 2-({2-[3-(2-Cyclopentanecarbonyl-4-methyl-phenyl)-ureido]-thiazol-5-ylmethyl}-amino)-oxazole-4-carboxylic acid
- 10 2-({2-[3-(2-Cyclopentanecarbonyl-4-methyl-phenyl)-ureido]-thiazol-5-ylmethyl}-amino)-1-methyl-1H-imidazole-4-carboxylic acid
- 6-({2-[3-(2-Cyclopentanecarbonyl-4-methyl-phenyl)-ureido]-thiazole-5-carbonyl}-amino)-nicotinic acid
- 15 2-({2-[3-(2-Cyclopentanecarbonyl-4-methyl-phenyl)-ureido]-thiazole-5-carbonyl}-amino)-pyrimidine-5-carboxylic acid
- 5-({2-[3-(2-Cyclopentanecarbonyl-4-methyl-phenyl)-ureido]-thiazole-5-carbonyl}-amino)-1-methyl-1H-pyrazole-3-carboxylic acid
- 2-(2-{2-[3-(2-Cyclopentanecarbonyl-4-methyl-phenyl)-ureido]-thiazol-5-yl}-ethyl)-5-methyl-1H-imidazole-4-carboxylic acid
- 20 2-(2-{2-[3-(2-Cyclopentanecarbonyl-4-methyl-phenyl)-ureido]-thiazol-5-yl}-ethyl)-5-trifluoromethyl-1H-imidazole-4-carboxylic acid
- 1-{5-[2-(1H-Benzimidazol-2-yl)-ethyl]-thiazol-2-yl}-3-(2-cyclopentanecarbonyl-4-methyl-phenyl)-urea
- 25 1-(2-Cyclopentanecarbonyl-4-methyl-phenyl)-3-{5-[2-(1H-imidazo[4,5-b]pyridin-2-yl)-ethyl]-thiazol-2-yl}-urea
- 2-{2-[3-(2-Cyclopentanecarbonyl-4-methyl-phenyl)-ureido]-thiazol-5-ylmethyl}-1H-imidazole-4-carboxylic acid
- 2-{2-[3-(2-Cyclopentanecarbonyl-4-methyl-phenyl)-ureido]-thiazol-5-ylmethyl}-1-methyl-1H-imidazole-4-carboxylic acid
- 30 2-{2-[3-(2-Cyclopentanecarbonyl-4-methyl-phenyl)-ureido]-thiazol-5-ylmethyl}-oxazole-4-carboxylic acid
- 1-[5-(1H-Benzimidazol-2-ylmethyl)-thiazol-2-yl]-3-(2-cyclopentanecarbonyl-4-methyl-phenyl)-urea

- 1-(2-Cyclopentanecarbonyl-4-methyl-phenyl)-3-[5-(1H-imidazo[4,5-b]pyridin-2-ylmethyl)-thiazol-2-yl]-urea
- 1-(5-Benzooxazol-2-ylmethyl-thiazol-2-yl)-3-(2-cyclopentanecarbonyl-4-methyl-phenyl)-urea
- 1-(2-Cyclopentanecarbonyl-4-methyl-phenyl)-3-(5-oxazolo[4,5-b]pyridin-2-ylmethyl-thiazol-2-yl)-urea
- 2-(2-{2-[3-(2-Cyclopentanecarbonyl-4-methyl-phenyl)-ureido]-thiazol-5-yl}-ethyl)-5-trifluoromethyl-oxazole-4-carboxylic acid
- 2-(2-{2-[3-(2-Cyclopentanecarbonyl-4-methyl-phenyl)-ureido]-thiazol-5-yl}-ethyl)-5-methyl-oxazole-4-carboxylic acid
- 1-(2-Cyclopentanecarbonyl-4-methyl-phenyl)-3-[5-[2-(4H-[1,2,4]triazol-3-yl)-ethyl]-thiazol-2-yl]-urea
- 1-(2-Cyclopentanecarbonyl-4-methyl-phenyl)-3-[5-[2-(4-methyl-4H-[1,2,4]triazol-3-yl)-ethyl]-thiazol-2-yl]-urea
- [3-(2-{2-[3-(2-Cyclopentanecarbonyl-4-methyl-phenyl)-ureido]-thiazol-5-yl}-ethyl)-[1,2,4]triazol-4-yl]-acetic acid
- [5-(2-{2-[3-(2-Cyclopentanecarbonyl-4-methyl-phenyl)-ureido]-thiazol-5-yl}-ethyl)-tetrazol-1-yl]-acetic acid
- 1-(2-Cyclopentanecarbonyl-4-methyl-phenyl)-3-[5-[2-(1-methyl-1H-tetrazol-5-yl)-ethyl]-thiazol-2-yl]-urea
- 1-(2-Cyclopentanecarbonyl-4-methyl-phenyl)-3-[5-[2-(1H-tetrazol-5-yl)-ethyl]-thiazol-2-yl]-urea
- 1-(2-Cyclopentanecarbonyl-4-methyl-phenyl)-3-[5-(2-[1,3,4]thiadiazol-2-yl-ethyl)-thiazol-2-yl]-urea
- 2-[3-(2-Cyclopentanecarbonyl-4-methyl-phenyl)-ureido]-thiazole-5-carboxylic acid
- [1,3,4]thiadiazol-2-ylamide
- 1-(2-Cyclopentanecarbonyl-4-methyl-phenyl)-3-[5-([1,3,4]thiadiazol-2-ylaminomethyl)-thiazol-2-yl]-urea
- [2-(2-{2-[3-(2-Cyclopentanecarbonyl-4-methyl-phenyl)-ureido]-thiazol-5-yl}-ethyl)-imidazol-1-yl]-acetic acid
- 2-[(2-{2-[3-(2-Methoxy-benzoyl)-4-methyl-phenyl]-ureido)-thiazole-5-carbonyl]-amino]-1H-imidazole-4-carboxylic acid
- 2-[(2-{2-[3-(2-Methoxy-benzoyl)-4-methyl-phenyl]-ureido)-thiazole-5-carbonyl]-amino]-oxazole-4-carboxylic acid
- 2-[(2-{2-[3-(2-Methoxy-benzoyl)-4-methyl-phenyl]-ureido)-thiazole-5-carbonyl]-amino]-1-methyl-1H-imidazole-4-carboxylic acid

- 2-[(2-{3-[2-(2-Methoxy-benzoyl)-4-methyl-phenyl]-ureido}-thiazol-5-ylmethyl)-amino]-1H-imidazole-4-carboxylic acid
- 2-[(2-{3-[2-(2-Methoxy-benzoyl)-4-methyl-phenyl]-ureido}-thiazol-5-ylmethyl)-amino]-oxazole-4-carboxylic acid
- 5 2-[(2-{3-[2-(2-Methoxy-benzoyl)-4-methyl-phenyl]-ureido}-thiazol-5-ylmethyl)-amino]-1-methyl-1H-imidazole-4-carboxylic acid
- 6-[(2-{3-[2-(2-Methoxy-benzoyl)-4-methyl-phenyl]-ureido}-thiazole-5-carbonyl)-amino]-nicotinic acid
- 2-[(2-{3-[2-(2-Methoxy-benzoyl)-4-methyl-phenyl]-ureido}-thiazole-5-carbonyl)-amino]-pyrimidine-5-carboxylic acid
- 10 5-[(2-{3-[2-(2-Methoxy-benzoyl)-4-methyl-phenyl]-ureido}-thiazole-5-carbonyl)-amino]-1-methyl-1H-pyrazole-3-carboxylic acid
- 2-[2-(2-{3-[2-(2-Methoxy-benzoyl)-4-methyl-phenyl]-ureido}-thiazol-5-yl)-ethyl]-5-methyl-1H-imidazole-4-carboxylic acid
- 15 2-[2-(2-{3-[2-(2-Methoxy-benzoyl)-4-methyl-phenyl]-ureido}-thiazol-5-yl)-ethyl]-5-trifluoromethyl-1H-imidazole-4-carboxylic acid
- 1-{5-[2-(1H-Benzimidazol-2-yl)-ethyl]-thiazol-2-yl}-3-[2-(2-methoxy-benzoyl)-4-methyl-phenyl]-urea
- 1-{5-[2-(1H-Imidazo[4,5-b]pyridin-2-yl)-ethyl]-thiazol-2-yl}-3-[2-(2-methoxy-benzoyl)-4-methyl-phenyl]-urea
- 20 2-(2-{3-[2-(2-Methoxy-benzoyl)-4-methyl-phenyl]-ureido}-thiazol-5-ylmethyl)-1H-imidazole-4-carboxylic acid
- 2-(2-{3-[2-(2-Methoxy-benzoyl)-4-methyl-phenyl]-ureido}-thiazol-5-ylmethyl)-1-methyl-1H-imidazole-4-carboxylic acid
- 25 2-(2-{3-[2-(2-Methoxy-benzoyl)-4-methyl-phenyl]-ureido}-thiazol-5-ylmethyl)-oxazole-4-carboxylic acid
- 1-[5-(1H-Benzimidazol-2-ylmethyl)-thiazol-2-yl]-3-[2-(2-methoxy-benzoyl)-4-methyl-phenyl]-urea
- 1-[5-(1H-Imidazo[4,5-b]pyridin-2-ylmethyl)-thiazol-2-yl]-3-[2-(2-methoxy-benzoyl)-4-methyl-phenyl]-urea
- 30 1-(5-Benzooxazol-2-ylmethyl-thiazol-2-yl)-3-[2-(2-methoxy-benzoyl)-4-methyl-phenyl]-urea
- 1-[2-(2-Methoxy-benzoyl)-4-methyl-phenyl]-3-(5-oxazolo[4,5-b]pyridin-2-ylmethyl-thiazol-2-yl)-urea
- 2-[2-(2-{3-[2-(2-Methoxy-benzoyl)-4-methyl-phenyl]-ureido}-thiazol-5-yl)-ethyl]-5-trifluoromethyl-oxazole-4-carboxylic acid
- 35

- 2-[2-(2-{3-[2-(2-Methoxy-benzoyl)-4-methyl-phenyl]-ureido]-thiazol-5-yl)-ethyl]-5-methyl-oxazole-4-carboxylic acid
- 1-[2-(2-Methoxy-benzoyl)-4-methyl-phenyl]-3-{5-[2-(4H-[1,2,4]triazol-3-yl)-ethyl]-thiazol-2-yl}-urea
- 5 1-[2-(2-Methoxy-benzoyl)-4-methyl-phenyl]-3-{5-[2-(4-methyl-4H-[1,2,4]triazol-3-yl)-ethyl]-thiazol-2-yl}-urea
- {3-[2-(2-{3-[2-(2-Methoxy-benzoyl)-4-methyl-phenyl]-ureido]-thiazol-5-yl)-ethyl]-[1,2,4]triazol-4-yl}-acetic acid
- {5-[2-(2-{3-[2-(2-Methoxy-benzoyl)-4-methyl-phenyl]-ureido]-thiazol-5-yl)-ethyl]-tetrazol-1-yl}-acetic acid
- 10 1-[2-(2-Methoxy-benzoyl)-4-methyl-phenyl]-3-{5-[2-(1-methyl-1H-tetrazol-5-yl)-ethyl]-thiazol-2-yl}-urea
- 1-[2-(2-Methoxy-benzoyl)-4-methyl-phenyl]-3-{5-[2-(1H-tetrazol-5-yl)-ethyl]-thiazol-2-yl}-urea
- 1-[2-(2-Methoxy-benzoyl)-4-methyl-phenyl]-3-{5-[2-[1,3,4]thiadiazol-2-yl-ethyl]-thiazol-2-yl}-urea
- 15 2-{3-[2-(2-Methoxy-benzoyl)-4-methyl-phenyl]-ureido]-thiazole-5-carboxylic acid
- [1,3,4]thiadiazol-2-ylamide
- 1-[2-(2-Methoxy-benzoyl)-4-methyl-phenyl]-3-{5-([1,3,4]thiadiazol-2-ylaminomethyl)-thiazol-2-yl}-urea
- 20 {2-[2-(2-{3-[2-(2-Methoxy-benzoyl)-4-methyl-phenyl]-ureido]-thiazol-5-yl)-ethyl]-imidazol-1-yl}-acetic acid
- 2-({2-[3-(2-Isobutoxy-4-methyl-phenyl)-ureido]-thiazole-5-carbonyl}-amino)-1H-imidazole-4-carboxylic acid
- 2-({2-[3-(2-Isobutoxy-4-methyl-phenyl)-ureido]-thiazole-5-carbonyl}-amino)-oxazole-4-carboxylic acid
- 25 2-({2-[3-(2-Isobutoxy-4-methyl-phenyl)-ureido]-thiazole-5-carbonyl}-amino)-1-methyl-1H-imidazole-4-carboxylic acid
- 2-({2-[3-(2-Isobutoxy-4-methyl-phenyl)-ureido]-thiazol-5-ylmethyl}-amino)-1H-imidazole-4-carboxylic acid
- 30 2-({2-[3-(2-Isobutoxy-4-methyl-phenyl)-ureido]-thiazol-5-ylmethyl}-amino)-oxazole-4-carboxylic acid
- 2-({2-[3-(2-Isobutoxy-4-methyl-phenyl)-ureido]-thiazol-5-ylmethyl}-amino)-1-methyl-1H-imidazole-4-carboxylic acid
- 6-({2-[3-(2-Isobutoxy-4-methyl-phenyl)-ureido]-thiazole-5-carbonyl}-amino)-nicotinic acid

- 2-({2-[3-(2-Isobutoxy-4-methyl-phenyl)-ureido]-thiazole-5-carbonyl}-amino)-pyrimidine-5-carboxylic acid
- 5-({2-[3-(2-Isobutoxy-4-methyl-phenyl)-ureido]-thiazole-5-carbonyl}-amino)-1-methyl-1H-pyrazole-3-carboxylic acid
- 5 2-(2-{2-[3-(2-Isobutoxy-4-methyl-phenyl)-ureido]-thiazol-5-yl}-ethyl)-5-methyl-1H-imidazole-4-carboxylic acid
- 2-(2-{2-[3-(2-Isobutoxy-4-methyl-phenyl)-ureido]-thiazol-5-yl}-ethyl)-5-trifluoromethyl-1H-imidazole-4-carboxylic acid
- 1-{5-[2-(1H-Benzimidazol-2-yl)-ethyl]-thiazol-2-yl}-3-(2-isobutoxy-4-methyl-phenyl)-urea
- 10 1-{5-[2-(1H-Imidazo[4,5-b]pyridin-2-yl)-ethyl]-thiazol-2-yl}-3-(2-isobutoxy-4-methyl-phenyl)-urea
- 2-{2-[3-(2-Isobutoxy-4-methyl-phenyl)-ureido]-thiazol-5-ylmethyl}-1H-imidazole-4-carboxylic acid
- 2-{2-[3-(2-Isobutoxy-4-methyl-phenyl)-ureido]-thiazol-5-ylmethyl}-1-methyl-1H-imidazole-4-carboxylic acid
- 15 2-{2-[3-(2-Isobutoxy-4-methyl-phenyl)-ureido]-thiazol-5-ylmethyl}-oxazole-4-carboxylic acid
- 1-[5-(1H-Benzimidazol-2-ylmethyl)-thiazol-2-yl]-3-(2-isobutoxy-4-methyl-phenyl)-urea
- 1-[5-(1H-Imidazo[4,5-b]pyridin-2-ylmethyl)-thiazol-2-yl]-3-(2-isobutoxy-4-methyl-phenyl)-urea
- 20 1-(5-Benzooxazol-2-ylmethyl-thiazol-2-yl)-3-(2-isobutoxy-4-methyl-phenyl)-urea
- 1-(2-Isobutoxy-4-methyl-phenyl)-3-(5-oxazolo[4,5-b]pyridin-2-ylmethyl-thiazol-2-yl)-urea
- 2-(2-{2-[3-(2-Isobutoxy-4-methyl-phenyl)-ureido]-thiazol-5-yl}-ethyl)-5-trifluoromethyl-oxazole-4-carboxylic acid
- 2-(2-{2-[3-(2-Isobutoxy-4-methyl-phenyl)-ureido]-thiazol-5-yl}-ethyl)-5-methyl-oxazole-4-carboxylic acid
- 25 1-(2-Isobutoxy-4-methyl-phenyl)-3-{5-[2-(4H-[1,2,4]triazol-3-yl)-ethyl]-thiazol-2-yl}-urea
- 1-(2-Isobutoxy-4-methyl-phenyl)-3-{5-[2-(4-methyl-4H-[1,2,4]triazol-3-yl)-ethyl]-thiazol-2-yl}-urea
- [3-(2-{2-[3-(2-Isobutoxy-4-methyl-phenyl)-ureido]-thiazol-5-yl}-ethyl)-[1,2,4]triazol-4-yl]-acetic acid
- 30 5-(2-{2-[3-(2-Isobutoxy-4-methyl-phenyl)-ureido]-thiazol-5-yl}-ethyl)-tetrazol-1-yl]-acetic acid
- 1-(2-Isobutoxy-4-methyl-phenyl)-3-{5-[2-(1-methyl-1H-tetrazol-5-yl)-ethyl]-thiazol-2-yl}-urea
- 1-(2-Isobutoxy-4-methyl-phenyl)-3-{5-[2-(1H-tetrazol-5-yl)-ethyl]-thiazol-2-yl}-urea
- 1-(2-Isobutoxy-4-methyl-phenyl)-3-{5-[2-[1,3,4]thiadiazol-2-yl)-ethyl]-thiazol-2-yl}-urea

- 2-[3-(2-Isobutoxy-4-methyl-phenyl)-ureido]-thiazole-5-carboxylic acid [1,3,4]thiadiazol-2-ylamide
- 1-(2-Isobutoxy-4-methyl-phenyl)-3-[5-([1,3,4]thiadiazol-2-ylaminomethyl)-thiazol-2-yl]-urea
- [2-(2-[3-(2-Isobutoxy-4-methyl-phenyl)-ureido]-thiazol-5-yl)-ethyl]-imidazol-1-yl]-acetic acid
- 5 2-((2-[3-(4-Methyl-2-piperidin-1-yl-phenyl)-ureido]-thiazole-5-carbonyl)-amino)-1H-imidazole-4-carboxylic acid
- 2-((2-[3-(4-Methyl-2-piperidin-1-yl-phenyl)-ureido]-thiazole-5-carbonyl)-amino)-oxazole-4-carboxylic acid
- 1-Methyl-2-((2-[3-(4-methyl-2-piperidin-1-yl-phenyl)-ureido]-thiazole-5-carbonyl)-amino)-1H-
- 10 imidazole-4-carboxylic acid
- 2-((2-[3-(4-Methyl-2-piperidin-1-yl-phenyl)-ureido]-thiazol-5-ylmethyl)-amino)-1H-imidazole-4-carboxylic acid
- 2-((2-[3-(4-Methyl-2-piperidin-1-yl-phenyl)-ureido]-thiazol-5-ylmethyl)-amino)-oxazole-4-carboxylic acid
- 15 1-Methyl-2-((2-[3-(4-methyl-2-piperidin-1-yl-phenyl)-ureido]-thiazol-5-ylmethyl)-amino)-1H-imidazole-4-carboxylic acid
- 6-((2-[3-(4-Methyl-2-piperidin-1-yl-phenyl)-ureido]-thiazole-5-carbonyl)-amino)-nicotinic acid
- 2-((2-[3-(4-Methyl-2-piperidin-1-yl-phenyl)-ureido]-thiazole-5-carbonyl)-amino)-pyrimidine-5-carboxylic acid
- 20 1-Methyl-5-((2-[3-(4-methyl-2-piperidin-1-yl-phenyl)-ureido]-thiazole-5-carbonyl)-amino)-1H-pyrazole-3-carboxylic acid
- 5-Methyl-2-(2-[3-(4-methyl-2-piperidin-1-yl-phenyl)-ureido]-thiazol-5-yl)-ethyl)-1H-imidazole-4-carboxylic acid
- 2-(2-[3-(4-Methyl-2-piperidin-1-yl-phenyl)-ureido]-thiazol-5-yl)-ethyl)-5-trifluoromethyl-1H-
- 25 imidazole-4-carboxylic acid
- 1-[5-[2-(1H-Benzoimidazol-2-yl)-ethyl]-thiazol-2-yl]-3-(4-methyl-2-piperidin-1-yl-phenyl)-urea
- 1-[5-[2-(1H-Imidazo[4,5-b]pyridin-2-yl)-ethyl]-thiazol-2-yl]-3-(4-methyl-2-piperidin-1-yl-phenyl)-urea
- 2-[2-[3-(4-Methyl-2-piperidin-1-yl-phenyl)-ureido]-thiazol-5-ylmethyl]-1H-imidazole-4-
- 30 carboxylic acid
- 1-Methyl-2-[2-[3-(4-methyl-2-piperidin-1-yl-phenyl)-ureido]-thiazol-5-ylmethyl]-1H-imidazole-4-carboxylic acid
- 2-[2-[3-(4-Methyl-2-piperidin-1-yl-phenyl)-ureido]-thiazol-5-ylmethyl]-oxazole-4-carboxylic acid
- 35 1-[5-(1H-Benzoimidazol-2-ylmethyl)-thiazol-2-yl]-3-(4-methyl-2-piperidin-1-yl-phenyl)-urea

- 1-[5-(1H-Imidazo[4,5-#b]pyridin-2-ylmethyl)-thiazol-2-yl]-3-(4-methyl-2-piperidin-1-yl-phenyl)-urea
- 1-(5-Benzooxazol-2-ylmethyl-thiazol-2-yl)-3-(4-methyl-2-piperidin-1-yl-phenyl)-urea
- 1-(4-Methyl-2-piperidin-1-yl-phenyl)-3-(5-oxazolo[4,5-#b]pyridin-2-ylmethyl-thiazol-2-yl)-urea
- 5 2-(2-{2-[3-(4-Methyl-2-piperidin-1-yl-phenyl)-ureido]-thiazol-5-yl}-ethyl)-5-trifluoromethyl-oxazole-4-carboxylic acid
- 5-Methyl-2-(2-{2-[3-(4-methyl-2-piperidin-1-yl-phenyl)-ureido]-thiazol-5-yl}-ethyl)-oxazole-4-carboxylic acid
- 1-(4-Methyl-2-piperidin-1-yl-phenyl)-3-{5-[2-(4H-[1,2,4]triazol-3-yl)-ethyl]-thiazol-2-yl}-urea
- 10 1-(4-Methyl-2-piperidin-1-yl-phenyl)-3-{5-[2-(4-methyl-4H-[1,2,4]triazol-3-yl)-ethyl]-thiazol-2-yl}-urea
- [3-(2-{2-[3-(4-Methyl-2-piperidin-1-yl-phenyl)-ureido]-thiazol-5-yl}-ethyl)-[1,2,4]triazol-4-yl]-acetic acid
- [5-(2-{2-[3-(4-Methyl-2-piperidin-1-yl-phenyl)-ureido]-thiazol-5-yl}-ethyl)-tetrazol-1-yl]-acetic acid
- 15 acid
- 1-(4-Methyl-2-piperidin-1-yl-phenyl)-3-{5-[2-(1-methyl-1H-tetrazol-5-yl)-ethyl]-thiazol-2-yl}-urea
- 1-(4-Methyl-2-piperidin-1-yl-phenyl)-3-{5-[2-(1H-tetrazol-5-yl)-ethyl]-thiazol-2-yl}-urea
- 1-(4-Methyl-2-piperidin-1-yl-phenyl)-3-{5-[2-[1,3,4]thiadiazol-2-yl-ethyl]-thiazol-2-yl}-urea
- 20 2-[3-(4-Methyl-2-piperidin-1-yl-phenyl)-ureido]-thiazole-5-carboxylic acid [1,3,4]thiadiazol-2-ylamide
- 1-(4-Methyl-2-piperidin-1-yl-phenyl)-3-{5-[1,3,4]thiadiazol-2-ylaminomethyl)-thiazol-2-yl}-urea
- 1-(2-Cyclopentanecarbonyl-phenyl)-3-[1,2,4]thiadiazol-5-yl-urea
- {5-[3-(2-Cyclopentanecarbonyl-phenyl)-ureido]-[1,2,4]thiadiazol-3-yl}-acetic acid
- 25 N-{5-[3-(2-Cyclopentanecarbonyl-phenyl)-ureido]-[1,2,4]thiadiazol-3-ylmethyl}-methanesulfonamide
- 1-(2-Cyclopentanecarbonyl-phenyl)-3-(3-methoxy-[1,2,4]thiadiazol-5-yl)-urea
- 1-(2-Cyclopentanecarbonyl-phenyl)-3-(3-ethyl-[1,2,4]thiadiazol-5-yl)-urea
- 1-(2-Cyclopentanecarbonyl-phenyl)-3-(3-methyl-[1,2,4]thiadiazol-5-yl)-urea
- 30 1-(2-Cyclopentanecarbonyl-phenyl)-3-(3-isopropyl-[1,2,4]thiadiazol-5-yl)-urea
- 1-(2-Cyclopentanecarbonyl-phenyl)-3-(3-phenyl-[1,2,4]thiadiazol-5-yl)-urea
- 1-(2-Isobutoxy-phenyl)-3-[1,2,4]thiadiazol-5-yl-urea
- {5-[3-(2-Isobutoxy-phenyl)-ureido]-[1,2,4]thiadiazol-3-yl}-acetic acid
- N-{5-[3-(2-Isobutoxy-phenyl)-ureido]-[1,2,4]thiadiazol-3-ylmethyl}-methanesulfonamide
- 35 1-(2-Isobutoxy-phenyl)-3-(3-methoxy-[1,2,4]thiadiazol-5-yl)-urea

- 1-(3-Ethyl-[1,2,4]thiadiazol-5-yl)-3-(2-isobutoxy-phenyl)-urea
1-(2-Isobutoxy-phenyl)-3-(3-methyl-[1,2,4]thiadiazol-5-yl)-urea
1-(2-Isobutoxy-phenyl)-3-(3-isopropyl-[1,2,4]thiadiazol-5-yl)-urea
1-(2-Isobutoxy-phenyl)-3-(3-phenyl-[1,2,4]thiadiazol-5-yl)-urea
5 1-[2-(2-Methoxy-benzoyl)-phenyl]-3-[1,2,4]thiadiazol-5-yl-urea
(5-{3-[2-(2-Methoxy-benzoyl)-phenyl]-ureido}-[1,2,4]thiadiazol-3-yl)-acetic acid
N-(5-{3-[2-(2-Methoxy-benzoyl)-phenyl]-ureido}-[1,2,4]thiadiazol-3-ylmethyl)-methanesulfonamide
1-[2-(2-Methoxy-benzoyl)-phenyl]-3-(3-methoxy-[1,2,4]thiadiazol-5-yl)-urea
10 1-(3-Ethyl-[1,2,4]thiadiazol-5-yl)-3-[2-(2-methoxy-benzoyl)-phenyl]-urea
1-[2-(2-Methoxy-benzoyl)-phenyl]-3-(3-methyl-[1,2,4]thiadiazol-5-yl)-urea
1-(3-Isopropyl-[1,2,4]thiadiazol-5-yl)-3-[2-(2-methoxy-benzoyl)-phenyl]-urea
1-[2-(2-Methoxy-benzoyl)-phenyl]-3-(3-phenyl-[1,2,4]thiadiazol-5-yl)-urea
1-(2-Piperidin-1-yl-phenyl)-3-[1,2,4]thiadiazol-5-yl-urea
15 {5-[3-(2-Piperidin-1-yl-phenyl)-ureido]-[1,2,4]thiadiazol-3-yl}-acetic acid
N-{5-[3-(2-Piperidin-1-yl-phenyl)-ureido]-[1,2,4]thiadiazol-3-ylmethyl}-methanesulfonamide
1-(3-Methoxy-[1,2,4]thiadiazol-5-yl)-3-(2-piperidin-1-yl-phenyl)-urea
1-(3-Ethyl-[1,2,4]thiadiazol-5-yl)-3-(2-piperidin-1-yl-phenyl)-urea
1-(3-Methyl-[1,2,4]thiadiazol-5-yl)-3-(2-piperidin-1-yl-phenyl)-urea
20 1-(3-Isopropyl-[1,2,4]thiadiazol-5-yl)-3-(2-piperidin-1-yl-phenyl)-urea
1-(3-Phenyl-[1,2,4]thiadiazol-5-yl)-3-(2-piperidin-1-yl-phenyl)-urea
1-(2-Cyclopentanecarbonyl-4-methyl-phenyl)-3-[1,2,4]thiadiazol-5-yl-urea
{5-[3-(2-Cyclopentanecarbonyl-4-methyl-phenyl)-ureido]-[1,2,4]thiadiazol-3-yl}-acetic acid
N-{5-[3-(2-Cyclopentanecarbonyl-4-methyl-phenyl)-ureido]-[1,2,4]thiadiazol-3-ylmethyl}-methanesulfonamide
25 1-(2-Cyclopentanecarbonyl-4-methyl-phenyl)-3-(3-methoxy-[1,2,4]thiadiazol-5-yl)-urea
1-(2-Cyclopentanecarbonyl-4-methyl-phenyl)-3-(3-ethyl-[1,2,4]thiadiazol-5-yl)-urea
1-(2-Cyclopentanecarbonyl-4-methyl-phenyl)-3-(3-methyl-[1,2,4]thiadiazol-5-yl)-urea
1-(2-Cyclopentanecarbonyl-4-methyl-phenyl)-3-(3-isopropyl-[1,2,4]thiadiazol-5-yl)-urea
30 1-(2-Cyclopentanecarbonyl-4-methyl-phenyl)-3-(3-phenyl-[1,2,4]thiadiazol-5-yl)-urea
1-(2-Isobutoxy-4-methyl-phenyl)-3-[1,2,4]thiadiazol-5-yl-urea
{5-[3-(2-Isobutoxy-4-methyl-phenyl)-ureido]-[1,2,4]thiadiazol-3-yl}-acetic acid
N-{5-[3-(2-Isobutoxy-4-methyl-phenyl)-ureido]-[1,2,4]thiadiazol-3-ylmethyl}-methanesulfonamide
35 1-(2-Isobutoxy-4-methyl-phenyl)-3-(3-methoxy-[1,2,4]thiadiazol-5-yl)-urea

- 1-(3-Ethyl-[1,2,4]thiadiazol-5-yl)-3-(2-isobutoxy-4-methyl-phenyl)-urea
1-(2-Isobutoxy-4-methyl-phenyl)-3-(3-methyl-[1,2,4]thiadiazol-5-yl)-urea
1-(2-Isobutoxy-4-methyl-phenyl)-3-(3-isopropyl-[1,2,4]thiadiazol-5-yl)-urea
1-(2-Isobutoxy-4-methyl-phenyl)-3-(3-phenyl-[1,2,4]thiadiazol-5-yl)-urea
5 1-[2-(2-Methoxy-benzoyl)-4-methyl-phenyl]-3-[1,2,4]thiadiazol-5-yl-urea
(5-{3-[2-(2-Methoxy-benzoyl)-4-methyl-phenyl]-ureido}-[1,2,4]thiadiazol-3-yl)-acetic acid
N-(5-{3-[2-(2-Methoxy-benzoyl)-4-methyl-phenyl]-ureido}-[1,2,4]thiadiazol-3-ylmethyl)-
methanesulfonamide
1-[2-(2-Methoxy-benzoyl)-4-methyl-phenyl]-3-(3-methoxy-[1,2,4]thiadiazol-5-yl)-urea
10 1-(3-Ethyl-[1,2,4]thiadiazol-5-yl)-3-[2-(2-methoxy-benzoyl)-4-methyl-phenyl]-urea
1-[2-(2-Methoxy-benzoyl)-4-methyl-phenyl]-3-(3-methyl-[1,2,4]thiadiazol-5-yl)-urea
1-(3-Isopropyl-[1,2,4]thiadiazol-5-yl)-3-[2-(2-methoxy-benzoyl)-4-methyl-phenyl]-urea
1-[2-(2-Methoxy-benzoyl)-4-methyl-phenyl]-3-(3-phenyl-[1,2,4]thiadiazol-5-yl)-urea
1-(4-Methyl-2-piperidin-1-yl-phenyl)-3-[1,2,4]thiadiazol-5-yl-urea
15 {5-[3-(4-Methyl-2-piperidin-1-yl-phenyl)-ureido]-[1,2,4]thiadiazol-3-yl}-acetic acid
N-{5-[3-(4-Methyl-2-piperidin-1-yl-phenyl)-ureido]-[1,2,4]thiadiazol-3-ylmethyl}-
methanesulfonamide
1-(3-Methoxy-[1,2,4]thiadiazol-5-yl)-3-(4-methyl-2-piperidin-1-yl-phenyl)-urea
1-(3-Ethyl-[1,2,4]thiadiazol-5-yl)-3-(4-methyl-2-piperidin-1-yl-phenyl)-urea
20 1-(4-Methyl-2-piperidin-1-yl-phenyl)-3-(3-methyl-[1,2,4]thiadiazol-5-yl)-urea
1-(3-Isopropyl-[1,2,4]thiadiazol-5-yl)-3-(4-methyl-2-piperidin-1-yl-phenyl)-urea
1-(4-Methyl-2-piperidin-1-yl-phenyl)-3-(3-phenyl-[1,2,4]thiadiazol-5-yl)-urea
1-(2-Cyclopentanecarbonyl-4-fluoro-phenyl)-3-[1,2,4]thiadiazol-5-yl-urea
{5-[3-(2-Cyclopentanecarbonyl-4-fluoro-phenyl)-ureido]-[1,2,4]thiadiazol-3-yl}-acetic acid
25 N-{5-[3-(2-Cyclopentanecarbonyl-4-fluoro-phenyl)-ureido]-[1,2,4]thiadiazol-3-ylmethyl}-
methanesulfonamide
1-(2-Cyclopentanecarbonyl-4-fluoro-phenyl)-3-(3-methoxy-[1,2,4]thiadiazol-5-yl)-urea
1-(2-Cyclopentanecarbonyl-4-fluoro-phenyl)-3-(3-ethyl-[1,2,4]thiadiazol-5-yl)-urea
1-(2-Cyclopentanecarbonyl-4-fluoro-phenyl)-3-(3-methyl-[1,2,4]thiadiazol-5-yl)-urea
30 1-(2-Cyclopentanecarbonyl-4-fluoro-phenyl)-3-(3-isopropyl-[1,2,4]thiadiazol-5-yl)-urea
1-(2-Cyclopentanecarbonyl-4-fluoro-phenyl)-3-(3-phenyl-[1,2,4]thiadiazol-5-yl)-urea
1-(4-Fluoro-2-isobutoxy-phenyl)-3-[1,2,4]thiadiazol-5-yl-urea
{5-[3-(4-Fluoro-2-isobutoxy-phenyl)-ureido]-[1,2,4]thiadiazol-3-yl}-acetic acid
N-{5-[3-(4-Fluoro-2-isobutoxy-phenyl)-ureido]-[1,2,4]thiadiazol-3-ylmethyl}-
35 methanesulfonamide

- 1-(4-Fluoro-2-isobutoxy-phenyl)-3-(3-methoxy-[1,2,4]thiadiazol-5-yl)-urea
1-(3-Ethyl-[1,2,4]thiadiazol-5-yl)-3-(4-fluoro-2-isobutoxy-phenyl)-urea
1-(4-Fluoro-2-isobutoxy-phenyl)-3-(3-methyl-[1,2,4]thiadiazol-5-yl)-urea
1-(4-Fluoro-2-isobutoxy-phenyl)-3-(3-isopropyl-[1,2,4]thiadiazol-5-yl)-urea
5 1-(4-Fluoro-2-isobutoxy-phenyl)-3-(3-phenyl-[1,2,4]thiadiazol-5-yl)-urea
1-[4-Fluoro-2-(2-methoxy-benzoyl)-phenyl]-3-[1,2,4]thiadiazol-5-yl-urea
(5-{3-[4-Fluoro-2-(2-methoxy-benzoyl)-phenyl]-ureido}-[1,2,4]thiadiazol-3-yl)-acetic acid
N-(5-{3-[4-Fluoro-2-(2-methoxy-benzoyl)-phenyl]-ureido}-[1,2,4]thiadiazol-3-ylmethyl)-methanesulfonamide
10 1-[4-Fluoro-2-(2-methoxy-benzoyl)-phenyl]-3-(3-methoxy-[1,2,4]thiadiazol-5-yl)-urea
1-(3-Ethyl-[1,2,4]thiadiazol-5-yl)-3-[4-fluoro-2-(2-methoxy-benzoyl)-phenyl]-urea
1-[4-Fluoro-2-(2-methoxy-benzoyl)-phenyl]-3-(3-methyl-[1,2,4]thiadiazol-5-yl)-urea
1-[4-Fluoro-2-(2-methoxy-benzoyl)-phenyl]-3-(3-isopropyl-[1,2,4]thiadiazol-5-yl)-urea
1-[4-Fluoro-2-(2-methoxy-benzoyl)-phenyl]-3-(3-phenyl-[1,2,4]thiadiazol-5-yl)-urea
15 1-(4-Fluoro-2-piperidin-1-yl-phenyl)-3-[1,2,4]thiadiazol-5-yl-urea
{5-[3-(4-Fluoro-2-piperidin-1-yl-phenyl)-ureido]-[1,2,4]thiadiazol-3-yl}-acetic acid
N-(5-[3-(4-Fluoro-2-piperidin-1-yl-phenyl)-ureido]-[1,2,4]thiadiazol-3-ylmethyl)-methanesulfonamide
1-(4-Fluoro-2-piperidin-1-yl-phenyl)-3-(3-methoxy-[1,2,4]thiadiazol-5-yl)-urea
20 1-(3-Ethyl-[1,2,4]thiadiazol-5-yl)-3-(4-fluoro-2-piperidin-1-yl-phenyl)-urea
1-(4-Fluoro-2-piperidin-1-yl-phenyl)-3-(3-methyl-[1,2,4]thiadiazol-5-yl)-urea
1-(4-Fluoro-2-piperidin-1-yl-phenyl)-3-(3-isopropyl-[1,2,4]thiadiazol-5-yl)-urea
1-(4-Fluoro-2-piperidin-1-yl-phenyl)-3-(3-phenyl-[1,2,4]thiadiazol-5-yl)-urea
1-(2-Cyclopentanecarbonyl-4-morpholin-4-yl-phenyl)-3-[1,2,4]thiadiazol-5-yl-urea
25 {5-[3-(2-Cyclopentanecarbonyl-4-morpholin-4-yl-phenyl)-ureido]-[1,2,4]thiadiazol-3-yl}-acetic acid
N-(5-[3-(2-Cyclopentanecarbonyl-4-morpholin-4-yl-phenyl)-ureido]-[1,2,4]thiadiazol-3-ylmethyl)-methanesulfonamide
1-(2-Cyclopentanecarbonyl-4-morpholin-4-yl-phenyl)-3-(3-methoxy-[1,2,4]thiadiazol-5-yl)-urea
30 urea
1-(2-Cyclopentanecarbonyl-4-morpholin-4-yl-phenyl)-3-(3-ethyl-[1,2,4]thiadiazol-5-yl)-urea
1-(2-Cyclopentanecarbonyl-4-morpholin-4-yl-phenyl)-3-(3-methyl-[1,2,4]thiadiazol-5-yl)-urea
1-(2-Cyclopentanecarbonyl-4-morpholin-4-yl-phenyl)-3-(3-isopropyl-[1,2,4]thiadiazol-5-yl)-urea
urea
35 1-(2-Cyclopentanecarbonyl-4-morpholin-4-yl-phenyl)-3-(3-phenyl-[1,2,4]thiadiazol-5-yl)-urea

- 1-(2-Isobutoxy-4-morpholin-4-yl-phenyl)-3-[1,2,4]thiadiazol-5-yl-urea
{5-[3-(2-Isobutoxy-4-morpholin-4-yl-phenyl)-ureido]-[1,2,4]thiadiazol-3-yl}-acetic acid
N-{5-[3-(2-Isobutoxy-4-morpholin-4-yl-phenyl)-ureido]-[1,2,4]thiadiazol-3-ylmethyl}-methanesulfonamide
- 5 1-(2-Isobutoxy-4-morpholin-4-yl-phenyl)-3-(3-methoxy-[1,2,4]thiadiazol-5-yl)-urea
1-(3-Ethyl-[1,2,4]thiadiazol-5-yl)-3-(2-isobutoxy-4-morpholin-4-yl-phenyl)-urea
1-(2-Isobutoxy-4-morpholin-4-yl-phenyl)-3-(3-methyl-[1,2,4]thiadiazol-5-yl)-urea
1-(2-Isobutoxy-4-morpholin-4-yl-phenyl)-3-(3-isopropyl-[1,2,4]thiadiazol-5-yl)-urea
1-(2-Isobutoxy-4-morpholin-4-yl-phenyl)-3-(3-phenyl-[1,2,4]thiadiazol-5-yl)-urea
- 10 1-[2-(2-Methoxy-benzoyl)-4-morpholin-4-yl-phenyl]-3-[1,2,4]thiadiazol-5-yl-urea
(5-[3-[2-(2-Methoxy-benzoyl)-4-morpholin-4-yl-phenyl]-ureido]-[1,2,4]thiadiazol-3-yl)-acetic acid
N-(5-[3-[2-(2-Methoxy-benzoyl)-4-morpholin-4-yl-phenyl]-ureido]-[1,2,4]thiadiazol-3-ylmethyl)-methanesulfonamide
- 15 1-[2-(2-Methoxy-benzoyl)-4-morpholin-4-yl-phenyl]-3-(3-methoxy-[1,2,4]thiadiazol-5-yl)-urea
1-(3-Ethyl-[1,2,4]thiadiazol-5-yl)-3-[2-(2-methoxy-benzoyl)-4-morpholin-4-yl-phenyl]-urea
1-[2-(2-Methoxy-benzoyl)-4-morpholin-4-yl-phenyl]-3-(3-methyl-[1,2,4]thiadiazol-5-yl)-urea
1-(3-Isopropyl-[1,2,4]thiadiazol-5-yl)-3-[2-(2-methoxy-benzoyl)-4-morpholin-4-yl-phenyl]-urea
1-[2-(2-Methoxy-benzoyl)-4-morpholin-4-yl-phenyl]-3-(3-phenyl-[1,2,4]thiadiazol-5-yl)-urea
- 20 1-(4-Morpholin-4-yl-2-piperidin-1-yl-phenyl)-3-[1,2,4]thiadiazol-5-yl-urea
{5-[3-(4-Morpholin-4-yl-2-piperidin-1-yl-phenyl)-ureido]-[1,2,4]thiadiazol-3-yl}-acetic acid
N-{5-[3-(4-Morpholin-4-yl-2-piperidin-1-yl-phenyl)-ureido]-[1,2,4]thiadiazol-3-ylmethyl}-methanesulfonamide
1-(3-Methoxy-[1,2,4]thiadiazol-5-yl)-3-(4-morpholin-4-yl-2-piperidin-1-yl-phenyl)-urea
- 25 1-(3-Ethyl-[1,2,4]thiadiazol-5-yl)-3-(4-morpholin-4-yl-2-piperidin-1-yl-phenyl)-urea
1-(3-Methyl-[1,2,4]thiadiazol-5-yl)-3-(4-morpholin-4-yl-2-piperidin-1-yl-phenyl)-urea
1-(3-Isopropyl-[1,2,4]thiadiazol-5-yl)-3-(4-morpholin-4-yl-2-piperidin-1-yl-phenyl)-urea
1-(4-Morpholin-4-yl-2-piperidin-1-yl-phenyl)-3-(3-phenyl-[1,2,4]thiadiazol-5-yl)-urea
4-{2-[3-(2-Cyclopentanecarbonyl-4-methyl-phenyl)-ureido]-thiazol-4-yl}-2-methyl-butyric acid
- 30 2-[3-(2-Cyclopentanecarbonyl-4-methyl-phenyl)-ureido]-4,5,6,7-terahydro-benzothiazole-7-carboxylic acid

BIOLOGICAL ASSAY

Glucokinase Activity Assay (I)

Glucokinase activity is assayed spectrometrically coupled to glucose 6-phosphate dehydrogenase to determine compound activation of glucokinase. The final assay contains
5 50 mM Hepes, pH 7.1, 50 mM KCl, 5 mM MgCl₂, 2 mM dithiothreitol, 0.6 mM NADP, 1 mM ATP, 0.195 μ M G-6-P dehydrogenase (from Roche, 127 671), 15 nM recombinant human glucokinase. The glucokinase is human liver glucokinase N-terminally truncated with an N-terminal His-tag ((His)₈-VEQILA.....Q466) and is expressed in E.coli as a soluble protein with enzymatic activity comparable to liver extracted GK.

10 The purification of His-tagged human glucokinase (hGK) was performed as follows:
The cell pellet from 50 ml E. coli culture was resuspended in 5 ml extraction buffer A (25 mM HEPES, pH 8.0, 1 mM MgCl₂, 150 mM NaCl, 2 mM mercaptoethanol) with addition of 0.25 mg/ml lysozyme and 50 μ g/ml sodium azide. After 5 minutes at room temperature 5 ml of extraction buffer B (1.5 M NaCl, 100 mM CaCl₂, 100 mM MgCl₂, 0.02 mg/ml DNase 1,
15 protease inhibitor tablet (Complete® 1697498): 1 tablet pr. 20 ml buffer) was added. The extract was then centrifugated at 15.000 g for 30 minutes. The resulting supernatant was loaded on a 1 ml Metal Chelate Affinity Chromatography (MCAC) Column charged with Ni²⁺. The column is washed with 2 volumes buffer A containing 20 mM imidazole and the bound his-tagged hGK is subsequently eluted using a 20 minute gradient of 20 to 500 mM
20 imidazol in buffer A. Fractions are examined using SDS-gel-electrophoresis, and fractions containing hGK (MW: 52 KDa) are pooled. Finally a gelfiltration step is used for final polishing and buffer exchange. hGK containing fractions are loaded onto a Superdex 75 (16/60) gelfiltration column and eluted with Buffer B (25 mM HEPES, pH 8.0, 1 mM MgCl₂, 150 mM NaCl, 1 mM Dithiothreitol). The purified hGK is examined by SDS-gel
25 electrophoresis and MALDI mass spectrometry and finally 20% glycerol is added before freezing. The yield from 50 ml E. coli culture is generally approximately 2-3 mg hGK with a purity >90%.

The compound to be tested is added into the well in final 2.5% DMSO concentration in an amount sufficient to give a desired concentration of compound, for instance 1, 5, 10, 25
30 or 50 μ M. The reaction starts after glucose is added to a final concentration of 2, 5, 10 or 15 mM. The assay uses a 96-well UV plate and the final assay volume used is 200 μ l/well. The plate is incubated at 25°C for 5 min and kinetics is measured at 340 nm in SpectraMax every 30 seconds for 5 minutes. Results for each compound are expressed as the fold activation of

the glucokinase activity compared to the activation of the glucokinase enzyme in an assay without compound after having been subtracted from a "blank", which is without glucokinase enzyme and without compound. The compounds in each of the Examples exhibits activation of glucokinase in this assay. A compound, which at a concentration of at or below 30 μ M gives 1.5 - fold higher glucokinase activity than the result from the assay without compound, is deemed to be an activator of glucokinase.

The glucose sensitivity of the compounds are measured at a compound concentration of 10 μ M and at glucose concentrations of 5 and 15 mM.

While the invention has been described and illustrated with reference to certain preferred embodiments thereof, those skilled in the art will appreciate that various changes, modifications and substitutions can be made therein without departing from the spirit and scope of the present invention. For example, effective dosages other than the preferred dosages as set forth herein may be applicable as a consequence of variations in the responsiveness of the mammal being treated for glucokinase-deficiency mediated disease(s). Likewise, the specific pharmacological responses observed may vary according to and depending on the particular active compound selected or whether there are present pharmaceutical carriers, as well as the type of formulation and mode of administration employed, and such expected variations or differences in the results are contemplated in accordance with the objects and practices of the present invention.

Glucokinase Activity Assay (II)

Determination of glycogen deposition in isolated rat hepatocytes:

Hepatocytes are isolated from rats fed *ad libitum* by a two-step perfusion technique. Cell viability, assessed by trypan blue exclusion, is consistently greater than 80%. Cells are plated onto collagen-coated 96-well plates in basal medium (Medium 199 (5.5 mM glucose) supplemented with 0.1 μ M dexamethasone, 100 units/ml penicillin, 100 mg/ml streptomycin, 2 mM L-glutamine and 1 nM insulin) with 4 % FCS at a cell density of 30,000 cells/well. The medium is replaced with basal medium 1 hour after initial plating in order to remove dead cells. Medium is changed after 24 hours to basal medium supplemented with 9.5 mM glucose and 10 nM insulin to induce glycogen synthesis, and experiments are performed the next day. The hepatocytes are washed twice with prewarmed (37°C) buffer A (117.6 mM NaCl, 5.4 mM KCl, 0.82 mM Mg_2SO_4 , 1.5 mM KH_2PO_4 , 20 mM HEPES, 9 mM $NaHCO_3$, 0.1% w/v HSA, and 2.25 mM $CaCl_2$, pH 7.4 at 37°C) and incubated in 100 μ l buffer A containing 15 mM glucose and increasing concentrations of the test compound, such as for

instance 1, 5, 10, 25, 50 or 100 μ M, for 180 minutes. Glycogen content is measured using standard procedures (Agius, L. et al, Biochem J. 266, 91-102 (1990)). A compound, which when used in this assay gives an significant increase in glycogen content compared to the result from the assay without compound, is deemed to have activity in this assay.


5 Glucokinase Activity Assay (III)

Stimulation of insulin secretion by glucokinase activators in INS-1E cells

- The glucose responsive β -cell line INS-1E is cultivated as described by Asfari M et al., Endocrinology, 130, 167-178 (1992). The cells are then seeded into 96 well cell culture plates and grown to a density of approximately 5×10^4 per well. Stimulation of glucose dependent insulin secretion is tested by incubation for 2 hours in Krebs Ringer Hepes buffer at glucose concentrations from 2.5 to 15 mM with or without addition of glucokinase activating compounds in concentrations of for instance 1, 5, 10, 25, 50 or 100 μ M, and the supernatants collected for measurements of insulin concentrations by ELISA (n= 4). A compound, which when used in this assay gives an significant increase in insulin secretion in response to glucose compared to the result from the assay without compound, is deemed to have activity in this assay.

CLAIMS

1. A compound of the general formula (I)



- 5 A¹ is selected from the group consisting of arylene, heteroarylene, fused cycloalkylarylene, fused heterocycliarylene, fused cycloalkylheteroarylene, or fused heterocycliheteroarylene; optionally substituted with one or more substituents R²³, R²⁴, R²⁵, R²⁶, and R²⁷, wherein R²³, R²⁴, R²⁵, R²⁶, and R²⁷ independently of each other are selected from the group consisting of
- 10 halogen, -C(O)OR², -C(O)R², -CN, -CF₃, -OCF₃, -NO₂, -OR², -NR²R³, C₁₋₆-alkyl-Z-, C₂₋₆-alkenyl-Z-, C₂₋₆-alkynyl-Z-, aryl-C₁₋₆-alkylene-Z-, heteroaryl-C₁₋₆-alkylene-Z-, heterocycliyl-C₁₋₆-alkylene-Z-, cycloalkyl-C₁₋₆-alkylene-Z-, N(R⁴R⁵)-C₁₋₆-alkylene-Z-, R⁶-W¹-Z-, R⁶-W¹-C₁₋₆-alkylene-Z-, R⁶-W¹-C₁₋₆-arylene-Z-, R⁶-W¹-heteroarylene-Z-, R⁶-W¹-heterocycliylene-Z-, R⁶-W¹-N(R⁴)-Z-, R⁶-N(R⁴)-Z-, R⁶-W¹-C₁₋₆-alkylene-Z-,
- 15 heterocycliyl-Z-C₁₋₆-alkylene-, heterocycliyl-C₁₋₆-alkylene-Z-C₁₋₆-alkylene-, C₃₋₁₀-cycloalkyl-Z-C₁₋₆-alkylene-, C₃₋₁₀-cycloalkyl-C₁₋₆-alkylene-Z-C₁₋₆-alkylene-, C₃₋₁₀-cycloalkyl-arylene-Z-, heterocycliyl-arylene-Z-, C₃₋₁₀-cycloalkyl-heteroarylene-Z-, heterocycliyl-heteroarylene-Z-, aryl-C₃₋₁₀-cycloalkylene-Z-, aryl-heterocycliylene-Z-, heteroaryl-C₃₋₁₀-cycloalkylene-Z-,
- 20 heteroaryl-heterocycliylene-Z-, heterocycliyl-C₃₋₁₀-cycloalkylene-Z-, aryl-heteroarylene-Z-, heteroaryl-arylene-Z-, aryl-arylene-Z-, heteroaryl-heteroarylene-Z-, wherein any mono- or divalent C₁₋₆-alkyl, C₃₋₁₀-cycloalkyl, heterocycliyl, aryl or heteroaryl moiety can optionally be substituted with one or more substituents independently selected from R², and wherein
- 25 R² and R³ independently of each other are hydrogen, halogen, hydroxy, -CN, -CF₃, -OCF₃, -NO₂, -C(O)OH, -NH₂, C₁₋₆-alkyl, C₁₋₆-alkoxy, aryloxy, R⁶-Z-, aryl-C₁₋₆-alkylene-, heteroaryl-C₁₋₆-alkylene-, C₁₋₆-alkyl-arylene-, C₁₋₆-alkyl-heteroarylene-, heteroaryl, or aryl;
- or

R^2 and R^3 , when attached to the same nitrogen atom, together with said nitrogen atom may form a 3 to 8 membered heterocyclic ring optionally containing one or two further heteroatoms selected from nitrogen, oxygen and sulfur, and optionally containing one or two double bonds;

5

Z and W^1 independently of each other are a direct bond, -O-, -N(R^7)-, -N(R^7)C(R^7R^8)-, -S-, -SO₂-, -C(O)N(R^7)-, -N(R^7)C(O)-, -N(R^7)C(O)C(R^7R^8)-, -N(R^7)CON(R^8)-, -N(R^7)SO₂-, -SO₂N(R^7)-, -C(O)-, -C(O)-O-, -N(R^7)SO₂N(R^8)-, or -O-C(O)-, wherein

10

R^7 and R^8 in each individual case independently of each other are hydrogen or C₁₋₆-alkyl; and

R^4 , R^5 , and R^6 independently of each other are selected from the group consisting of hydrogen, cyano, halogen, aryl, heteroaryl, heteroaryl-C₁₋₆-alkylene-, aryl-C₁₋₆-alkylene-, C₃₋₁₀-cycloalkyl, heterocyclyl optionally substituted with one or more C₁₋₆-alkyl, or C₁₋₆-alkyl optionally substituted with halogen, -S(O)₂CH₃ or COOH; or


15

R^4 and R^5 may be taken together to form a ring having the formula -(CH₂)_j-Q-(CH₂)_k- bonded to the nitrogen atom to which R^4 and R^5 are attached, wherein

20

j and k independently of each other is 1, 2, 3, or 4; and

Q is a direct bond, -CH₂-, -O-, -S-, -S(O)₂-, -C(O)-, -C(O)NH-, -NHC(O)-, -NHC(O)NH-, -NHSO₂-, -SO₂NH-, -C(O)-O-, -O-C(O)-, -NHSO₂NH-,



wherein

25

R^9 and R^{10} independently of each other are selected from the group consisting of hydrogen, aryl, C₁₋₆-alkyl, and aryl-alkylene-;

30

L^1 is a bond, -D-C₁₋₆-alkylene-E-, -D-C₂₋₆-alkenylene-E-, -D-C₂₋₆-alkynylene-E-, -D-cycloalkylene-E-, -D-heterocyclylene-E-, -O-, -S-, -S(O)-, -S(O)₂-,

-C(O)-, -N(R¹¹)-, or -C(=N-OR¹²)-, wherein

D and E independently of each other are a direct bond, -O- or -S-;

R¹¹ is selected from hydrogen, C₁₋₆-alkyl, aryl, carbamoyl, aryl-C₁₋₆-alkylene-,

heteroaryl-C₁₋₆-alkylene-, C₁₋₆-alkyl-O-C(O)-, aryl-C₁₋₆-alkylene-O-C(O)-,

5 heteroaryl-C₁₋₆-alkylene-O-C(O)-, C₁₋₆-alkyl-NH-C(O)-, aryl-C₁₋₆-alkylene-NH-C(O)-,

heteroaryl-C₁₋₆-alkylene-NH-C(O)-, C₁₋₆-alkyl-SO₂-, aryl-C₁₋₆-alkylene-SO₂-,

heteroaryl-C₁₋₆-alkylene-SO₂-, aryl-SO₂-, heteroaryl-SO₂-, C₁₋₆-alkyl-NH-SO₂-,

aryl-C₁₋₆-alkylene-NH-SO₂-, heteroaryl-C₁₋₆-alkylene-NH-SO₂-, C₁₋₆-alkyl-C(O)-,

aryl-C₁₋₆-alkylene-C(O)-, heteroaryl-C₁₋₆-alkylene-C(O)-, C₁₋₆-alkyl-Y-, aryl-Y-,

10 heteroaryl-Y-, aryl-C₁₋₆-alkylene-Y-, heteroaryl-C₁₋₆-alkylene-Y-,

N(R¹³)(R¹⁴)-C₁₋₆-alkylene-Y-, and R¹⁵-W²-C₁₋₆-alkylene-Y-, wherein

Y and W² independently of each other are a direct bond, -CH₂-, -SO₂-,

-N(H)CO-, -N(H)SO₂-, or -O-C(O)-;

R¹³ and R¹⁴ independently of each other are selected from hydrogen, aryl,

15 heteroaryl, C₁₋₆-alkyl, C₁₋₆-alkoxy, aryl-C₁₋₆-alkylene-,

heteroaryl-C₁₋₆-alkylene-, aryl-C₁₋₆-alkoxy-, heteroaryl-C₁₋₆-alkoxy-,

C₁₋₆-alkyl-arylene-, C₁₋₆-alkyl-heteroarylene-, C₁₋₆-alkoxy-heteroarylene-, or

C₁₋₆-alkoxy-arylene-;

or

20 R₁₃ and R₁₄ may be taken together to form a ring having the formula

-(CH₂)_o-X-(CH₂)_p- bonded to the nitrogen atom to which R₁₃ and R₁₄ are attached, wherein


o and p are independently of each other are 1, 2, 3, or 4; and

X is a direct bond, -CH₂-, -O-, -S-, -S(O₂)-, -C(O)-, -CON(H)-,

25 -NHC(O)-, -NHCON(H)-, -NHSO₂-, -SO₂N(H)-, -C(O)-O-, -O-C(O)-,

-NHSO₂NH-,

540



wherein

R^{16} and R^{17} are selected from hydrogen, aryl, heteroaryl, C_{1-6} -alkyl, C_{1-6} -alkoxy, aryl- C_{1-6} -alkylene-, heteroaryl- C_{1-6} -alkylene-, C_{1-6} -alkyl-arylene-, C_{1-6} -alkyl-heteroarylene-, C_{1-6} -alkoxy-arylene-, C_{1-6} -alkoxy-heteroarylene-, heteroarylaryl- C_{1-6} -alkoxy-, or aryl- C_{1-6} -alkoxy-; and

R^{15} is selected from the group consisting of aryl, heteroaryl, cycloalkyl,

heterocyclyl, C_{1-6} -alkyl, heteroaryl- C_{1-6} -alkylene-, or aryl- C_{1-6} -alkylene-; and

R^{12} is selected from hydrogen, aryl, heteroaryl, C_{1-6} -alkyl, aryl- C_{1-6} -alkylene-,

heteroaryl- C_{1-6} -alkylene-, C_{1-6} -alkyl-arylene-, C_{1-6} -alkyl-heteroarylene-,

C_{1-6} -alkoxy-heteroarylene-, or C_{1-6} -alkoxy-arylene-;

G^1 is C_{1-6} -alkyl, C_{3-10} -cycloalkyl, C_{3-10} -cycloalkyl- C_{1-6} -alkylene-, C_{2-6} -alkenyl, or C_{2-6} -alkynyl, all

of which may optionally be substituted with one or more substituents independently selected from the group consisting of -CN, -CF₃, -OCF₃, -OR¹⁸, -NR¹⁸R¹⁹, C_{3-10} -cycloalkyl and

C_{1-6} -alkyl, wherein

R^{18} and R^{19} independently of each other are hydrogen, C_{1-6} -alkyl,

heteroaryl- C_{1-6} -alkylene-, aryl- C_{1-6} -alkylene-, C_{1-6} -alkyl-arylene-,

C_{1-6} -alkyl-heteroarylene-, heteroaryl, or aryl;

or

R^{18} and R^{19} , when attached to the same nitrogen atom, together with the said nitrogen atom may form a 3 to 8 membered heterocyclic ring optionally containing one or two further heteroatoms selected from nitrogen, oxygen and sulfur, and optionally containing one or two double bonds;

or

G¹ is aryl, heteroaryl, heterocyclyl, fused cycloalkylheteroaryl, fused heterocyclylaryl, fused arylheterocyclyl, or fused cycloalkylaryl, all of which may optionally be substituted with one or more substituents selected from R⁴⁰, R⁴¹, and R⁴²;

- 5 L² is a direct bond, C₁₋₆-alkylene, C₂₋₆-alkenylene, C₂₋₆-alkynylene, -N(R²⁰)-, -C₁₋₆-alkylene-N(R²⁰)-, -C₂₋₆-alkenylene-N(R²⁰)-, -C₂₋₆-alkynylene-N(R²⁰)-, wherein

R²⁰ is hydrogen, or

R²⁰ is C₁₋₆-alkyl, C₂₋₆-alkenyl, C₂₋₆-alkynyl, cycloalkyl-W³-, heterocyclyl-W³-, aryl-W³-, heteroaryl-W³-, optionally substituted with one or more substituents R³⁰, R³¹, and

- 10 R³², wherein

W³ is C₁₋₆-alkylene or a direct bond;

wherein L¹ and L² are attached to adjacent atoms in A¹;

L³ is -C(O)-, -C(O)-C(O)-, -C(O)CH₂C(O)- or -S(O)₂-;

R¹ is hydrogen, or

- 15 R¹ is C₁₋₆-alkyl, C₂₋₆-alkenyl, C₂₋₆-alkynyl, cycloalkyl-W⁴-, heterocyclyl-W⁴-, aryl-W⁴-, or heteroaryl-W⁴-,


optionally substituted with one or more substituents R³³, R³⁴, and R³⁵, wherein

W⁴ is C₁₋₆-alkylene or a direct bond;

G² is heteroaryl, fused heterocyclylheteroaryl, or fused cycloalkylheteroaryl,

- 20 optionally substituted with one or more substituents R⁴³, R⁴⁴, and R⁴⁵, wherein said heteroaryl group possesses a nitrogen atom adjacent to the atom joining said heteroaryl group to -N(R¹)-;

or a group of the formula



- 25 wherein

G³ and G⁵ independently of each other are C₁₋₆-alkyl, C₂₋₆-alkenyl, C₂₋₆-alkynyl, cycloalkyl-R²²-, heterocyclyl-R²²-, aryl-R²²-, heteroaryl-R²²-,

optionally substituted with one or more substituents R⁴⁶, R⁴⁷, and R⁴⁸, wherein

R²² is alkylene or a direct bond; and

- 30 R²¹ is hydrogen, C₁₋₆-alkyl, C₂₋₆-alkenyl, C₂₋₆-alkynyl, cycloalkyl-W⁵-, or heterocyclyl-W⁵-,

optionally substituted with one or more substituents R³⁶, R³⁷, and R³⁸, or

R^{21} is aryl- W^5 -, or heteroaryl- W^5 -, optionally substituted with one or more substituents R^{49} , R^{50} , and R^{51} , wherein

W^5 is C_{1-6} -alkylene or a direct bond;

wherein

- 5 R^{30} , R^{31} , R^{32} , R^{33} , R^{34} , R^{35} , R^{36} , R^{37} , and R^{38} independently of each other are selected from -CHF₂-, -CF₃-, -OCF₃-, -OCHF₂-, -OCH₂CF₃-, -OCF₂CHF₂-, -S(O)₂CF₃-, -SCF₃-, -OR⁵²-, -NR⁵²R⁵³-, -SR⁵²-, -NR⁵²S(O)₂R⁵³-, -S(O)₂NR⁵²R⁵³-, -S(O)NR⁵²R⁵³-, -S(O)R⁵²-, -S(O)₂R⁵²-, -C(O)NR⁵²R⁵³-, -OC(O)NR⁵²R⁵³-, -NR⁵²C(O)R⁵³-, -CH₂C(O)NR⁵²R⁵³-, -OCH₂C(O)NR⁵²R⁵³-, -CH₂OR⁵²-, -CH₂NR⁵²R⁵³-, -OC(O)R⁵²-, -C(O)R⁵² and -C(O)OR⁵²-, or
- 10 C_{2-6} -alkenyl and C_{2-6} -alkynyl, which may optionally be substituted with one or more substituents selected from -CN-, -CF₃-, -OCF₃-, -OR⁵²-, -NR⁵²R⁵³ and C_{1-6} -alkyl; or C_{3-10} -cycloalkyl, C_{4-8} -cycloalkenyl, heterocyclyl, C_{3-10} -cycloalkyl- C_{1-6} -alkylene-, C_{3-10} -cycloalkyl- C_{1-6} -alkoxy-, C_{3-10} -cycloalkyloxy, C_{3-10} -cycloalkyl- C_{1-6} -alkylthio-, C_{3-10} -cycloalkylthio, C_{3-10} -cycloalkyl- C_{2-6} -alkenylene-, C_{3-10} -cycloalkyl- C_{2-6} -alkynylene-,
- 15 C_{4-8} -cycloalkenyl- C_{1-6} -alkylene-, C_{4-8} -cycloalkenyl- C_{2-6} -alkenylene-, C_{4-8} -cycloalkenyl- C_{2-6} -alkynylene-, heterocyclyl- C_{1-6} -alkylene-, heterocyclyl- C_{2-6} -alkenylene-, heterocyclyl- C_{2-6} -alkynylene-, aryl, aryloxy, aryloxycarbonyl, aroyl, aryl- C_{1-6} -alkoxy-, aryl- C_{1-6} -alkylene-, aryl- C_{2-6} -alkenylene-, aryl- C_{2-6} -alkynylene-, heteroaryl, heteroaryl- C_{1-6} -alkylene-, heteroaryl- C_{2-6} -alkenylene- and heteroaryl- C_{2-6} -alkynylene-, of
- 20 which the aryl and heteroaryl moieties optionally may be substituted with one or more substituents selected from halogen, -C(O)OR⁵²-, -CN-, -CF₃-, -OCF₃-, -NO₂-, -OR⁵²-, -NR⁵²R⁵³ and C_{1-6} -alkyl, wherein
 R^{52} and R^{53} independently of each other are hydrogen, C_{1-6} -alkyl, aryl- C_{1-6} -alkylene-, heteroaryl- C_{1-6} -alkylene- heteroaryl, or aryl;
- 25 or
 R^{52} and R^{53} , when attached to the same nitrogen atom, together with the said nitrogen atom may form a 3 to 8 membered heterocyclic ring optionally containing one or two further heteroatoms selected from nitrogen, oxygen and sulfur, and optionally containing one or two double bonds;
- 30 R^{40} , R^{41} , R^{42} , R^{43} , R^{44} , R^{45} , R^{46} , R^{47} , R^{48} , R^{49} , R^{50} and R^{51} independently of each other are -CN-, -NO₂-, -S(O)₂CF₃-, -SCF₃-, -OR⁵⁴-, -NR⁵⁴R⁵⁵-, -SR⁵⁴-, -NR⁵⁴S(O)₂R⁵⁵-, -S(O)₂NR⁵⁴R⁵⁵-, -S(O)NR⁵⁴R⁵⁵-, -S(O)R⁵⁴-, -S(O)₂R⁵⁴-, -C(O)NR⁵⁴R⁵⁵-, -OC(O)NR⁵⁴R⁵⁵-, -NR⁵⁴C(O)R⁵⁵-, halogen, -S- C_{1-6} -alkylene-OR⁵⁴-, -S(O)₂- C_{1-6} -alkylene-OR⁵⁴-, - C_{1-6} -alkylene-S-R⁵⁴-, - C_{1-6} -alkylene-S(O)R⁵⁴-, - C_{1-6} -alkylene-S(O)₂R⁵⁴-, - C_{1-6} -alkylene-N(R⁵⁴)S(O)₂R⁵⁵-,
- 35 -N(R⁵⁴)S(O)₂R⁵⁵-, - C_{1-6} -alkylene-CN,

- $-C_{1-6}$ -alkylene- $C(O)NR^{54}R^{55}$, $-C_{1-6}$ -alkylene- $N(R^{54})C(O)R^{55}$, $-N(R^{54})C(O)R^{55}$,
 $-C_{1-6}$ -alkylene- $N(R^{54})C(O)NR^{55}R^{56}$, $-C_{1-6}$ -alkylene- $NHC(=NR^{54})NR^{55}R^{56}$,
 $-C_{1-6}$ -alkylene- $N(R^{54})C(O)OR^{55}$, $-N(R^{54})C(O)OR^{55}$, $-C_{1-6}$ -alkylene- $C(O)OR^{54}$,
 $-C_{1-6}$ -alkylene- $N(R^{54})S(O)_2R^{55}$, $-OCH_2C(O)NR^{54}R^{55}$, $-O(CH_2)_{1-3}OR^{54}$, $-C_{1-6}$ -alkylene- $O-R^{54}$,
5 $-C_{1-6}$ -alkylene- $C(O)R^{54}$, $-C_{1-6}$ -alkylene- $NR^{54}R^{55}$, $-C(=NR^{54})-O-R^{55}$, $-C(=N(OR^{54}))C(O)OR^{55}$,
 $-C(=N(OR^{54}))C(O)R^{55}$, $-C_{1-6}$ -alkylene- $C(=N(OR^{54}))C(O)R^{55}$, $-C_{1-6}$ -alkylene- $N-O-R^{54}$,
 $-C_{1-6}$ -alkylene- $N(R^{54})S(O)_2NR^{55}R^{56}$, $-N(R^{54})S(O)_2NR^{55}R^{56}$, $-N(R^{54})S(O)_2NR^{55}R^{56}$, $-OC(O)R^{54}$,
 $-C_{1-6}$ -alkylene- $C(O)N(R^{54})S(O)_2R^{55}$, $-C(O)N(R^{54})S(O)_2R^{55}$, $-C_{1-6}$ -alkylene- $C(R^{54})=N-OR^{55}$,
 $-NHC(=NR^{54})NR^{55}R^{56}$, $-C_{1-6}$ -alkylene- $NHC(=NR^{54})NR^{54}R^{55}$, $-C_{1-6}$ -alkylene- $N=C(N(R^{54}R^{55}))_2$,
10 $-N=C(N(R^{54}R^{55}))_2$, $-C(O)R^{54}$ and $-C(O)OR^{54}$; or
 C_{1-6} -alkyl, C_{2-6} -alkenyl and C_{2-6} -alkynyl, each of which may optionally be substituted with one
or more substituents independently selected from halogen, R^{54} , $-CN$, $-CF_3$, $-OCF_3$, $-OR^{54}$, $-$
 $C(O)OR^{54}$, $-NR^{54}R^{55}$ and C_{1-6} -alkyl; or
 C_{3-10} -cycloalkyl, C_{4-8} -cycloalkenyl, heterocyclyl, C_{3-10} -cycloalkyl- C_{1-6} -alkylene-,
15 C_{3-10} -cycloalkyl- C_{1-6} -alkoxy-, C_{3-10} -cycloalkyloxy, C_{3-10} -cycloalkyl- C_{1-6} -alkylthio-,
 C_{3-10} -cycloalkylthio, C_{3-10} -cycloalkyl- C_{2-6} -alkenylene-, C_{3-10} -cycloalkyl- C_{2-6} -alkynylene-,
 C_{4-8} -cycloalkenyl- C_{1-6} -alkylene-, C_{4-8} -cycloalkenyl- C_{2-6} -alkenylene-,
 C_{4-8} -cycloalkenyl- C_{2-6} -alkynylene-, heterocyclyl- C_{1-6} -alkylene-, heterocyclyl- C_{2-6} -alkenylene-,
heterocyclyl- C_{2-6} -alkynylene- of which the heterocyclyl moieties optionally may be substituted
20 with one or more substituents independently selected from R^{70} ; or
aryl, aryloxy, aryloxycarbonyl, aroyl, aryl- C_{1-6} -alkoxy-, aryl- C_{1-6} -alkylene-,
aryl- C_{2-6} -alkenylene-, aryl- C_{2-6} -alkynylene-, heteroaryl, heteroaryl- C_{1-6} -alkylene-, heteroaryl-S-
 C_{1-6} -alkylene-, heteroaryl- C_{2-6} -alkenylene- and heteroaryl- C_{2-6} -alkynylene-, of which the aryl
and heteroaryl moieties optionally may be substituted with one or more substituents selected
25 from halogen, $-C(O)OR^{54}$, $-CN$, $-CF_3$, $-OCF_3$, $-NO_2$, $-OR^{54}$, $-NR^{54}R^{55}$ or C_{1-6} -alkyl, wherein
 R^{54} , R^{55} and R^{56} independently of each other are hydrogen, C_{1-6} -alkyl, $-Si(C_{1-6}$ -alkyl)₃,
 C_{1-6} -alkyl-arylene-, C_{1-6} -alkyl-heteroarylene-, aryl- C_{1-6} -alkylene-,
heteroaryl- C_{1-6} -alkylene-, heterocyclyl, heterocyclyl- C_{1-6} -alkylene-, heteroaryl, or aryl,
each of which is optionally substituted with one or more substituents independently
30 selected from R^{71} ;
or
 R^{54} and R^{55} independently of each other are hydrogen or $-(CHR^{72})_u-(CHR^{73})_v-W^6$,
wherein

u is 0, 1 or 2;

35

v is 0, 1 or 2;

R^{72} and R^{73} independently of each other are hydrogen, C_{1-6} -alkyl, C_{1-6} -alkyl-arylene-aryl, hydroxy, hydroxyalkyl, $-C(O)O-R^{75}$, amino, or aminoalkyl;

W^6 is hydrogen, $-O-R^{75}$, $-C(O)O-R^{75}$, $-C(O)-R^{75}$, $-CONR^{75}R^{76}$, $-NR^{75}R^{76}$, $-NHCH_2C(O)R^{75}$, $-NHC(O)R^{75}$, $-NHC(O)OR^{75}$, $-S(O)_2R^{75}$, $-NHS(O)_2R^{75}$, alkylamino, or dialkylamino, or

W^6 is a five or six membered ring wherein at least one ring atom is nitrogen and the remaining ring atoms are either carbon or oxygen or optionally substituted with C_{1-6} -alkyl, $-C(O)O-R^{75}$, $-(CH_2)_{1-3}C(O)O-R^{75}$, $=O$;

or

W^6 is phthalimido or heterocyclyl.

or

R^{54} and R^{55} , when attached to the same nitrogen atom, together with the said nitrogen atom may form a 3 to 8 membered heterocyclic ring optionally containing one or two further heteroatoms selected from nitrogen, oxygen and sulfur, and optionally containing one or two double bonds, and optionally substituted with one or more C_{1-6} -alkyl groups;

R^{70} is $=O$, $-C(O)CH_3$, $-S(O)_2CH_3$, $-CF_3$, $-C(O)O-R^{75}$, $-(CH_2)_{1-3}C(O)O-R^{75}$ or C_{1-6} -alkyl;

R^{71} is $=O$, C_{1-6} -alkyl, cycloalkyl, $-C(O)O-R^{75}$, $-(CH_2)_{1-3}C(O)O-R^{75}$, $-(CH_2)_{1-3}NR^{75}R^{76}$, $-OH$ or amino;

R^{75} and R^{76} independently of each other is hydrogen, halogen, $-OH$, $-CF_3$, or C_{1-6} -alkyl optionally substituted with $-NH_2$;

or a pharmaceutically acceptable salt, solvate, or prodrug thereof

2. A compound according to claim 1, wherein

A^1 is arylene or heteroarylene, optionally substituted with one or more substituents R^{23} , R^{24} , R^{25} , R^{26} , and R^{27} , wherein

R^{23} , R^{24} , R^{25} , R^{26} , and R^{27} independently of each other are selected from the group consisting of

halogen, $-C(O)OR^2$, $-CN$, $-CF_3$, $-OCF_3$, $-NO_2$, $-OR^2$, $-NR^2R^3$, C_{1-6} -alkyl-Z-, C_{2-6} -alkenyl-Z-, C_{2-6} -alkynyl-Z-, cycloalkyl-Z-, heterocyclyl-Z-, aryl-Z-, heteroaryl-Z-, aryl- C_{1-6} -alkylene-Z-, heteroaryl- C_{1-6} -alkylene-Z-, heterocyclyl- C_{1-6} -alkylene-Z-,

cycloalkyl-C₁₋₆-alkylene-Z-, N(R⁴R⁵)-C₁₋₆-alkylene-Z-, R⁶-W¹-Z-, R⁶-W¹-N(R⁴)-Z-, R⁶-N(R⁴)-Z, and R⁶-W¹-C₁₋₆-alkylene-Z-, wherein

R², R³, R⁴, R⁵, R⁶, Z, and W¹ are as defined in claim 1.

3. A compound according to claim 2, wherein

5 A¹ is C₆₋₁₀-arylene or C₄₋₁₀-heteroarylene, optionally substituted with one or more substituents R²³, R²⁴, R²⁵, R²⁶, and R²⁷, wherein

R²³, R²⁴, R²⁵, R²⁶, and R²⁷ independently of each other are selected from the group consisting of

10 halogen, -C(O)OR², -CN, -CF₃, -OCF₃, -NO₂, -OR², -NR²R³, C₁₋₆-alkyl-Z-, C₂₋₆-alkenyl-Z-, C₂₋₆-alkynyl-Z-, cycloalkyl-Z-, heterocyclyl-Z-, aryl-Z-, heteroaryl-Z-, aryl-C₁₋₆-alkylene-Z-, heteroaryl-C₁₋₆-alkylene-Z-, heterocyclyl-C₁₋₆-alkylene-Z-, cycloalkyl-C₁₋₆-alkylene-Z-, N(R⁴R⁵)-C₁₋₆-alkylene-Z-, R⁶-W¹-Z-, R⁶-W¹-N(R⁴)-Z-, R⁶-N(R⁴)-Z, and R⁶-W¹-C₁₋₆-alkylene-Z-, wherein

R², R³, R⁴, R⁵, R⁶, Z, and W¹ are as defined in claim 1.

15

4. A compound according to claim 3 wherein


A¹ is phenylene optionally substituted with one or more substituents R²³, R²⁴, R²⁵, R²⁶, and R²⁷, wherein

20 R²³, R²⁴, R²⁵, R²⁶, and R²⁷ independently of each other are selected from the group consisting of

25 halogen, -C(O)OR², -CN, -CF₃, -OCF₃, -NO₂, -OR², -NR²R³, C₁₋₆-alkyl-Z-, C₂₋₆-alkenyl-Z-, C₂₋₆-alkynyl-Z-, cycloalkyl-Z-, heterocyclyl-Z-, aryl-Z-, heteroaryl-Z-, aryl-C₁₋₆-alkylene-Z-, heteroaryl-C₁₋₆-alkylene-Z-, heterocyclyl-C₁₋₆-alkylene-Z-, cycloalkyl-C₁₋₆-alkylene-Z-, N(R⁴R⁵)-C₁₋₆-alkylene-Z-, R⁶-W¹-Z-, R⁶-W¹-N(R⁴)-Z-, R⁶-N(R⁴)-Z, and R⁶-W¹-C₁₋₆-alkylene-Z-, wherein

R², R³, R⁴, R⁵, R⁶, Z, and W¹ are as defined in claim 1.


5. A compound according to claim 4 of the formula (Ia)



Formula (Ia)

wherein L^1 , G^1 , L^2 , L^3 , R^1 , G^2 and R^{25} are as defined in claim 1.

6. A compound according to claim 4 of the formula (Ib)



Formula (Ib)

wherein L^1 , G^1 , L^2 , L^3 , R^1 , G^2 and R^{24} are as defined in claim 1.

7. A compound according to any one of the claims 1 to 6, wherein

R^{23} , R^{24} , R^{25} , R^{26} , and R^{27} independently of each other are selected from the group consisting of

halogen, $-C(O)OR^2$, $-CN$, $-CF_3$, $-OCF_3$, $-NO_2$, $-OR^2$, $-NR^2R^3$, C_{1-6} -alkyl-Z-, cycloalkyl-Z-, heterocyclyl-Z-, aryl-Z-, or heteroaryl-Z-, $N(R^4R^5)-C_{1-6}$ -alkylene-Z-, R^6-W^1-Z -, $R^6-W^1-N(R^4)-Z$ -, $R^6-N(R^4)-Z$ -, and $R^6-W^1-C_{1-6}$ -alkylene-Z-, wherein

R^2 , R^3 , R^4 , R^5 , R^6 , Z , and W^1 are as defined in claim 1.

8. A compound according to claim 7 wherein

R^{23} , R^{24} , R^{25} , R^{26} , and R^{27} independently of each other are selected from the group consisting of

halogen, $-CN$, $-CF_3$, $-OR^2$, $-NR^2R^3$, C_{1-6} -alkyl-Z-, cycloalkyl-Z-, heterocyclyl-Z-, aryl-Z-, or heteroaryl-Z-, R^6-W^1-Z -, $R^6-W^1-N(R^4)-Z$ -, $R^6-N(R^4)-Z$ -, and $R^6-W^1-C_{1-6}$ -alkylene-Z-, wherein

R^2 , R^3 , R^4 , R^5 , R^6 , Z , and W^1 are as defined in claim 1.

9. a compound according to claim 8 wherein

R^{23} , R^{24} , R^{25} , R^{26} , and R^{27} independently of each other are selected from the group consisting of

halogen, $-\text{OR}^2$, $-\text{NR}^2\text{R}^3$, $\text{C}_{1-6}\text{-alkyl-Z-}$, $\text{R}^6\text{-W}^1\text{-Z-}$, and $\text{R}^6\text{-W}^1\text{-C}_{1-6}\text{-alkylene-Z-}$, wherein R^2 , R^3 , R^4 , R^5 , R^6 , Z , and W^1 are as defined in claim 1.

10. A compound according to claim 9 wherein

5 R^{23} , R^{24} , R^{25} , R^{26} , and R^{27} independently of each other are selected from the group consisting of F , Cl , Br , and methyl.

11. A compound according to any one of the claims 1 to 10, wherein

R^4 , R^5 , and R^6 independently of each other are selected from the group consisting of hydrogen, aryl, $\text{C}_{1-6}\text{-alkyl}$, heteroaryl- $\text{C}_{1-6}\text{-alkylene-}$, and aryl- $\text{C}_{1-6}\text{-alkylene-}$.

10 12. A compound according to claim 11, wherein

R^4 , R^5 , and R^6 independently of each other are hydrogen or $\text{C}_{1-6}\text{-alkyl}$.


13. A compound according to claim 12, wherein

R^4 , R^5 , and R^6 are hydrogen.

14. A compound according to any one of the claims 1 to 9, wherein

15 R^4 and R^5 is taken together to form a ring having the formula $-(\text{CH}_2)_j\text{-Q-(CH}_2)_k\text{-}$ bonded to the nitrogen atom to which R^4 and R^5 are attached, wherein j and k independently of each other is 1, 2, 3, or 4;

Q is a direct bond, $-\text{CH}_2\text{-}$, $-\text{O-}$, $-\text{S-}$, $-\text{S}(\text{O}_2)\text{-}$, $-\text{C}(\text{O})\text{-}$, $-\text{C}(\text{O})\text{NH-}$, $-\text{NHC}(\text{O})\text{-}$, $-\text{NHC}(\text{O})\text{NH-}$, $-\text{NHSO}_2\text{-}$, $-\text{SO}_2\text{NH-}$, $-\text{C}(\text{O})\text{-O-}$, $-\text{O-C}(\text{O})\text{-}$, $-\text{NHSO}_2\text{NH-}$,




20

wherein

R^9 and R^{10} independently of each other are selected from the group consisting of hydrogen, aryl, $\text{C}_{1-6}\text{-alkyl}$, and aryl- $\text{C}_{1-6}\text{-alkyl-}$.

15. A compound according to claim 14, wherein

Q is a direct bond, $-\text{CH}_2-$, $-\text{O}-$, $-\text{S}-$, $-\text{S}(\text{O}_2)-$, $-\text{C}(\text{O})-$, $-\text{C}(\text{O})\text{NH}-$, $-\text{NHC}(\text{O})-$, $-\text{NHC}(\text{O})\text{NH}-$, $-\text{NHSO}_2-$, $-\text{SO}_2\text{NH}-$, $-\text{C}(\text{O})\text{O}-$, $-\text{O}-\text{C}(\text{O})-$, $-\text{NHSO}_2\text{NH}-$,




wherein

- 5 R^9 and R^{10} independently of each other are hydrogen or C_{1-6} -alkyl.

16. A compound according to claim 15, wherein

Q is a direct bond, $-\text{CH}_2-$, $-\text{O}-$, $-\text{S}-$, $-\text{S}(\text{O}_2)-$, $-\text{C}(\text{O})-$, $-\text{C}(\text{O})\text{NH}-$, $-\text{NHC}(\text{O})-$, $-\text{NHC}(\text{O})\text{NH}-$, $-\text{NHSO}_2-$, $-\text{SO}_2\text{NH}-$, $-\text{C}(\text{O})\text{O}-$, $-\text{O}-\text{C}(\text{O})-$, $-\text{NHSO}_2\text{NH}-$,



- 10 17. A compound according to claim 16, wherein Q is a direct bond.

18. A compound according to any one of the claims 1 to 17, wherein

W^1 is a direct bond, $-\text{O}-$, $-\text{C}(\text{O})-$, $-\text{NH}-$, $-\text{S}-$, $-\text{SO}_2-$, $-\text{C}(\text{O})\text{NH}-$, $-\text{NHC}(\text{O})-$, $-\text{N}(\text{H})\text{CON}(\text{H})-$, $-\text{N}(\text{H})\text{SO}_2-$, $-\text{SO}_2\text{N}(\text{H})-$, $-\text{C}(\text{O})\text{O}-$, $-\text{N}(\text{H})\text{SO}_2\text{N}(\text{H})-$, or $-\text{O}-\text{C}(\text{O})-$.

19. A compound according to claim 18, wherein


- 15 W^1 is a direct bond, $-\text{O}-$, $-\text{C}(\text{O})-$, $-\text{SO}_2-$, $-\text{C}(\text{O})\text{NH}-$, $-\text{NHC}(\text{O})-$, $-\text{N}(\text{H})\text{SO}_2-$, $-\text{C}(\text{O})\text{O}-$, or $-\text{O}-\text{C}(\text{O})-$.

20. A compound according to claim 19, wherein W^1 is a direct bond or $-\text{C}(\text{O})\text{O}-$.

21. A compound according to claim 20, wherein W^1 is a direct bond.

22. A compound according to claim 7, wherein
at least one of R^{23} , R^{24} , R^{25} , R^{26} , and R^{27} is halogen, $-C(O)OR^2$, $-CN$, $-CF_3$, $-OCF_3$,
 $-NO_2$, $-OR^2$, $-NR^2R^3$, C_{1-6} -alkyl-Z-, cycloalkyl-Z-, heterocyclyl-Z-, aryl-Z-, or
heteroaryl-Z-, wherein
5 R^2 , R^3 , and Z are as defined in claim 1.
23. A compound according to claim 22, wherein
at least one of R^{23} , R^{24} , R^{25} , R^{26} , and R^{27} is halogen, $-C(O)OR^2$, $-CN$, $-CF_3$, $-OCF_3$,
 $-NO_2$, $-OR^2$, $-NR^2R^3$, C_{1-6} -alkyl-Z-, C_{3-10} -cycloalkyl-Z-, C_{3-10} -heterocyclyl-Z-,
 C_{3-10} -aryl-Z-, or C_{3-10} -heteroaryl-Z-, wherein
10 R^2 , R^3 , and Z are as defined in claim 1.
24. A compound according to claim 23, wherein
at least one of R^{23} , R^{24} , R^{25} , R^{26} , and R^{27} is halogen, $-C(O)OR^2$, $-CN$, $-NO_2$, $-OR^2$,
 $-NR^2R^3$, C_{1-6} -alkyl-Z-, C_{3-10} -cycloalkyl-Z-, C_{3-10} -heterocyclyl-Z-, C_{3-10} -aryl-Z-, or
 C_{3-10} -heteroaryl-Z-, wherein
15 R^2 , R^3 , and Z are as defined in claim 1.
25. A compound according to claim 24, wherein
at least one of R^{23} , R^{24} , R^{25} , R^{26} , and R^{27} is F, Cl, Br, or methyl.
26. A compound according to claim 24, wherein
at least one of R^{23} , R^{24} , R^{25} , R^{26} , and R^{27} is halogen, $-C(O)OR^2$, $-CN$, $-NO_2$, $-OR^2$, or
20 $-NR^2R^3$, wherein
 R^2 , R^3 , and Z are as defined in claim 1.
27. A compound according to any one of the claims 1 to 26, wherein
 R^2 and R^3 independently of each other are hydrogen, C_{1-6} -alkyl, aryl- C_{1-6} -alkylene-,
heteroaryl- C_{1-6} -alkylene-, C_{1-6} -alkyl-arylene-, C_{1-6} -alkyl-heteroarylene-, heteroaryl, or
25 aryl.
28. A compound according to claim 27, wherein
 R^2 and R^3 independently of each other, are hydrogen, C_{1-6} -alkyl, aryl- C_{1-6} -alkylene-
or aryl.

29. A compound according to claim 28, wherein
R² is hydrogen or C₁₋₆-alkyl.
30. A compound according to claim 29, wherein
R² is hydrogen.
- 5 31. A compound according to any one of the claims 1 to 30, wherein
R³ is hydrogen or C₁₋₆-alkyl.
32. A compound according to claim 31, wherein
R³ is hydrogen.
- 10 33. A compound according to any one of the claims 1 to 22, wherein
R² and R³, when attached to the same nitrogen atom, together with said nitrogen atom may form a 3 to 8 membered heterocyclic ring optionally containing one or two further heteroatoms selected from nitrogen, oxygen and sulfur, and optionally containing one or two double bonds.
- 15 34. A compound according to any one of the claims 1 to 33, wherein
Z is a direct bond, -O-, -NH-, -NHCH₂-, -S-, -SO₂-, -C(O)NH-, -NHC(O)-, -N(H)CON(H)-, -N(CH₃)CONH-, -N(H)SO₂-, -SO₂N(H)-, -C(O)-O-, -N(H)SO₂N(H)-, or -O-C(O)-.
- 20 35. A compound according to claim 34, wherein
Z is a direct bond, -O-, -S-, -SO₂-, -C(O)NH-, -NHC(O)-, -N(H)SO₂-, -C(O)-O-, -N(H)SO₂N(H)-, or -O-C(O)-.
36. A compound according to claim 35, wherein
Z is a direct bond, -NHC(O)-, or -NHS(O)₂-.
37. A compound according to claim 2, wherein
A¹ is



, wherein

R²³, R²⁴, R²⁵, and R²⁶, independently of each other, are hydrogen or as defined in claim 1.

38. A compound according to claim 37, wherein

- 5 R²³, R²⁴, R²⁵, and R²⁶ independently of each other are selected from the group consisting of
 halogen, -C(O)OR², -CN, -CF₃, -OCF₃, -NO₂, -OR², -NR²R³, C₁₋₆-alkyl-Z-,
 cycloalkyl-Z-, heterocyclyl-Z-, aryl-Z-, or heteroaryl-Z-, N(R⁴R⁵)-C₁₋₆-alkylene-Z-,
 R⁶-W¹-Z-, R⁶-W¹-N(R⁴)-Z-, R⁶-N(R⁴)-Z-, and R⁶-W¹-C₁₋₆-alkylene-Z-, wherein
 R², R³, R⁴, R⁵, R⁶, Z, and W¹ are as defined in claim 1.

10 39. A compound according to claim 37 or 38, wherein

R⁴, R⁵, and R⁶ independently of each other are selected from the group consisting of
 hydrogen, aryl, C₁₋₆-alkyl, heteroaryl-C₁₋₆-alkylene-, and aryl-C₁₋₆-alkylene-.

40. A compound according to claim 39, wherein

R⁴, R⁵, and R⁶ independently of each other are hydrogen or C₁₋₆-alkyl.


15 41. A compound according to claim 40, wherein

R⁴, R⁵, and R⁶ are hydrogen.

42. A compound according to claim 37 or 38, wherein

- 20 R⁴ and R⁵ is taken together to form a ring having the formula -(CH₂)_j-Q-(CH₂)_k-
 bonded to the nitrogen atom to which R⁴ and R⁵ are attached, wherein
 j and k independently of each other is 1, 2, 3, or 4;
 Q is a direct bond, -CH₂-, -O-, -S-, -S(O₂)-, -C(O)-, -C(O)NH-, -NHC(O)-,
 -NHC(O)NH-, -NHSO₂-, -SO₂NH-, -C(O)-O-, -O-C(O)-, -NHSO₂NH-,

552




wherein

R^9 and R^{10} independently of each other are selected from the group consisting of hydrogen, aryl, C_{1-6} -alkyl, and arylalkyl-.

5 43. A compound according to claim 42, wherein

Q is a direct bond, $-\text{CH}_2-$, $-\text{O}-$, $-\text{S}-$, $-\text{S}(\text{O}_2)-$, $-\text{C}(\text{O})-$, $-\text{C}(\text{O})\text{NH}-$, $-\text{NHC}(\text{O})-$, $-\text{NHC}(\text{O})\text{NH}-$, $-\text{NHSO}_2-$, $-\text{SO}_2\text{NH}-$, $-\text{C}(\text{O})-\text{O}-$, $-\text{O}-\text{C}(\text{O})-$, $-\text{NHSO}_2\text{NH}-$,




wherein

10 R^9 and R^{10} independently of each other are hydrogen or alkyl.

44. A compound according to claim 43, wherein

Q is a direct bond, $-\text{CH}_2-$, $-\text{O}-$, $-\text{S}-$, $-\text{S}(\text{O}_2)-$, $-\text{C}(\text{O})-$, $-\text{C}(\text{O})\text{NH}-$, $-\text{NHC}(\text{O})-$, $-\text{NHC}(\text{O})\text{NH}-$, $-\text{NHSO}_2-$, $-\text{SO}_2\text{NH}-$, $-\text{C}(\text{O})-\text{O}-$, $-\text{O}-\text{C}(\text{O})-$, $-\text{NHSO}_2\text{NH}-$,



15 45. A compound according to claim 44, wherein

Q is a direct bond.

46. A compound according to any one of the claims 37 to 45, wherein
W¹ is a direct bond, -O-, -C(O)-, -NH-, -S-, -SO₂-, -C(O)NH-, -NHC(O)-,
-N(H)CON(H)-, -N(H)SO₂-, -SO₂N(H)-, -C(O)-O-, -N(H)SO₂N(H)-, or -O-C(O)-.
- 5 47. A compound according to claim 46 wherein
W¹ is a direct bond, -O-, -C(O)-, -SO₂-, -C(O)NH-, -NHC(O)-, -N(H)SO₂-, -C(O)-O-,
or -O-C(O)-.
48. A compound according to claim 47, wherein
W¹ is a direct bond, or -C(O)-O-.
- 10 49. A compound according to claim 48, wherein
W¹ is a direct bond.
50. A compound according to claim 37, wherein
at least one of R²³, R²⁴, R²⁵, and R²⁶ is halogen, -C(O)OR², -CN, -CF₃, -OCF₃, -NO₂,
-OR², -NR²R³, C₁₋₆-alkyl-Z-, cycloalkyl-Z-, heterocyclyl-Z-, aryl-Z-, or heteroaryl-Z-,
15 wherein
R², R³, and Z are as defined in claim 37.
51. A compound according to claim 50, wherein
at least one of R²³, R²⁴, R²⁵, and R²⁶ is halogen, -C(O)OR², -CN, -CF₃, -OCF₃, -NO₂,
-OR², -NR²R³, C₁₋₆-alkyl-Z-, C₃₋₁₀-cycloalkyl-Z-, C₃₋₁₀-heterocyclyl-Z-, C₃₋₁₀-aryl-Z-, or
20 C₃₋₁₀-heteroaryl-Z-, wherein
R², R³, and Z are as defined in claim 37.
52. A compound according to claim 51, wherein
at least one of R²³, R²⁴, R²⁵, and R²⁶ is halogen, -C(O)OR², -CN, -NO₂, -OR²,
-NR²R³, C₁₋₆-alkyl-Z-, C₃₋₁₀-cycloalkyl-Z-, C₃₋₁₀-heterocyclyl-Z-, C₃₋₁₀-aryl-Z-, or
25 C₃₋₁₀-heteroaryl-Z-, wherein
R², R³, and Z are as defined in claim 37.
53. A compound according to any one of the claims 37 to 52, wherein

Z is a direct bond, -O-, -NH-, -NHCH₂-, -S-, -SO₂-, -C(O)NH-, -NHC(O)-, -N(H)CON(H)-, -N(CH₃)CON(H)-, -N(H)SO₂-, -SO₂N(H)-, -C(O)-O-, -N(H)SO₂N(H)-, or -O-C(O)-.

54. A compound according to claim 53, wherein

5 Z is a direct bond, -O-, -S-, -SO₂-, -C(O)NH-, -NHC(O)-, -N(H)SO₂-, -C(O)-O-, -N(H)SO₂N(H)-, or -O-C(O)-.

55. A compound according to claim 54, wherein

Z is a direct bond, -NHC(O)-, or -NHS(O)₂.

56. A compound according to claim 52, wherein

10 at least one of R²³, R²⁴, R²⁵, and R²⁶ is halogen, -C(O)OR², -CN, -NO₂, -OR², -NR²R³, or C₁₋₆-alkyl, wherein R², and R³ are as defined in claim 37.

57. A compound according to any one of the claims 37 to 56, wherein

15 R² and R³ independently of each other are hydrogen, C₁₋₆-alkyl, aryl-C₁₋₆-alkylene-, heteroaryl-C₁₋₆-alkylene-, C₁₋₆-alkyl-arylene-, C₁₋₆-alkyl-heteroarylene-, heteroaryl, or aryl.

58. A compound according to claim 57, wherein

R² and R³ independently of each other, are hydrogen, C₁₋₆-alkyl, aryl-C₁₋₆-alkylene- or aryl.

20 59. A compound according to claim 58, wherein

R² is hydrogen or C₁₋₆-alkyl.

60. A compound according to claim 59, wherein

R² is hydrogen.

61. A compound according to any one of the claims 37 to 60, wherein

25 R³ is hydrogen or C₁₋₆-alkyl.

62. A compound according to claim 61, wherein

R^3 is hydrogen.

63. A compound according to any one of the claims 37 to 50, wherein
 R^2 and R^3 , when attached to the same nitrogen atom, together with said nitrogen
atom may form a 3 to 8 membered heterocyclic ring optionally containing one or two
5 further heteroatoms selected from nitrogen, oxygen and sulfur, and optionally
containing one or two double bonds.
64. A compound according to any one of the claims 37 to 55, wherein
at least one of R^{23} , R^{24} , R^{25} , and R^{26} are hydrogen.
65. A compound according to claim 64, wherein
10 at least two of R^{23} , R^{24} , R^{25} , and R^{26} are hydrogen.
66. A compound according to claim 65, wherein
 R^{23} and R^{26} are hydrogen.
67. A compound according to claim 65 or claim 66, wherein
at least three of R^{23} , R^{24} , R^{25} , and R^{26} are hydrogen.
- 15 68. A compound according to any one of the claims 37 to 67, wherein R^{24} or R^{25} is halogen.
69. A compound according to claim 68, wherein R^{24} or R^{25} is fluoro.
70. A compound according to any one of the claims 37 to 67, wherein R^{24} or R^{25} is C_{1-6} -alkyl.
71. A compound according to claim 68, wherein R^{24} or R^{25} is methyl.
72. A compound according to any one of the claims 37 to 67, wherein
20 R^{24} is hydrogen.
73. A compound according to any one of the claims 37 to 72, wherein
 R^{25} is hydrogen.
74. A compound according to claim 37, wherein
 R^{23} , R^{24} , R^{25} , and R^{26} are hydrogen.

75. A compound according to any one of the claims 1 to 74, wherein
L¹ is a bond, -D-alkylene-E-, -O-, -S-, -S(O)-, -S(O)₂-, -C(O)-, -N(R¹¹)-, or -C(=N-OR¹²), wherein
D, E, R¹¹ and R¹² are as defined in claim 1.
- 5 76. A compound according to claim 75, wherein
L¹ is a bond, -D-alkylene-E-, -O-, -C(O)-, -N(R¹¹)-, or -C(=N-OR¹²)-, wherein
D, E, R¹¹ and R¹² are as defined in claim 1.
77. A compound according to claim 75, wherein
L¹ is -O-.
- 10 78. A compound according to claim 75, wherein
L¹ is -S-.
79. A compound according to claim 75, wherein
L¹ is a bond.
80. A compound according to claim 75, wherein
15 L¹ is -C(O)-.
81. A compound according to any one of the claims 1 to 76, wherein
D is a direct bond or -O-;
E is a direct bond or -O-; and
R¹¹ and R¹² are as defined in claim 1.
- 20 82. A compound according to claim 81, wherein
D is a direct bond.
83. A compound according to claim 81, wherein
D is -O-.
84. A compound according to any one of the claims 81 to 83, wherein
25 E is a direct bond.
85. A compound according to any one of the claims 81 to 83, wherein

E is -O-.

86. A compound according to any one of the claims 1 to 85, wherein

R¹¹ is selected from hydrogen, C₁₋₆-alkyl, aryl, carbamoyl, aryl-C₁₋₆-alkylene-,
C₁₋₆-alkyl-NH-C(O)-, aryl-C₁₋₆-alkylene-NH-C(O)-, C₁₋₆-alkyl-SO₂-,
5 aryl-C₁₋₆-alkylene-SO₂-, aryl-SO₂-, SO₂-, C₁₋₆-alkyl-C(O)-, aryl-C₁₋₆-alkylene-C(O)-,
N(R¹³)(R¹⁴)-C₁₋₆-alkylene-Y-, and R¹⁵-W²-C₁₋₆-alkylene-Y-, wherein




Y and W² independently of each other are a direct bond, -CH₂-, -SO₂-,
-N(H)CO-, -N(H)SO₂-, or -O-C(O)-;

R¹³ and R¹⁴ independently of each other are selected from hydrogen, aryl,
10 C₁₋₆-alkyl, or aryl-C₁₋₆-alkylene-;

or

R¹³ and R¹⁴ may be taken together to form a ring having the formula
-(CH₂)_o-X-(CH₂)_p- bonded to the nitrogen atom to which R¹³ and R¹⁴ are
attached, wherein

15 o and p are independently of each other are 1, 2, 3, or 4; and
X is a direct bond, -CH₂-, -O-, -S-, -S(O₂)-, -C(O)-, -CON(H)-,
-NHC(O)-, -NHCON(H)-, -NHSO₂-, -SO₂N(H)-, -C(O)-O-, -O-C(O)-,
-NHSO₂NH-,



20 wherein

R¹⁶ and R¹⁷ are selected from hydrogen, aryl, heteroaryl,
C₁₋₆-alkyl, C₁₋₆-alkoxy, aryl-C₁₋₆-alkylene-,
heteroaryl-C₁₋₆-alkylene-, C₁₋₆-alkyl-arylene-,
C₁₋₆-alkyl-heteroarylene-, C₁₋₆-alkoxy-arylene-,

C₁₋₆-alkoxy-heteroarylene-, heteroarylaryl-C₁₋₆-alkoxy-, or
aryl-C₁₋₆-alkoxy-; and

R¹⁵ is selected from the group consisting of aryl, heteroaryl, cycloalkyl, heterocyclyl,
C₁₋₆-alkyl, or aryl-C₁₋₆-alkylene-.

5 87. A compound according to claim 86, wherein

R¹¹ is selected from hydrogen, C₁₋₆-alkyl, aryl, carbamoyl, aryl-C₁₋₆-alkylene-,
C₁₋₆-alkyl-NH-C(O)-, aryl-C₁₋₆-alkylene-NH-C(O)-, C₁₋₆-alkyl-SO₂-,
aryl-C₁₋₆-alkylene-SO₂-, aryl-SO₂-, SO₂-, C₁₋₆-alkyl-C(O)-, aryl-C₁₋₆-alkylene-C(O)-,
N(R¹³)(R¹⁴)-C₁₋₆-alkylene-Y-, and R¹⁵-W²-C₁₋₆-alkylene-Y-, wherein

10 Y and W² independently of each other are a direct bond, -CH₂-, -SO₂-,
-N(H)CO-, -N(H)SO₂-, or -O-C(O)-;




R¹³ and R¹⁴ independently of each other are selected from hydrogen, aryl,
C₁₋₆-alkyl, or aryl-C₁₋₆-alkylene-;

or

15 R¹³ and R¹⁴ may be taken together to form a ring having the formula
-(CH₂)_o-X-(CH₂)_p- bonded to the nitrogen atom to which R¹³ and R¹⁴ are
attached, wherein

o and p are independently of each other are 1, 2, 3, or 4; and

20 X is a direct bond, -CH₂-, -O-, -S-, -S(O₂)-, -C(O)-, -CON(H)-,
-NHC(O)-, -NHCON(H)-, -NHSO₂-, -SO₂N(H)-, -C(O)-O-, -O-C(O)-,
-NHSO₂NH-,



wherein

R¹⁶ and R¹⁷ are hydrogen; and

R^{15} is selected from the group consisting of aryl, heteroaryl, cycloalkyl, heterocyclyl, C_{1-6} -alkyl, or aryl- C_{1-6} -alkylene-.

88. A compound according to claim 87, wherein

R^{11} is selected from hydrogen, C_{1-6} -alkyl, aryl, carbamoyl, aryl- C_{1-6} -alkylene-,
 5 C_{1-6} -alkyl-NH-C(O)-, aryl- C_{1-6} -alkylene-NH-C(O)-, C_{1-6} -alkyl-SO₂-,
 aryl- C_{1-6} -alkylene-SO₂-, aryl-SO₂-, SO₂-, C_{1-6} -alkyl-C(O)-, aryl- C_{1-6} -alkylene-C(O)-,
 N(R^{13})(R^{14})- C_{1-6} -alkylene-Y-, and R^{15} -W²- C_{1-6} -alkylene-Y-, wherein
 Y and W² independently of each other are a direct bond, -CH₂-, -SO₂-,
 -N(H)CO-, -N(H)SO₂-, or -O-C(O)-;
 10 R^{13} and R^{14} independently of each other are selected from hydrogen, aryl,
 C_{1-6} -alkyl, or aryl- C_{1-6} -alkylene-;
 or
 R^{13} and R^{14} may be taken together to form a ring having the formula
 -(CH₂)_o-X-(CH₂)_p- bonded to the nitrogen atom to which R^{13} and R^{14} are
 15 attached, wherein

o and p are independently of each other are 1, 2, 3, or 4;

X is a direct bond; and

R^{15} is selected from the group consisting of aryl, heteroaryl, cycloalkyl, heterocyclyl, C_{1-6} -alkyl, or aryl- C_{1-6} -alkylene-.

20 89. A compound according to claim 88, wherein

R^{11} is selected from hydrogen, C_{1-6} -alkyl, aryl, carbamoyl, aryl- C_{1-6} -alkylene-,
 C_{1-6} -alkyl-NH-C(O)-, aryl- C_{1-6} -alkylene-NH-C(O)-, C_{1-6} -alkyl-SO₂-,
 aryl- C_{1-6} -alkylene-SO₂-, aryl-SO₂-, SO₂-, C_{1-6} -alkyl-C(O)-, aryl- C_{1-6} -alkylene-C(O)-,
 N(R^{13})(R^{14})- C_{1-6} -alkylene-Y-, and R^{15} -W²- C_{1-6} -alkylene-Y-, wherein
 25 Y and W² independently of each other are a direct bond, -CH₂-, -SO₂-,
 -N(H)CO-, -N(H)SO₂-, or -O-C(O)-;
 R^{13} and R^{14} independently of each other are selected from hydrogen, aryl,
 C_{1-6} -alkyl, or aryl- C_{1-6} -alkylene-; and
 R^{15} is selected from the group consisting of aryl, heteroaryl, cycloalkyl,
 30 heterocyclyl, C_{1-6} -alkyl, or aryl- C_{1-6} -alkylene-.

90. A compound according to claim 89, wherein

R¹¹ is selected from hydrogen, C₁₋₆-alkyl, aryl, carbamoyl, aryl-C₁₋₆-alkylene-, C₁₋₆-alkyl-NH-C(O)-, aryl-C₁₋₆-alkylene-NH-C(O)-, C₁₋₆-alkyl-SO₂-, aryl-C₁₋₆-alkylene-SO₂-, aryl-SO₂-, SO₂-, C₁₋₆-alkyl-C(O)-, aryl-C₁₋₆-alkylene-C(O)-, N(R¹³)(R¹⁴)-C₁₋₆-alkylene-Y-, and R¹⁵-W²-C₁₋₆-alkylene-Y-, wherein

- 5 Y and W² independently of each other are a direct bond, -CH₂-, -SO₂-, -N(H)CO-, -N(H)SO₂-, or -O-C(O)-;
 R¹³ and R¹⁴ are hydrogen; and
 R¹⁵ is selected from the group consisting of aryl, heteroaryl, cycloalkyl, heterocyclyl, C₁₋₆-alkyl, or aryl-C₁₋₆-alkylene-.

- 10 91. A compound according to claim 90, wherein

R¹¹ is selected from hydrogen, C₁₋₆-alkyl, aryl, carbamoyl, aryl-C₁₋₆-alkylene-, C₁₋₆-alkyl-NH-C(O)-, aryl-C₁₋₆-alkylene-NH-C(O)-, C₁₋₆-alkyl-SO₂-, aryl-C₁₋₆-alkylene-SO₂-, aryl-SO₂-, SO₂-, C₁₋₆-alkyl-C(O)-, aryl-C₁₋₆-alkylene-C(O)-, N(R¹³)(R¹⁴)-C₁₋₆-alkylene-Y-, and R¹⁵-W²-C₁₋₆-alkylene-Y-, wherein

- 15 Y is a direct bond;
 W² is a direct bond, -CH₂-, -SO₂-, -N(H)CO-, -N(H)SO₂-, or -O-C(O)-;
 R¹³ and R¹⁴ are hydrogen; and
 R¹⁵ is selected from the group consisting of aryl, heteroaryl, cycloalkyl, heterocyclyl, C₁₋₆-alkyl, or aryl-C₁₋₆-alkylene-.

- 20 92. A compound according to claim 90, wherein

R¹¹ is selected from hydrogen, C₁₋₆-alkyl, aryl, carbamoyl, aryl-C₁₋₆-alkylene-, C₁₋₆-alkyl-NH-C(O)-, aryl-C₁₋₆-alkylene-NH-C(O)-, C₁₋₆-alkyl-SO₂-, aryl-C₁₋₆-alkylene-SO₂-, aryl-SO₂-, SO₂-, C₁₋₆-alkyl-C(O)-, aryl-C₁₋₆-alkylene-C(O)-, N(R¹³)(R¹⁴)-C₁₋₆-alkylene-Y-, and R¹⁵-W²-C₁₋₆-alkylene-Y-, wherein

- 25 Y is a direct bond, -CH₂-, -SO₂-, -N(H)CO-, -N(H)SO₂-, or -O-C(O)-;
 W² is a direct bond;
 R¹³ and R¹⁴ are hydrogen; and
 R¹⁵ is selected from the group consisting of aryl, heteroaryl, cycloalkyl, heterocyclyl, C₁₋₆-alkyl, or aryl-C₁₋₆-alkylene-.

- 30 93. A compound according to any one of the claims 90 to 92, wherein

R¹¹ is selected from hydrogen, C₁₋₆-alkyl, aryl, carbamoyl, aryl-C₁₋₆-alkylene-, C₁₋₆-alkyl-NH-C(O)-, aryl-C₁₋₆-alkylene-NH-C(O)-, C₁₋₆-alkyl-SO₂-,

aryl-C₁₋₆-alkylene-SO₂-, aryl-SO₂-, SO₂-, C₁₋₆-alkyl-C(O)-, aryl-C₁₋₆-alkylene-C(O)-, NH₂-C₁₋₆-alkylene-, and R¹⁵-C₁₋₆-alkylene-, wherein

R¹⁵ is selected from the group consisting of aryl, heteroaryl, cycloalkyl, heterocyclyl, C₁₋₆-alkyl, or aryl-C₁₋₆-alkylene-.

5 94. A compound according to claim 93, wherein

R¹¹ is selected from hydrogen, C₁₋₆-alkyl, aryl, carbamoyl, aryl-C₁₋₆-alkylene-, C₁₋₆-alkyl-NH-C(O)-, aryl-C₁₋₆-alkylene-NH-C(O)-, C₁₋₆-alkyl-SO₂-, aryl-C₁₋₆-alkylene-SO₂-, aryl-SO₂-, SO₂-, C₁₋₆-alkyl-C(O)-, aryl-C₁₋₆-alkylene-C(O)-, and NH₂-C₁₋₆-alkylene-.

10 95. A compound according to claim 94, wherein

R¹¹ is hydrogen or C₁₋₆-alkyl.

96. A compound according to claim 95, wherein

R¹¹ is hydrogen.

97. A compound according to any one of the claims 1 to 96, wherein

15 R¹² is hydrogen or C₁₋₆-alkyl.

98. A compound according to claim 97, wherein

R¹² is hydrogen.

99. A compound according to any one of the claims 1 to 98, wherein

20 G¹ is C₁₋₆-alkyl, C₂₋₆-alkenyl, C₂₋₆-alkynyl, cycloalkyl or heterocyclyl, optionally substituted with one or more substituents selected from the group consisting of -CN, -CF₃, -OCF₃, -OR¹⁸, -NR¹⁸R¹⁹, C₃₋₁₀-cycloalkyl and C₁₋₆-alkyl, wherein

R¹⁸ and R¹⁹ are as defined in claim 1.

100. A compound according to claim 99, wherein

25 G¹ is C₁₋₆-alkyl, C₂₋₆-alkenyl, C₂₋₆-alkynyl, cycloalkyl or heterocyclyl, optionally substituted with one or more substituents selected from the group consisting of -CN, -CF₃, -OCF₃, -OR¹⁸, -NR¹⁸R¹⁹ and C₁₋₆-alkyl, wherein

R¹⁸ and R¹⁹ are as defined in claim 1.

101. A compound according to claim 99, wherein
G¹ is C₁₋₈-alkyl, C₂₋₆-alkenyl, C₂₋₆-alkynyl, C₃₋₁₀-cycloalkyl or C₃₋₁₀-heterocyclyl,
optionally substituted with one or more substituents selected from the group
consisting of -CN, -CF₃, -OCF₃, -OR¹⁸, -NR¹⁸R¹⁹, C₃₋₁₀-cycloalkyl and C₁₋₈-alkyl,
5 wherein
R¹⁸ and R¹⁹ are as defined in claim 1.
102. A compound according to claim 100 or claim 101, wherein
G¹ is C₁₋₈-alkyl, C₂₋₆-alkenyl, C₂₋₆-alkynyl, C₃₋₁₀-cycloalkyl or C₃₋₁₀-heterocyclyl,
optionally substituted with one or more substituents selected from the group
10 consisting of -CN, -CF₃, -OCF₃, -OR¹⁸, -NR¹⁸R¹⁹ and C₁₋₈-alkyl, wherein
R¹⁸ and R¹⁹ are as defined in claim 1.
103. A compound according to any one of the claims 99 to 102, wherein
G¹ is selected from the group consisting of methyl, ethyl, propyl, butyl, isopropyl,
isobutyl, *sec*-butyl, *tert*-butyl, 3-pentyl, 2-pentyl, 3-methyl-butyl, 2-propenyl,
15 cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, cycloheptyl, oxetanyl,
tetrahydrofuranyl, tetrahydropyranyl, azetidyl, pyrrolidyl, piperidyl,
hexahydroazepinyl, thiolanyl, tetrahydrothiopyranyl, thiepanyl, 1,4-oxathianyl, 1,3-
dioxolanyl, 1,2-dithiolanyl, 1,3-dithiolanyl, hexahydro-pyridazinyl, imidazolidyl, 1,3-
dioxanyl, morpholinyl, 1,3-dithianyl, 1,4-dioxanyl, 1,4-dithianyl, or thiomorpholinyl.
- 20 104. A compound according to claim 103 wherein
G¹ is selected from the group consisting of methyl, ethyl, propyl, butyl, isopropyl,
isobutyl, *sec*-butyl, *tert*-butyl, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl,
cycloheptyl, tetrahydrofuranyl, tetrahydropyranyl, pyrrolidyl, piperidyl,
hexahydroazepinyl, thiolanyl, tetrahydrothiopyranyl, or thiepanyl .
- 25 105. A compound according to claim 104 wherein
G¹ is selected from the group consisting of methyl, ethyl, propyl, butyl, isopropyl,
isobutyl, cyclopentyl, cyclohexyl, tetrahydrofuranyl, tetrahydropyranyl, piperidyl, or
hexahydroazepinyl.
106. A compound according to claim 105 wherein
30 G¹ is selected from the group consisting of isobutyl, cyclopentyl, and piperidyl.
107. A compound according to claim 105 wherein
G¹ is isobutyl.
108. A compound according to claim 105 wherein
G¹ is cyclopentyl.

109. A compound according to claim 105 wherein
G¹ is piperidyl.
110. A compound according to any one of the claims 99 to 102, wherein
R¹⁸ and R¹⁹, independently of each other, are hydrogen, C₁₋₆-alkyl,
5 heteroaryl-C₁₋₆-alkylene-, aryl-C₁₋₆-alkylene-, C₁₋₆-alkyl-arylene-,
C₁₋₆-alkyl-heteroarylene-, heteroaryl, or aryl.
111. A compound according to claim 110, wherein
R¹⁸ and R¹⁹, independently of each other, are hydrogen, C₁₋₆-alkyl,
C₃₋₁₀-heteroaryl-C₁₋₆-alkylene-, C₃₋₁₀-aryl-C₁₋₆-alkylene-, C₁₋₆-alkyl-C₃₋₁₀-arylene-,
10 C₁₋₆-alkyl-C₃₋₁₀-heteroarylene-, C₃₋₁₀-heteroaryl, or C₃₋₁₀-aryl.
112. A compound according to claim 111, wherein
R¹⁸ and R¹⁹, independently of each other, are hydrogen or C₁₋₆-alkyl.
113. A compound according to claim 112, wherein
R¹⁸ is hydrogen.
- 15 114. A compound according to claim 112 or 113, wherein
R¹⁹ is hydrogen.
115. A compound according to claim 100 or claim 102, wherein
R¹⁸ and R¹⁹, when attached to the same nitrogen atom, together with the said
nitrogen atom forms a 3 to 8 membered heterocyclic ring optionally containing one
20 or two further heteroatoms selected from nitrogen, oxygen and sulfur, and optionally
containing one or two double bonds.
116. A compound according to any one of the claims 1 to 98, wherein
G¹ is alkyl or cycloalkyl, optionally substituted with one or more substituents
selected from the group consisting of -CN, -CF₃, -OCF₃, -OR¹⁸, -NR¹⁸R¹⁹ and
25 C₁₋₆-alkyl,
or G¹ is aryl optionally substituted with one or more substituents R⁴⁰, R⁴¹, and R⁴²,
wherein R¹⁸, R¹⁹, R⁴⁰, R⁴¹, and R⁴² are as defined in claim 1.

117. A compound according to claim 116, wherein

G¹ is C₁₋₆-alkyl or C₃₋₁₀-cycloalkyl, optionally substituted with one or more substituents selected from the group consisting of -CN, -CF₃, -OCF₃, -OR¹⁸, -NR¹⁸R¹⁹ and C₁₋₆-alkyl,

5 or G¹ is C₃₋₁₀-aryl optionally substituted with one or more substituents R⁴⁰, R⁴¹, and R⁴², wherein R¹⁸, R¹⁹, R⁴⁰, R⁴¹, and R⁴² are as defined in claim 1.

118. A compound according to claim 117, wherein

G¹ is C₁₋₆-alkyl or C₃₋₁₀-cycloalkyl, optionally substituted with one or more substituents selected from the group consisting of -CN, -CF₃, -OCF₃, -OR¹⁸, -NR¹⁸R¹⁹ and C₁₋₆-alkyl,

10 or G¹ is phenyl optionally substituted with one or more substituents R⁴⁰, R⁴¹, and R⁴², wherein R¹⁸, R¹⁹, R⁴⁰, R⁴¹, and R⁴² are as defined in claim 1.

119. A compound according to claim 116, 117, or 118, wherein

15 R¹⁸ and R¹⁹, independently of each other, are hydrogen, C₁₋₆-alkyl, heteroaryl-C₁₋₆-alkylene-, aryl-C₁₋₆-alkylene-, C₁₋₆-alkyl-arylene-, C₁₋₆-alkyl-heteroarylene-, heteroaryl, or aryl.

120. A compound according to claim 119, wherein

20 R¹⁸ and R¹⁹, independently of each other, are hydrogen, C₁₋₆-alkyl, C₃₋₁₀-heteroaryl-C₁₋₆-alkylene-, C₃₋₁₀-aryl-C₁₋₆-alkylene-, C₁₋₆-alkyl-C₃₋₁₀-arylene-, C₁₋₆-alkyl-C₃₋₁₀-heteroarylene-, C₃₋₁₀-heteroaryl, or C₃₋₁₀-aryl.

121. A compound according to claim 120, wherein

R¹⁸ and R¹⁹, independently of each other, are hydrogen or C₁₋₆-alkyl.

122. A compound according to claim 121, wherein

R¹⁸ is hydrogen.

25 123. A compound according to claim 121 or 122, wherein

R¹⁹ is hydrogen.

124. A compound according to claim 116 or claim 117, wherein

R¹⁸ and R¹⁹, when attached to the same nitrogen atom, together with the said nitrogen atom forms a 3 to 8 membered heterocyclic ring optionally containing one

or two further heteroatoms selected from nitrogen, oxygen and sulfur, and optionally containing one or two double bonds.

125. A compound according to any one of the claims 116 to 124, wherein

- 5 R^{40} , R^{41} , and R^{42} independently of each other are
 halogen, -CN, -NO₂, C₁₋₆-alkyl, -CHF₂, -CF₃, -OR⁵⁴, -NR⁵⁴R⁵⁵, -SR⁵⁴, -NR⁵⁴S(O)₂R⁵⁵,
 -S(O)₂NR⁵⁴R⁵⁵, -S(O)NR⁵⁴R⁵⁵, -S(O)R⁵⁴, -S(O)₂R⁵⁴, -C(O)NR⁵⁴R⁵⁵, -OC(O)NR⁵⁴R⁵⁵,
 -NR⁵⁴C(O)R⁵⁵, -CH₂C(O)NR⁵⁴R⁵⁵, -CH₂C(O)OR⁵⁴, -OCH₂C(O)NR⁵⁴R⁵⁵, -CH₂OR⁵⁴,
 -CH₂NR⁵⁴R⁵⁵, and -C(O)OR⁵⁴; or
 C₂₋₆-alkenyl and C₂₋₆-alkynyl, which may optionally be substituted with one or more
 10 substituents selected from -CN, -CF₃, -OCF₃, -OR⁵⁴, -NR⁵⁴R⁵⁵ and C₁₋₆-alkyl,
 wherein

- R^{54} and R^{55} independently of each other are hydrogen, C₁₋₆-alkyl,
 C₁₋₆-alkyl-arylene-, C₁₋₆-alkyl-heteroarylene-, aryl-C₁₋₆-alkylene-,
 heteroaryl-C₁₋₆-alkylene-, heteroaryl, or aryl;
 15 or
 R^{54} and R^{55} , when attached to the same nitrogen atom, together with the
 said nitrogen atom may form a 3 to 8 membered heterocyclic ring optionally
 containing one or two further heteroatoms selected from nitrogen, oxygen
 and sulfur, and optionally containing one or two double bonds.

20 126. A compound according to claim 125, wherein

- R^{40} , R^{41} , and R^{42} independently of each other are
 halogen, -CN, C₁₋₆-alkyl, -CF₃, -OR⁵⁴, -NR⁵⁴R⁵⁵, -SR⁵⁴, -NR⁵⁴S(O)₂R⁵⁵, -S(O)R⁵⁴,
 -S(O)₂R⁵⁴, -CH₂C(O)OR⁵⁴, -CH₂OR⁵⁴, -CH₂NR⁵⁴R⁵⁵, and -C(O)OR⁵⁴;
 wherein

- 25 R^{54} and R^{55} independently of each other are hydrogen, C₁₋₆-alkyl,
 C₁₋₆-alkyl-arylene-, C₁₋₆-alkyl-heteroarylene-, aryl-C₁₋₆-alkylene-,
 heteroaryl-C₁₋₆-alkylene-, heteroaryl, or aryl;
 or
 R^{54} and R^{55} , when attached to the same nitrogen atom, together with the
 30 said nitrogen atom may form a 3 to 8 membered heterocyclic ring optionally
 containing one or two further heteroatoms selected from nitrogen, oxygen
 and sulfur, and optionally containing one or two double bonds.

127. A compound according to claim 126 wherein

R^{40} , R^{41} , and R^{42} independently of each other are

halogen, or -OR⁵⁴

wherein

R⁵⁴ is hydrogen, C₁₋₆-alkyl, C₁₋₆-alkyl-arylene-, C₁₋₆-alkyl-heteroarylene-, aryl-C₁₋₆-alkylene-, heteroaryl-C₁₋₆-alkylene-, heteroaryl, or aryl.

5 128. A compound according to claim 126, wherein

R⁵⁴ and R⁵⁵ independently of each other are hydrogen, or C₁₋₆-alkyl, C₁₋₆-alkyl-C₃₋₁₀-arylene-, C₁₋₆-alkyl-C₃₋₁₀-heteroarylene-, C₃₋₁₀-aryl-C₁₋₆-alkylene-, C₃₋₁₀-heteroaryl-C₁₋₆-alkylene-, C₃₋₁₀-heteroaryl, or C₃₋₁₀-aryl.

129. A compound according to claim 128, wherein

10 R⁵⁴ and R⁵⁵ independently of each other are hydrogen or C₁₋₆-alkyl.

130. A compound according to claim 129, wherein

R⁵⁴ is hydrogen.

131. A compound according to claim 129 or claim 130, wherein

R⁵⁵ is hydrogen.

15 132. A compound according to claim 127 wherein

R⁵⁴ is methyl.

133. A compound according to any one of the claims 1 to 98, wherein

G¹ is aryl or heteroaryl, optionally substituted with one or more substituents R⁴⁰, R⁴¹, and R⁴², wherein

20 R⁴⁰, R⁴¹, and R⁴² are as defined in claim 1.

134. A compound according to claim 133, wherein

G¹ is C₃₋₁₀-aryl or C₃₋₁₀-heteroaryl, optionally substituted with one or more substituents R⁴⁰, R⁴¹, and R⁴², wherein

R⁴⁰, R⁴¹, and R⁴² are as defined in claim 1.

25 135. A compound according to claim 133, wherein

G¹ is aryl, optionally substituted with one or more substituents R⁴⁰, R⁴¹, and R⁴², wherein

R⁴⁰, R⁴¹, and R⁴² are as defined in claim 1.

136. A compound according to claim 135, wherein

G¹ is C₃₋₁₀-aryl, optionally substituted with one or more substituents R⁴⁰, R⁴¹, and R⁴², wherein

R⁴⁰, R⁴¹, and R⁴² are as defined in claim 1.

5 137. A compound according to claim 136, wherein

G¹ is phenyl, optionally substituted with one or more substituents R⁴⁰, R⁴¹, and R⁴², wherein

R⁴⁰, R⁴¹, and R⁴² are as defined in claim 1.

138. A compound according to any one of the claims 133 to 137, wherein

10 R⁴⁰, R⁴¹, and R⁴² independently of each other are

halogen, -CN, -NO₂, C₁₋₆-alkyl, -CHF₂, -CF₃, -OR⁵⁴, -NR⁵⁴R⁵⁵, -SR⁵⁴, -NR⁵⁴S(O)₂R⁵⁵, -S(O)₂NR⁵⁴R⁵⁵, -S(O)NR⁵⁴R⁵⁵, -S(O)R⁵⁴, -S(O)₂R⁵⁴, -C(O)NR⁵⁴R⁵⁵, -OC(O)NR⁵⁴R⁵⁵, -NR⁵⁴C(O)R⁵⁵, -CH₂C(O)NR⁵⁴R⁵⁵, -CH₂C(O)OR⁵⁴, -OCH₂C(O)NR⁵⁴R⁵⁵, -CH₂OR⁵⁴, -CH₂NR⁵⁴R⁵⁵, and -C(O)OR⁵⁴; or

15 C₂₋₆-alkenyl and C₂₋₆-alkynyl, which may optionally be substituted with one or more substituents selected from -CN, -CF₃, -OCF₃, -OR⁵⁴, -NR⁵⁴R⁵⁵ and C₁₋₆-alkyl, wherein

R⁵⁴ and R⁵⁵ independently of each other are hydrogen, C₁₋₆-alkyl,

C₁₋₆-alkyl-arylene-, C₁₋₆-alkyl-heteroarylene-, aryl-C₁₋₆-alkylene-,

20 heteroaryl-C₁₋₆-alkylene-, heteroaryl, or aryl;

or

R⁵⁴ and R⁵⁵, when attached to the same nitrogen atom, together with the said nitrogen atom may form a 3 to 8 membered heterocyclic ring optionally containing one or two further heteroatoms selected from nitrogen, oxygen, and sulfur, and optionally containing one or two double bonds.

25

139. A compound according to claim 138, wherein

R⁴⁰, R⁴¹, and R⁴² independently of each other are

halogen, -CN, C₁₋₆-alkyl, -CF₃, -OR⁵⁴, -NR⁵⁴R⁵⁵, -SR⁵⁴, -NR⁵⁴S(O)₂R⁵⁵, -S(O)R⁵⁴, -S(O)₂R⁵⁴, -CH₂C(O)OR⁵⁴, -CH₂OR⁵⁴, -CH₂NR⁵⁴R⁵⁵, and -C(O)OR⁵⁴;

30

wherein

R⁵⁴ and R⁵⁵ independently of each other are hydrogen, C₁₋₆-alkyl,

C₁₋₆-alkyl-arylene-, C₁₋₆-alkyl-heteroarylene-, aryl-C₁₋₆-alkylene-,

heteroaryl-C₁₋₆-alkylene-, heteroaryl, or aryl;

or

R⁵⁴ and R⁵⁵, when attached to the same nitrogen atom, together with the said nitrogen atom may form a 3 to 8 membered heterocyclic ring optionally containing one or two further heteroatoms selected from nitrogen, oxygen and sulfur, and optionally containing one or two double bonds.

5

140. A compound according to claim 139 wherein

R⁴⁰, R⁴¹, and R⁴² independently of each other are halogen, or -OR⁵⁴

wherein

10

R⁵⁴ is hydrogen, C₁₋₆-alkyl, C₁₋₆-alkyl-aryl-, C₁₋₆-alkyl-heteroaryl-, aryl-C₁₋₆-alkylene-, heteroaryl-C₁₋₆-alkylene-, heteroaryl, or aryl.

141. A compound according to any one of the claims 138 to 139, wherein

R⁵⁴ and R⁵⁵ independently of each other are hydrogen, or C₁₋₆-alkyl, C₁₋₆-alkyl-C₃₋₁₀-aryl-, C₁₋₆-alkyl-C₃₋₁₀-heteroaryl-, C₃₋₁₀-aryl-C₁₋₆-alkylene-, C₃₋₁₀-heteroaryl-C₁₋₆-alkylene-, C₃₋₁₀-heteroaryl, or C₃₋₁₀-aryl.

15

142. A compound according to claim 141, wherein

R⁵⁴ and R⁵⁵ independently of each other are hydrogen or C₁₋₆-alkyl.

143. A compound according to claim 142, wherein

R⁵⁴ is hydrogen.

20

144. A compound according to claim 142 or claim 143, wherein

R⁵⁵ is hydrogen.


145. A compound according to claim 140 wherein

R⁵⁴ is methyl.

146. A compound according to claim 135, wherein

25

G¹ is



, wherein

- R⁵⁶, R⁵⁷, R⁵⁸, R⁵⁹, and R⁶⁰, independently of each other, are hydrogen or halogen, -CN, -NO₂, C₁₋₆-alkyl, -CHF₂, -CF₃, -OCF₃, -OCHF₂, -OCH₂CF₃, -OCF₂CHF₂, -S(O)₂CF₃, -SCF₃, -OR⁶¹, -NR⁶¹R⁶², -SR⁶¹, -NR⁶¹S(O)₂R⁶², -S(O)₂NR⁶¹R⁶², -S(O)NR⁶¹R⁶², -S(O)R⁶¹, -S(O)₂R⁶¹, -C(O)NR⁶¹R⁶², -OC(O)NR⁶¹R⁶², -NR⁶¹C(O)R⁶², -CH₂C(O)NR⁶¹R⁶², -CH₂C(O)OR⁶¹, -OCH₂C(O)NR⁶¹R⁶², -CH₂OR⁶¹, -CH₂NR⁶¹R⁶², -OC(O)R⁶¹, -C(O)R⁶¹ and -C(O)OR⁶¹; or C₂₋₆-alkenyl and C₂₋₆-alkynyl, which may optionally be substituted with one or more substituents selected from -CN, -CF₃, -OCF₃, -OR⁶¹, -NR⁶¹R⁶² and C₁₋₆-alkyl; C₃₋₁₀-cycloalkyl, C₄₋₈-cycloalkenyl, heterocyclyl, C₃₋₁₀-cycloalkyl-C₁₋₆-alkylene-, C₃₋₁₀-cycloalkyl-C₁₋₆-alkoxy-, C₃₋₁₀-cycloalkyloxy, C₃₋₁₀-cycloalkyl-C₁₋₆-alkylthio-, C₃₋₁₀-cycloalkylthio, C₃₋₁₀-cycloalkyl-C₂₋₆-alkenylene-, C₃₋₁₀-cycloalkyl-C₂₋₆-alkynylene-, C₄₋₈-cycloalkenyl-C₁₋₆-alkylene-, C₄₋₈-cycloalkenyl-C₂₋₆-alkenylene-, C₄₋₈-cycloalkenyl-C₂₋₆-alkynylene-, heterocyclyl-C₁₋₆-alkylene-, heterocyclyl-C₂₋₆-alkenylene-, heterocyclyl-C₂₋₆-alkynylene-; or aryl, aryloxy, aryloxy carbonyl, aroyl, aryl-C₁₋₆-alkoxy-, aryl-C₁₋₆-alkylene-, aryl-C₂₋₆-alkenylene-, aryl-C₂₋₆-alkynylene-, heteroaryl, heteroaryl-C₁₋₆-alkylene-, heteroaryl-C₂₋₆-alkenylene- and heteroaryl-C₂₋₆-alkynylene-, of which the aryl and heteroaryl moieties optionally may be substituted with one or more substituents selected from halogen, -C(O)OR⁶¹, -CN, -CF₃, -OCF₃, -NO₂, -OR⁶¹, -NR⁶¹R⁶² or C₁₋₆-alkyl, wherein
- R⁶¹ and R⁶² independently of each other are hydrogen, C₁₋₆-alkyl, C₁₋₆-alkyl-arylene-, C₁₋₆-alkyl-heteroarylene-, aryl-C₁₋₆-alkylene-, heteroaryl-C₁₋₆-alkylene-, heteroaryl, or aryl;
- or
- R⁶¹ and R⁶², when attached to the same nitrogen atom, together with the said nitrogen atom may form a 3 to 8 membered heterocyclic ring optionally containing one or two further heteroatoms selected from nitrogen, oxygen and sulfur, and optionally containing one or two double bonds.

147. A compound according to claim 146, wherein

R⁵⁶, R⁵⁷, R⁵⁸, R⁵⁹, and R⁶⁰ independently of each other are

halogen, -CN, -NO₂, C₁₋₆-alkyl, -CHF₂, -CF₃, -OR⁶¹, -NR⁶¹R⁶², -SR⁶¹, -NR⁶¹S(O)₂R⁶², -S(O)₂NR⁶¹R⁶², -S(O)NR⁶¹R⁶², -S(O)R⁶¹, -S(O)₂R⁶¹, -C(O)NR⁶¹R⁶², -OC(O)NR⁶¹R⁶², -NR⁶¹C(O)R⁶², -CH₂C(O)NR⁶¹R⁶², -CH₂C(O)OR⁶¹, -OCH₂C(O)NR⁶¹R⁶², -CH₂OR⁶¹, -CH₂NR⁶¹R⁶², and -C(O)OR⁶¹; or

- 5 C₂₋₆-alkenyl and C₂₋₆-alkynyl, which may optionally be substituted with one or more substituents selected from -CN, -CF₃, -OCF₃, -OR⁶¹, -NR⁶¹R⁶² and C₁₋₆-alkyl, wherein

- R⁶¹ and R⁶² independently of each other are hydrogen, C₁₋₆-alkyl, C₁₋₆-alkyl-arylene-, C₁₋₆-alkyl-heteroarylene-, aryl-C₁₋₆-alkylene-, heteroaryl-C₁₋₆-alkylene-, heteroaryl, or aryl;
- 10 or
- R⁶¹ and R⁶², when attached to the same nitrogen atom, together with the said nitrogen atom may form a 3 to 8 membered heterocyclic ring optionally containing one or two further heteroatoms selected from nitrogen, oxygen
- 15 and sulfur, and optionally containing one or two double bonds.

148. A compound according to claim 146 or claim 147, wherein R⁵⁶ or R⁵⁷ is -OR⁶¹.

149. A compound according to claim 148 wherein R⁶¹ is methyl.

150. A compound according to claim 147, wherein

- R⁶¹ and R⁶² independently of each other are hydrogen, or C₁₋₆-alkyl, C₁₋₆-alkyl-C₃₋₁₀-arylene-, C₁₋₆-alkyl-C₃₋₁₀-heteroarylene-, C₃₋₁₀-aryl-C₁₋₆-alkylene-, C₃₋₁₀-heteroaryl-C₁₋₆-alkylene-, C₃₋₁₀-heteroaryl, or C₃₋₁₀-aryl.
- 20

151. A compound according to claim 150, wherein

R⁶¹ and R⁶² independently of each other are hydrogen or C₁₋₆-alkyl.

152. A compound according to claim 151, wherein

- 25 R⁶¹ is hydrogen.

153. A compound according to claim 151 or claim 152, wherein

R⁶² is hydrogen.

154. A compound according to any one of the claims 146 to 153, wherein

at least one of R^{56} , R^{57} , R^{58} , R^{59} , and R^{60} are hydrogen.

155. A compound according to claim 154, wherein
at least two of R^{23} , R^{24} , R^{25} , and R^{26} are hydrogen.

5 156. A compound according to claim 155, wherein
at least three of R^{23} , R^{24} , R^{25} , and R^{26} are hydrogen.

157. A compound according to any one of the claims 1 to 156, wherein
 L^2 is a direct bond, C_{1-6} -alkylene, C_{2-6} -alkenylene, C_{2-6} -alkynylene, $-N-R^{20}-$,
 $-C_{1-6}$ -alkylene- $N(R^{20})-$, $-C_{2-6}$ -alkenylene- $N(R^{20})-$, or $-C_{2-6}$ -alkynylene- $N(R^{20})-$, wherein
 R^{20} is as defined in claim 1.

10 158. A compound according to any one of the claims 1 to 153, wherein
 L^2 is $-N-R^{20}-$, $-alkylene-N(R^{20})-$, $-alkenylene-N(R^{20})-$, or $-alkynylene-N(R^{20})-$, wherein
 R^{20} is as defined in claim 1.

15 159. A compound according to claim 157 or claim 158, wherein
 L^2 is $-N-R^{20}-$, $-C_{1-6}$ -alkylene- $N(R^{20})-$, $-C_{2-6}$ -alkenylene- $N(R^{20})-$, or
 $-C_{2-6}$ -alkynylene- $N(R^{20})-$, wherein
 R^{20} is as defined in claim 1.

160. A compound according to claim 159, wherein
 L^2 is $-N-R^{20}-$, wherein
 R^{20} is as defined in claim 1.

20 161. A compound according to claim 157, wherein
 L^2 is a direct bond.

25 162. A compound according to any one of the claims 1 to 160, wherein
 R^{20} is hydrogen, or
 R^{20} is C_{1-6} -alkyl, C_{2-6} -alkenyl, C_{2-6} -alkynyl, C_{3-10} -cycloalkyl- W^3- ,
 C_{3-10} -heterocyclyl- W^3- , C_{3-10} -aryl- W^3- , or C_{4-10} -heteroaryl- W^3- , optionally substituted
with one or more substituents R^{30} , R^{31} , and R^{32} , wherein
 W^3 , R^{30} , R^{31} , and R^{32} are as defined in claim 1.

163. A compound according to claim 162, wherein
W³ is alkylene.
164. A compound according to claim 163, wherein
W³ is C₂₋₆-alkylene.
- 5 165. A compound according to claim 162, wherein
W³ is a direct bond.
166. A compound according to any one of the claims 1 to 160, wherein
R²⁰ is hydrogen, C₁₋₆-alkyl, C₂₋₆-alkenyl, or C₂₋₆-alkynyl, optionally substituted with
one or more substituents R³⁰, R³¹, and R³², wherein
10 R³⁰, R³¹, and R³² are as defined in claim 1.
167. A compound according to any one of the claims 162 to 166, wherein
R²⁰ is hydrogen, or
R²⁰ is C₁₋₆-alkyl, C₁₋₆-alkenyl, or C₁₋₆-alkynyl, optionally substituted with one or more
substituents R³⁰, R³¹, and R³², wherein
15 R³⁰, R³¹, and R³² are as defined in claim 1.
168. A compound according to any one of the claims 1 to 167, wherein
R³⁰, R³¹, and R³² independently of each other are selected from -CHF₂, -CF₃, -OCF₃,
-OCHF₂, -OCH₂CF₃, -OCF₂CHF₂, -S(O)₂CF₃, -SCF₃, -OR⁵², -NR⁵²R⁵³, -SR⁵²,
-NR⁵²S(O)₂R⁵³, -S(O)₂NR⁵²R⁵³, -S(O)NR⁵²R⁵³, -S(O)R⁵², -S(O)₂R⁵², -C(O)NR⁵²R⁵³,
20 -OC(O)NR⁵²R⁵³, -NR⁵²C(O)R⁵³, -CH₂C(O)NR⁵²R⁵³, -OCH₂C(O)NR⁵²R⁵³, -CH₂OR⁵²,
-CH₂NR⁵²R⁵³, -OC(O)R⁵², -C(O)R⁵² and -C(O)OR⁵², wherein
R⁵² and R⁵³ are as defined in claim 1.
169. A compound according to claim 168, wherein
R⁵² and R⁵³, when attached to the same nitrogen atom, together with the said
25 nitrogen atom may form a 3 to 8 membered heterocyclic ring optionally containing
one or two further heteroatoms selected from nitrogen, oxygen and sulfur, and
optionally containing one or two double bonds.
170. A compound according to any one of the claims 1 to 168, wherein

R^{52} and R^{53} independently of each other are hydrogen, C_{1-6} -alkyl, aryl- C_{1-6} -alkylene-heteroaryl- C_{1-6} -alkylene- heteroaryl, or aryl.

171. A compound according to claim 170, wherein

5 R^{52} and R^{53} independently of each other are hydrogen, C_{1-6} -alkyl, aryl- C_{1-6} -alkylene-or aryl.

172. A compound according to claim 171, wherein

R^{52} and R^{53} independently of each other are hydrogen or C_{1-6} -alkyl.

173. A compound according to claim 172, wherein

R^{52} is hydrogen.

10 174. A compound according to claim 172 or claim 173, wherein

R^{53} is hydrogen.

175. A compound according to claim 167, wherein

R^{20} is hydrogen.

176. A compound according to any one of the claims 1 to 175, wherein

15 L^3 is -C(O)-, -C(O)-C(O)- or -C(O)CH₂C(O)-.

177. A compound according to claim 176 wherein

L^3 is -C(O)-.

178. A compound according to claim 176 wherein

L^3 is -C(O)-C(O)-.

20 179. A compound according to claim 176 wherein

L^3 is -C(O)CH₂C(O)-.

180. A compound according to any one of the claims 1 to 179, wherein

R^1 is hydrogen, or

25 R^1 is C_{1-6} -alkyl, C_{2-6} -alkenyl, C_{2-6} -alkynyl, C_{3-10} -cycloalkyl- W^4 -, C_{3-10} -heterocyclidyl- W^4 -, C_{3-10} -aryl- W^4 -, or C_{4-10} -heteroaryl- W^4 -, optionally substituted with one or more substituents R^{33} , R^{34} , and R^{35} , wherein

W^4 , R^{33} , R^{34} , and R^{35} are as defined in claim 1.

181. A compound according to claim 180, wherein

W^4 is alkylene.

182. A compound according to claim 181, wherein

5 W^4 is C_{2-6} -alkylene.

183. A compound according to claim 180, wherein

W^4 is a direct bond.

184. A compound according to any one of the claims 1 to 176, wherein

10 R^1 is hydrogen, C_{1-6} -alkyl, C_{2-6} -alkenyl, or C_{2-6} -alkynyl, optionally substituted with one or more substituents R^{33} , R^{34} , and R^{35} , wherein

R^{33} , R^{34} , and R^{35} are as defined in claim 1.

185. A compound according to any one of the claims 180 to 184, wherein

R^1 is hydrogen, C_{1-6} -alkyl, C_{2-6} -alkenyl, or C_{2-6} -alkynyl, optionally substituted with one or more substituents R^{33} , R^{34} , and R^{35} , wherein

15 R^{33} , R^{34} , and R^{35} are as defined in claim 1.

186. A compound according to claim 185, wherein

R^1 is hydrogen or C_{1-6} -alkyl optionally substituted with one or more substituents R^{33} , R^{34} , and R^{35} , wherein

R^{33} , R^{34} , and R^{35} are as defined in claim 1.

20 187. A compound according to any one of the claims 1 to 186, wherein

R^{33} , R^{34} , and R^{35} independently of each other are selected from $-\text{CHF}_2$, $-\text{CF}_3$, $-\text{OCF}_3$, $-\text{OCHF}_2$, $-\text{OCH}_2\text{CF}_3$, $-\text{OCF}_2\text{CHF}_2$, $-\text{S}(\text{O})_2\text{CF}_3$, $-\text{SCF}_3$, $-\text{OR}^{52}$, $-\text{NR}^{52}\text{R}^{53}$, $-\text{SR}^{52}$, $-\text{NR}^{52}\text{S}(\text{O})_2\text{R}^{53}$, $-\text{S}(\text{O})_2\text{NR}^{52}\text{R}^{53}$, $-\text{S}(\text{O})\text{NR}^{52}\text{R}^{53}$, $-\text{S}(\text{O})\text{R}^{52}$, $-\text{S}(\text{O})_2\text{R}^{52}$, $-\text{C}(\text{O})\text{NR}^{52}\text{R}^{53}$, $-\text{OC}(\text{O})\text{NR}^{52}\text{R}^{53}$, $-\text{NR}^{52}\text{C}(\text{O})\text{R}^{53}$, $-\text{CH}_2\text{C}(\text{O})\text{NR}^{52}\text{R}^{53}$, $-\text{OCH}_2\text{C}(\text{O})\text{NR}^{52}\text{R}^{53}$, $-\text{CH}_2\text{OR}^{52}$, $-\text{CH}_2\text{NR}^{52}\text{R}^{53}$, $-\text{OC}(\text{O})\text{R}^{52}$, $-\text{C}(\text{O})\text{R}^{52}$ and $-\text{C}(\text{O})\text{OR}^{52}$, wherein

25 R^{52} and R^{53} are as defined in claim 1.

188. A compound according to claim 186, wherein

R⁵² and R⁵³, when attached to the same nitrogen atom, together with the said nitrogen atom may form a 3 to 8 membered heterocyclic ring optionally containing one or two further heteroatoms selected from nitrogen, oxygen and sulfur, and optionally containing one or two double bonds.

5 189. A compound according to claim 186, wherein
R⁵² and R⁵³ independently of each other are hydrogen, C₁₋₆-alkyl, aryl-C₁₋₆-alkylene-
heteroaryl-C₁₋₆-alkylene- heteroaryl, or aryl.

190. A compound according to claim 189, wherein
R⁵² and R⁵³ independently of each other are hydrogen, C₁₋₆-alkyl,
10 aryl-C₁₋₆-alkylene-or aryl.

191. A compound according to claim 190, wherein
R⁵² and R⁵³ independently of each other are hydrogen or C₁₋₆-alkyl.

192. A compound according to claim 191, wherein
R⁵² is hydrogen.

15 193. A compound according to claim 191 or claim 192, wherein
R⁵³ is hydrogen.

194. A compound according to claim 186, wherein
R¹ is hydrogen.

195. A compound according to any one of the claims 1 to 194, wherein
20 G² is heteroaryl, fused heterocycl/heteroaryl, or fused cycloalkyl/heteroaryl,
optionally substituted with one or more substituents R⁴³, R⁴⁴, and R⁴⁵, wherein said
heteroaryl group possesses a nitrogen atom adjacent to the atom joining G² with
-N(R¹)-, and wherein
R⁴³, R⁴⁴, and R⁴⁵ are as defined in claim 1.

25 196. A compound according to claim 195, wherein

G² is fused cycloalkylheteroaryl optionally substituted with one or more substituents R⁴³, R⁴⁴, and R⁴⁵, wherein said heteroaryl group possesses a nitrogen atom adjacent to the atom joining G² with -N(R¹)-, and wherein

R⁴³, R⁴⁴, and R⁴⁵ are as defined in claim 1.

5

197. A compound according to claim 196 wherein

G² is fused cycloalkylthiazolyl optionally substituted with one or more substituents R⁴³, R⁴⁴, and R⁴⁵, and wherein

R⁴³, R⁴⁴, and R⁴⁵ are as defined in claim 1.

10

198. A compound according to claim 197, wherein

G² is fused cycloalkylthiazolyl optionally substituted with -COOH

199. A compound according to claim 195, wherein

G² is heteroaryl optionally substituted with one or more substituents R⁴³, R⁴⁴, and R⁴⁵, wherein said heteroaryl group possesses a nitrogen atom adjacent to the atom joining G² with -N(R¹)-, and wherein

R⁴³, R⁴⁴, and R⁴⁵ are as defined in claim 1.

15

200. A compound according to claim 199, wherein

G² is furanyl, thienyl, thiophenyl, pyrrolyl, imidazolyl, pyrazolyl, triazolyl, tetrazolyl, thiazolyl, oxazolyl, isoxazolyl, oxadiazolyl, thiadiazolyl, isothiazolyl, pyridinyl, pyridazinyl, pyrazinyl, pyrimidinyl, quinolinyl, isoquinolinyl, benzofuranyl, benzothiophenyl, indolyl, or indazolyl, optionally substituted with one or more substituents R⁴³, R⁴⁴, and R⁴⁵, wherein

R⁴³, R⁴⁴, and R⁴⁵ are as defined in claim 1.

20


201. A compound according to claim 200 wherein G² is thiazolyl, optionally substituted with one or more substituents R⁴³, R⁴⁴, and R⁴⁵, wherein

R⁴³, R⁴⁴, and R⁴⁵ are as defined in claim 1.

25

202. A compound according to claim 201 wherein G² is


577



wherein

R^{43} and R^{44} are as defined in claim 1.


203. A compound according to claim 202 wherein G^2 is



wherein

R^{43} is as defined in claim 1.


204. A compound according to claim 202 wherein G^2 is



wherein

R^{43} and R^{44} are as defined in claim 1.


205. A compound according to claim 200 wherein G^2 is



wherein

R^{43} is as defined in claim 1.

206. A compound according to claim 202 wherein G^2 is



wherein

R^{44} is as defined in claim 1.

207. A compound according to any one of the claims 1 to 206, wherein

R^{43} , R^{44} , and R^{45} independently of each other are selected from
 -CN, -NO₂, -SCF₃, -OR⁵⁴, -NR⁵⁴R⁵⁵, -SR⁵⁴, -S(O)₂NR⁵⁴R⁵⁵,
 -S(O)NR⁵⁴R⁵⁵, -S(O)R⁵⁴, -S(O)₂R⁵⁴, -C(O)NR⁵⁴R⁵⁵, -OC(O)NR⁵⁴R⁵⁵, -NR⁵⁴C(O)R⁵⁵,
 halogen, -S-C₁₋₆-alkylene-OR⁵⁴, -S(O)₂-C₁₋₆-alkylene-OR⁵⁴, -C₁₋₆-alkylene-S-R⁵⁴,
 -C₁₋₆-alkylene-S(O)R⁵⁴, -C₁₋₆-alkylene-S(O)₂R⁵⁴, -C₁₋₆-alkylene-N(R⁵⁴)S(O)₂R⁵⁵,
 -N(R⁵⁴)S(O)₂R⁵⁵, -C₁₋₆-alkylene-C(O)NR⁵⁴R⁵⁵, -C₁₋₆-alkylene-N(R⁵⁴)C(O)R⁵⁵,
 -N(R⁵⁴)C(O)R⁵⁵, -C₁₋₆-alkylene-N(R⁵⁴)C(O)NR⁵⁵R⁵⁶,
 C₁₋₆-alkylene-NHC(=NR⁵⁴)NR⁵⁵R⁵⁶, -C₁₋₆-alkylene-N(R⁵⁴)C(O)OR⁵⁵,
 -N(R⁵⁴)C(O)OR⁵⁵, -C₁₋₆-alkylene-C(O)OR⁵⁴, -OCH₂C(O)NR⁵⁴R⁵⁵, -O(CH₂)₁₋₃OR⁵⁴,
 -C₁₋₆-alkylene-O-R⁵⁴, -C₁₋₆-alkylene-C(O)R⁵⁴, -C₁₋₆-alkylene-NR⁵⁴R⁵⁵,
 -C₁₋₆-alkylene=N-O-R⁵⁴, -C₁₋₆-alkylene-N(R⁵⁴)S(O)₂NR⁵⁵R⁵⁶, -N(R⁵⁴)S(O)₂NR⁵⁵R⁵⁶,
 -N(R⁵⁴)S(O)₂NR⁵⁵R⁵⁶, -OC(O)R⁵⁴, -C(O)N(R⁵⁴)S(O)₂R⁵⁵,
 -C₁₋₆-alkylene-C(R⁵⁴)=N-OR⁵⁵, -NHC(=NR⁵⁴)NR⁵⁵R⁵⁶,
 -C₁₋₆-alkylene-N=C(N(R⁵⁴R⁵⁵))₂, -N=C(N(R⁵⁴R⁵⁵))₂, -C(O)R⁵⁴ and -C(O)OR⁵⁴; or
 C₁₋₆-alkyl or C₂₋₆-alkenyl, each of which may optionally be substituted with one or
 more substituents independently selected from halogen, R⁵⁴, -CN, -CF₃, -OCF₃, -
 OR⁵⁴, -C(O)OR⁵⁴, -NR⁵⁴R⁵⁵ and C₁₋₆-alkyl; or
 C₃₋₁₀-cycloalkyl, heterocyclyl, C₃₋₁₀-cycloalkyl-C₁₋₆-alkylene-,
 C₃₋₁₀-cycloalkyl-C₁₋₆-alkoxy-, C₃₋₁₀-cycloalkyloxy, heterocyclyl-C₁₋₆-alkylene-, of which
 the heterocyclyl moieties optionally may be substituted with one or more
 substituents independently selected from R⁷⁰; or
 aryl, aryloxy, aryl-C₁₋₆-alkoxy-, aryl-C₁₋₆-alkylene-, heteroaryl,
 heteroaryl-C₁₋₆-alkylene-, of which the aryl and heteroaryl moieties optionally may be
 substituted with one or more substituents selected from halogen, -C(O)OR⁵⁴, -CN,
 -CF₃, -OCF₃, -NO₂, -OR⁵⁴, -NR⁵⁴R⁵⁵ or C₁₋₆-alkyl, wherein
 R⁵⁴, R⁵⁵ and R⁵⁶ are as defined in claim 1.

208. A compound according to claim 207, wherein

R^{43} , R^{44} , and R^{45} independently of each other are selected from
 -OR⁵⁴, -NR⁵⁴R⁵⁵, -SR⁵⁴, -S(O)R⁵⁴, -S(O)₂R⁵⁴, -C(O)NR⁵⁴R⁵⁵, halogen,
 -S-C₁₋₆-alkylene-OR⁵⁴, -C₁₋₆-alkylene-S-R⁵⁴, -C₁₋₆-alkylene-S(O)R⁵⁴,
 -C₁₋₆-alkylene-S(O)₂R⁵⁴, -C₁₋₆-alkylene-N(R⁵⁴)S(O)₂R⁵⁵, -C₁₋₆-alkylene-C(O)NR⁵⁴R⁵⁵,
 -C₁₋₆-alkylene-N(R⁵⁴)C(O)R⁵⁵, -C₁₋₆-alkylene-N(R⁵⁴)C(O)NR⁵⁵R⁵⁶,
 C₁₋₆-alkylene-NHC(=NR⁵⁴)NR⁵⁵R⁵⁶, -C₁₋₆-alkylene-N(R⁵⁴)C(O)OR⁵⁵,
 -C₁₋₆-alkylene-C(O)OR⁵⁴, -C₁₋₆-alkylene-O-R⁵⁴, -C₁₋₆-alkylene-C(O)R⁵⁴,

$-\text{C}_{1-6}\text{-alkylene-NR}^{54}\text{R}^{55}$, $-\text{C}_{1-6}\text{-alkylene=N-O-R}^{54}$, $-\text{C}_{1-6}\text{-alkylene-N=C(N(R}^{54}\text{R}^{55}))_2$,
 $-\text{N=C(N(R}^{54}\text{R}^{55}))_2$, $-\text{C(O)R}^{54}$ and $-\text{C(O)OR}^{54}$; or

$\text{C}_{1-6}\text{-alkyl}$ optionally substituted with one or more substituents independently
 selected from halogen, R^{54} , $-\text{CN}$, $-\text{CF}_3$, $-\text{OCF}_3$, $-\text{OR}^{54}$, $-\text{C(O)OR}^{54}$, $-\text{NR}^{54}\text{R}^{55}$ and

5 $\text{C}_{1-6}\text{-alkyl}$; or

Heterocyclyl or heterocyclyl- $\text{C}_{1-6}\text{-alkylene-}$, of which the heterocyclyl moieties
 optionally may be substituted with one or more substituents independently selected
 from R^{70} ; or

10 aryl, aryl- $\text{C}_{1-6}\text{-alkylene-}$, heteroaryl, heteroaryl- $\text{C}_{1-6}\text{-alkylene-}$, of which the aryl and
 heteroaryl moieties optionally may be substituted with one or more substituents
 selected from halogen, $-\text{C(O)OR}^{54}$, $-\text{CN}$, $-\text{CF}_3$, $-\text{OCF}_3$, $-\text{NO}_2$, $-\text{OR}^{54}$, $-\text{NR}^{54}\text{R}^{55}$ or
 $\text{C}_{1-6}\text{-alkyl}$, wherein

R^{54} , R^{55} , R^{56} , and R^{70} are as defined in claim 1.

209. A compound according to claim 208, wherein

15 R^{43} , R^{44} , and R^{45} independently of each other are selected from

$-\text{SR}^{54}$, $-\text{S(O)R}^{54}$, $-\text{S(O)}_2\text{R}^{54}$, halogen, $-\text{C}_{1-6}\text{-alkylene-S-R}^{54}$, $-\text{C}_{1-6}\text{-alkylene-S(O)R}^{54}$,
 $-\text{C}_{1-6}\text{-alkylene-S(O)}_2\text{R}^{54}$, $-\text{C}_{1-6}\text{-alkylene-N(R}^{54})\text{C(O)OR}^{55}$, $-\text{C}_{1-6}\text{-alkylene-C(O)OR}^{54}$,
 $-\text{C}_{1-6}\text{-alkylene-O-R}^{54}$, $-\text{C}_{1-6}\text{-alkylene-NR}^{54}\text{R}^{55}$, $-\text{C}_{1-6}\text{-alkylene-C(O)R}^{54}$, $-\text{C(O)R}^{54}$ and
 $-\text{C(O)OR}^{54}$; or

20 $\text{C}_{1-6}\text{-alkyl}$ optionally substituted with one or more substituents independently
 selected from halogen, R^{54} , $-\text{CN}$, $-\text{CF}_3$, $-\text{OCF}_3$, $-\text{OR}^{54}$, $-\text{C(O)OR}^{54}$, $-\text{NR}^{54}\text{R}^{55}$ and
 $\text{C}_{1-6}\text{-alkyl}$; or

Heterocyclyl or heterocyclyl- $\text{C}_{1-6}\text{-alkylene-}$, of which the heterocyclyl moieties
 optionally may be substituted with one or more substituents independently selected
 25 from R^{70} ; or

heteroaryl or heteroaryl- $\text{C}_{1-6}\text{-alkylene-}$, of which the aryl and heteroaryl moieties
 optionally may be substituted with one or more substituents selected from halogen,
 $-\text{C(O)OR}^{54}$, $-\text{CN}$, $-\text{CF}_3$, $-\text{OCF}_3$, $-\text{NO}_2$, $-\text{OR}^{54}$, $-\text{NR}^{54}\text{R}^{55}$ or $\text{C}_{1-6}\text{-alkyl}$, wherein
 R^{54} , R^{55} , R^{56} , and R^{70} are as defined in claim 1.

30 210. A compound according to claim 209, wherein

R^{43} , R^{44} , and R^{45} independently of each other are selected from
 $-\text{SR}^{54}$, $-\text{S(O)}_2\text{R}^{54}$, halogen, $-\text{C}_{1-6}\text{-alkylene-C(O)OR}^{54}$, and $-\text{C(O)OR}^{54}$; or

Heterocyclyl or heterocyclyl-C₁₋₆-alkylene-, of which the heterocyclyl moieties optionally may be substituted with one or more substituents independently selected from R⁷⁰; or

- 5 heteroaryl or heteroaryl-C₁₋₆-alkylene-, of which the heteroaryl moieties optionally may be substituted with one or more substituents selected from halogen, -C(O)OR⁵⁴, -CN, -CF₃, -OCF₃, -NO₂, -OR⁵⁴, -NR⁵⁴R⁵⁵ or C₁₋₆-alkyl, wherein R⁵⁴, R⁵⁵, R⁵⁶, and R⁷⁰ are as defined in claim 1.

211. A compound according to claim 210, wherein

- 10 R⁴³, R⁴⁴, and R⁴⁵ independently of each other are selected from -SR⁵⁴, -S(O)₂R⁵⁴, halogen, -C₁₋₆-alkylene-C(O)OR⁵⁴, and -C(O)OR⁵⁴; or heterocyclyl-C₁₋₆-alkylene-, wherein heterocyclyl is selected from imidazolyl, piperidyl, piperazinyl, and morpholinyl, and of which the heterocyclyl moieties optionally may be substituted with one or more substituents independently selected from R⁷⁰; or
- 15 heteroaryl or heteroaryl-C₁₋₆-alkylene-, wherein heteroaryl is selected from thiazolyl, triazolyl, or tetrazolyl, and of which the heteroaryl moieties optionally may be substituted with one or more substituents selected from halogen, -C(O)OR⁵⁴, -CN, -CF₃, -OCF₃, -NO₂, -OR⁵⁴, -NR⁵⁴R⁵⁵ or C₁₋₆-alkyl, wherein R⁵⁴, R⁵⁵, R⁵⁶, and R⁷⁰ are as defined in claim 1.

20 212. A compound according to claim 209, wherein

R⁴³, R⁴⁴, and R⁴⁵ independently of each other are selected from -C₁₋₆-alkylene-S-R⁵⁴, -C₁₋₆-alkylene-O-R⁵⁴, -C₁₋₆-alkylene-S(O)₂R⁵⁴ or -C₁₋₆-alkylene-NR⁵⁴R⁵⁵, wherein R⁵⁴ and R⁵⁵ are as defined in claim 1.

25 213. A compound according to claim 209, wherein

R⁴³ is -C₁₋₆-alkylene-S-R⁵⁴, wherein R⁵⁴ is as defined in claim 1.

214. R⁴³ is -C₁₋₆-alkylene-O-R⁵⁴, wherein R⁵⁴ is as defined in claim 1.

30 215. R⁴³ is -C₁₋₆-alkylene-NR⁵⁴R⁵⁵, wherein R⁵⁴ and R⁵⁵ are as defined in claim 1.

216. R⁴³ is -C₁₋₆-alkylene-S(O)₂R⁵⁴, wherein R⁵⁴ is as defined in claim 1.

217. A compound according to any one of the claims 207 to 210 wherein R^{43} is heteroaryl or heteroaryl- C_{1-6} -alkylene- optionally substituted with one or more substituents selected from halogen, $-C(O)OR^{54}$, $-CN$, $-CF_3$, $-OCF_3$, $-NO_2$, $-OR^{54}$, $-NR^{54}R^{55}$ or C_{1-6} -alkyl, wherein R^{54} and R^{55} are as defined in claim 1.
- 5 218. A compound according to claim 217 wherein R^{43} is heteroaryl optionally substituted with one or more substituents selected from halogen, $-C(O)OR^{54}$, $-CN$, $-CF_3$, $-OCF_3$, $-NO_2$, $-OR^{54}$, $-NR^{54}R^{55}$ or C_{1-6} -alkyl, wherein R^{54} and R^{55} are as defined in claim 1.
219. A compound according to claim 217 wherein R^{43} is heteroaryl- C_{1-6} -alkylene- optionally substituted with one or more substituents selected from halogen, $-C(O)OR^{54}$, $-CN$, $-CF_3$,
 10 $-OCF_3$, $-NO_2$, $-OR^{54}$, $-NR^{54}R^{55}$ or C_{1-6} -alkyl, wherein R^{54} and R^{55} are as defined in claim 1.
220. A compound according to any one of the claims 207 to 211 wherein R^{70} is $=O$, methyl or $C(O)OR^{75}$.
221. A compound according to any one of the claims 207 to 211 wherein R^{70} is $-C(O)OH$
222. A compound according to any one of the claims 207 to 211 wherein R^{70} is
 15 $-(CH_2)_{1-3}C(O)OH$.
223. A compound according to any one of the claims 207 to 211 wherein R^{70} is $-S(O)_2CH_3$.
224. A compound according to claim 211 wherein R^{43} is $-C_{1-6}$ -alkylene- $C(O)OR^{54}$.
225. A compound according to claim 224 wherein R^{43} is $-CH_2-C(O)OR^{54}$.
226. A compound according to claim 211 wherein R^{44} is $-SR^{54}$.
- 20 227. A compound according to claim 211 wherein R^{44} is $-S(O)_2R^{54}$.
228. A compound according to claim 211 wherein R^{44} is halogen.
229. A compound according to any one of the claims 1 to 228 wherein
 R^{54} , R^{55} and R^{56} independently of each other are hydrogen, C_{1-6} -alkyl,
 aryl- C_{1-6} -alkylene-, heteroaryl- C_{1-6} -alkylene-, heterocyclyl,
 25 heterocyclyl- C_{1-6} -alkylene-, heteroaryl, or aryl, each of which is optionally substituted
 with one or more substituents independently selected from R^{71} ;
 or
 R^{54} and R^{55} independently of each other are hydrogen or
 $-(CHR^{72})_u-(CHR^{73})_v-W^6$,
 30 or
 R^{54} and R^{55} , when attached to the same nitrogen atom, together with the said
 nitrogen atom may form a 3 to 8 membered heterocyclic ring optionally containing
 one or two further heteroatoms selected from nitrogen, oxygen and sulfur, and
 optionally containing one or two double bonds, and optionally substituted with one or
 35 more C_{1-6} -alkyl groups,

wherein R^{71} , R^{72} , R^{73} , u , v , and W^6 are as defined in claim 1.

230. A compound according to claim 229 wherein

R^{54} , R^{55} and R^{56} independently of each other are hydrogen, C_{1-6} -alkyl, heterocyclyl, heterocyclyl- C_{1-6} -alkylene-, heteroaryl, or aryl, each of which is optionally substituted with one or more substituents independently selected from R^{71} ;

or

R^{54} and R^{55} independently of each other are hydrogen or $-(CHR^{72})_u-(CHR^{73})_v-W^6$,

wherein R^{71} , R^{72} , R^{73} , u , v , and W^6 are as defined in claim 1.

231. A compound according to any one of the claims 1 to 230 wherein

R^{54} , R^{55} and R^{56} independently of each other are hydrogen, methyl, ethyl, propyl, butyl, isopropyl, isobutyl, sec-butyl, *tert*-butyl, 3-pentyl, 2-pentyl, or 3-methyl-butyl.

232. A compound according to claim 231 wherein

R^{54} , R^{55} and R^{56} independently of each other are hydrogen, methyl, ethyl, propyl, butyl, isopropyl, isobutyl, sec-butyl, or *tert*-butyl.

233. A compound according to claim 232 wherein

R^{54} , R^{55} and R^{56} independently of each other are hydrogen, methyl, ethyl, propyl, butyl, isopropyl, isobutyl, sec-butyl, or *tert*-butyl.

234. A compound according to claim 233 wherein

R^{54} , R^{55} and R^{56} independently of each other are hydrogen, methyl or ethyl.

235. A compound according to claim 234 wherein

R^{54} is hydrogen.

236. A compound according to any one of the claims 1 to 230 wherein

R^{54} , R^{55} and R^{56} independently of each other are oxetanyl, tetrahydrofuranyl, tetrahydropyranyl, oxepanyl, azetidyl, pyrrolidyl, piperidyl, hexahydroazepinyl, thietanyl, thiolanyl, tetrahydrothiopyranyl, thiepanyl, 1,4-oxathianyl, 1,3-dioxolanyl, 1,2-dithiolanyl, 1,3-dithiolanyl, hexahydro-pyridazinyl, imidazolidyl, 1,3-dioxanyl, morpholinyl, 1,3-dithianyl, 1,4-dioxanyl, 1,4-dithianyl, or thiomorpholinyl, each of which is optionally substituted with one or more substituents independently selected from R^{71} , wherein R^{71} is as defined in claim 1.

237. A compound according to claim 236 wherein

R^{54} , R^{55} and R^{56} independently of each other are tetrahydropyranyl, oxepanyl, piperidyl, hexahydroazepinyl, tetrahydrothiopyranyl, thiepanyl, 1,4-oxathianyl, 1,3-dithiolanyl, hexahydro-pyridazinyl, 1,3-dioxanyl, morpholinyl, 1,3-dithianyl, 1,4-dioxanyl, 1,4-dithianyl, or thiomorpholinyl, each of which is optionally substituted

with one or more substituents independently selected from R^{71} , wherein R^{71} is as defined in claim 1.

238. A compound according to claim 237 wherein

R^{54} , R^{55} and R^{56} independently of each other are tetrahydropyranyl, piperidyl,

5 tetrahydrothiopyranyl, 1,4-oxathianyl, hexahydro-pyridazinyl, morpholinyl, 1,4-dioxanyl, 1,4-dithianyl, or thiomorpholinyl, each of which is optionally substituted with one or more substituents independently selected from R^{71} , wherein R^{71} is as defined in claim 1.

239. A compound according to claim 238 wherein

10 R^{54} , R^{55} and R^{56} independently of each other are tetrahydropyranyl, piperidyl, tetrahydrothiopyranyl, or morpholinyl, each of which is optionally substituted with one or more substituents independently selected from R^{71} , wherein R^{71} is as defined in claim 1.

240. A compound according to claim 239 wherein

15 R^{54} , R^{55} and R^{56} independently of each other are piperidyl or morpholinyl, each of which is optionally substituted with one or more substituents independently selected from R^{71} , wherein R^{71} is as defined in claim 1.

241. A compound according to claim 240 wherein

R^{54} , R^{55} and R^{56} independently of each other are piperidyl or morpholinyl.

242.

20 243. A compound according to any one of the claims 1 to 230 wherein

R^{54} , R^{55} and R^{56} independently of each other are furanyl, thienyl, thiophenyl, pyrrolyl, imidazolyl, pyrazolyl, triazolyl, tetrazolyl, thiazolyl, oxazolyl, isoxazolyl, oxadiazolyl, thiadiazolyl, isothiazolyl, pyridinyl, pyridazinyl, pyrazinyl, pyrimidinyl, quinolinyl, isoquinolinyl, benzofuranyl, benzothiophenyl, indolyl, purinyl, or indazolyl, each of
25 which is optionally substituted with one or more substituents independently selected from R^{71} , wherein R^{71} is as defined in claim 1.

244. A compound according to claim 243 wherein

R^{54} , R^{55} and R^{56} independently of each other are furanyl, thienyl, thiophenyl, pyrrolyl, imidazolyl, pyrazolyl, triazolyl, tetrazolyl, thiazolyl, oxazolyl, isoxazolyl, oxadiazolyl, thiadiazolyl, isothiazolyl, pyridinyl, pyridazinyl, pyrazinyl, pyrimidinyl, benzofuranyl, indolyl, or purinyl, each of which is optionally substituted with one or more
30 substituents independently selected from R^{71} , wherein R^{71} is as defined in claim 1.

245. A compound according to claim 244 wherein

35 R^{54} , R^{55} and R^{56} independently of each other are imidazolyl, triazolyl, tetrazolyl, thiazolyl, pyridinyl, pyrimidinyl, benzofuranyl, indolyl, or purinyl, each of which is

optionally substituted with one or more substituents independently selected from R^{71} , wherein R^{71} is as defined in claim 1.

246. A compound according to claim 245 wherein

5 R^{54} , R^{55} and R^{56} independently of each other are imidazolyl, triazolyl, tetrazolyl, thiazolyl, pyridinyl, pyrimidinyl, or purinyl, each of which is optionally substituted with one or more substituents independently selected from R^{71} , wherein R^{71} is as defined in claim 1.

247. A compound according to claim 246 wherein

10 R^{54} , R^{55} and R^{56} independently of each other are imidazolyl, triazolyl, tetrazolyl, thiazolyl, pyridinyl, pyrimidinyl, each of which is optionally substituted with one or more substituents independently selected from R^{71} , wherein R^{71} is as defined in claim 1.

248. A compound according to claim 247 wherein R^{54} is imidazolyl optionally substituted with one or more substituents independently selected from R^{71} , wherein R^{71} is as defined in claim 1.

249. A compound according to claim 247 wherein R^{54} is triazolyl optionally substituted with one or more substituents independently selected from R^{71} , wherein R^{71} is as defined in claim 1.

20 250. A compound according to claim 247 wherein R^{54} is tetrazolyl optionally substituted with one or more substituents independently selected from R^{71} , wherein R^{71} is as defined in claim 1.

251. A compound according to claim 247 wherein R^{54} is thiazolyl optionally substituted with one or more substituents independently selected from R^{71} , wherein R^{71} is as defined in claim 1.

25 252. A compound according to claim 247 wherein R^{54} is pyridinyl optionally substituted with one or more substituents independently selected from R^{71} , wherein R^{71} is as defined in claim 1.

30 253. A compound according to claim 247 wherein R^{54} is pyrimidinyl optionally substituted with one or more substituents independently selected from R^{71} , wherein R^{71} is as defined in claim 1.

254. A compound according to claim 246 wherein R^{54} is purinyl optionally substituted with one or more substituents independently selected from R^{71} , wherein R^{71} is as defined in claim 1.

255. A compound according to any one of the claims 1 to 254 wherein R^{71} is methyl or =O.

35 256. A compound according to any one of the claims 1 to 254 wherein R^{71} is $-C(O)OH$

257. A compound according to any one of the claims 1 to 254 wherein R^{71} is $-(CH_2)_{1-3}C(O)OH$.
258. A compound according to any one of the claims 1 to 254 wherein R^{70} is $-(CH_2)_{1-3}NR^{75}R^{76}$.
- 5 259. A compound according to any one of the claims 1 to 230 wherein u is 0 or 1.
260. A compound according to claim 259 wherein u is 0.
261. A compound according to claim 259 wherein u is 1.
262. A compound according to any one of the claims 1 to 230 wherein v is 0 or 1.
263. A compound according to claim 262 wherein v is 0.
- 10 264. A compound according to claim 262 wherein v is 1.
265. A compound according to any one of the claims 1 to 230 wherein u is 0 and v is 1.
266. A compound according to any one of the claims 1 to 230 wherein u and v are both 0.
267. A compound according to any one of the claims 1 to 230 or 259 to 266 wherein R^{72} and R^{73} are independently selected from hydrogen, hydroxy or $-C(O)OR^{75}$.
- 15 268. A compound according to claim 267 wherein R^{72} and R^{73} are independently selected from hydrogen or $-C(O)OR^{75}$, wherein R^{75} is as defined in claim 1.
269. A compound according to claim 267 wherein R^{72} and R^{73} are hydrogen.
270. A compound according to any one of the claims 1 to 230 or 259 to 269 wherein
 W^6 is $-O-R^{75}$, $-C(O)O-R^{75}$, $-C(O)-R^{75}$, $-NR^{75}R^{76}$, $-NHCH_2C(O)R^{75}$,
 20 $-NHC(O)R^{75}$, $-S(O)_2R^{75}$, $-NHS(O)_2R^{75}$, or
 W^6 is heterocyclyl, wherein R^{75} and R^{76} are as defined in claim 1.
271. A compound according to claim 270 wherein
 W^6 is $-O-R^{75}$, $-C(O)O-R^{75}$, $-C(O)-R^{75}$, $-NR^{75}R^{76}$, $-NHCH_2C(O)R^{75}$,
 $-NHC(O)R^{75}$, $-S(O)_2R^{75}$, $-NHS(O)_2R^{75}$, or
 25 W^6 is oxetanyl, tetrahydrofuranyl, tetrahydropyranyl, oxepanyl, azetidyl,
 pyrrolidyl, piperidyl, hexahydroazepinyl, thietanyl, thiolanyl,
 tetrahydrothiopyranyl, thiepanyl, 1,4-oxathianyl, 1,3-dioxolanyl, 1,2-
 dithiolanyl, 1,3-dithiolanyl, hexahydro-pyridazinyl, imidazolidyl, 1,3-dioxanyl,
 morpholinyl, 1,3-dithianyl, 1,4-dioxanyl, 1,4-dithianyl, or thiomorpholinyl,
 30 wherein R^{75} and R^{76} are as defined in claim 1.
272. A compound according to claim 270 wherein
 W^6 is $-O-R^{75}$, $-C(O)O-R^{75}$, $-C(O)-R^{75}$, $-NR^{75}R^{76}$, $-NHCH_2C(O)R^{75}$,
 $-NHC(O)R^{75}$, $-S(O)_2R^{75}$, $-NHS(O)_2R^{75}$, or
 W^6 is tetrahydropyranyl, oxepanyl, piperidyl, hexahydroazepinyl,
 35 tetrahydrothiopyranyl, thiepanyl, 1,4-oxathianyl, morpholinyl, 1,4-dioxanyl,

1,4-dithianyl, or thiomorpholinyl, wherein R^{75} and R^{76} are as defined in claim 1.

273. A compound according to claim 272 wherein

5 W^6 is $-O-R^{75}$, $-C(O)O-R^{75}$, $-C(O)-R^{75}$, $-NR^{75}R^{76}$, $-NHCH_2C(O)R^{75}$,
 $-NHC(O)R^{75}$, $-S(O)_2R^{75}$, $-NHS(O)_2R^{75}$, or
 W^6 is tetrahydropyranyl, piperidyl, tetrahydrothiopyranyl, or morpholinyl,
wherein R^{75} and R^{76} are as defined in claim 1.

274. A compound according to claim 273 wherein

10 W^6 is $-O-R^{75}$, $-C(O)O-R^{75}$, $-NR^{75}R^{76}$, $-NHC(O)R^{75}$, $-S(O)_2R^{75}$, or
 W^6 is tetrahydropyranyl, piperidyl, tetrahydrothiopyranyl, or morpholinyl,
wherein R^{75} and R^{76} are as defined in claim 1.

275. A compound according to claim 274 wherein

W^6 is $-O-R^{75}$, or $-C(O)O-R^{75}$, wherein R^{75} is as defined in claim 1.

15 276. A compound according to claim 275 wherein W^6 is $-C(O)O-R^{75}$, wherein R^{75} is as
defined in claim 1..

277. A compound according to any one of the claims 1 to 230 or 259 to 276 wherein R^{75} and
 R^{76} are independently selected from hydrogen, $-OH$, or C_{1-8} -alkyl optionally substituted with
 $-NH_2$.

20 278. A compound according to claim 277 wherein R^{75} and R^{76} are independently selected
from hydrogen, $-OH$, or methyl.

279. A compound according to claim 278 wherein R^{75} and R^{76} are independently selected
from hydrogen or $-OH$.

280. A compound according to claim 279 wherein R^{75} is hydrogen.

25 281. A compound according to any one of the claims 1 to 280, which compound is an
activator of glucokinase, when tested in the Glucokinase Activation Assay (I) disclosed
herein at a glucose concentration of 2 mM.

282. A compound according to any one of the claims 1 to 281, which compound is an
activator of glucokinase, when tested in the Glucokinase Activation Assay (I) disclosed
herein at a glucose concentration of from 10 to 15 mM.

30 283. A compound according to any one of the claims 1 to 282, which compound, at a
concentration of 30 μM , is capable of providing an at least 1.5, such as at least 1.7, for

instance at least 2.0 fold activation of glucokinase in the Glucokinase Activation Assay (I) disclosed herein at a glucose concentration of 2 mM.

284. A compound according to any one of the claims 1 to 283, which compound, at a concentration of 30 μ M, is capable of providing an at least 1.5, such as at least 1.7, for instance at least 2.0 fold activation of glucokinase in the Glucokinase Activation Assay (I) disclosed herein at a glucose concentration of from 10 to 15 mM.

285. A compound according to any one of the claims 1 to 284, which at a concentration of 5 μ M is capable of providing an at least 1.5, such as at least 1.7, for instance at least 2.0 fold activation of glucokinase in the Glucokinase Activation Assay (I) disclosed herein at a glucose concentration of 2 mM.

286. A compound according to any one of the claims 1 to 285, which at a concentration of 5 μ M is capable of providing an at least 1.5, such as at least 1.7, for instance at least 2.0 fold activation of glucokinase in the Glucokinase Activation Assay (I) disclosed herein at a glucose concentration of from 10 to 15 mM.

287. A glucose kinase activator compound defined as a compound which at a compound concentration of at or below 30 μ M at a glucose concentration of 2 mM in the Glucokinase Activation Assay (I) disclosed herein gives 1.5 - fold higher glucokinase activity than measured at a glucose concentration of 2 mM in the Glucokinase Activation Assay (I) without compound, where the increase in glucokinase activity provided by the compound increases with increasing concentrations of glucose.

288. A compound according to any one of the claims 1 to 286, which compound provides an increase in glucokinase activity, where the increase in glucokinase activity provided by the compound increases with increasing concentrations of glucose.

289. A compound according to claim 287 or claim 288, which provides an increase in glucokinase activity in Glucokinase Activation Assay (I) disclosed herein at a glucose concentration of 15 mM, which increase is significantly higher than the increase in glucokinase activity provided by the compound in Glucokinase Activation Assay (I) disclosed herein at a glucose concentration of 5 mM.

290. A compound according to any one of the claims 287 to 289, which at a compound concentration of 10 μ M provides an increase in glucokinase activity in Glucokinase Activation Assay (I) disclosed herein at a glucose concentration of 15 mM, which increase is significantly higher than the increase in glucokinase activity provided by the compound at a compound concentration of 10 μ M in Glucokinase Activation Assay (I) disclosed herein at a glucose concentration of 5 mM.

291. A compound according to any one of the claims 287 to 290, which at a compound concentration of 10 μ M provides an increase in glucokinase activity in Glucokinase Activation Assay (I) disclosed herein at a glucose concentration of 15 mM, which increase is at least 1.1 fold higher, such as at least 1.2 fold higher, for instance at least 1.3 fold higher, such as at least 1.4 fold higher, for instance 1.5 fold higher, such as at least 1.6 fold higher, for instance at least 1.7 fold higher, such as at least 1.8 fold higher, for instance at least 1.9 fold higher, such as at least 2.0 fold higher than the increase in glucokinase activity provided by the compound at a compound concentration of 10 μ M in Glucokinase Activation Assay (I) disclosed herein at a glucose concentration of 5 mM.

292. A glucose kinase activator compound defined as a compound which at a compound concentration of at or below 30 μ M at a glucose concentration of 2 mM in the Glucokinase Activation Assay (I) disclosed herein gives 1.5 - fold higher glucokinase activity than measured at a glucose concentration of 2 mM in the Glucokinase Activation Assay (I) without compound, which glucose kinase activator compound increases glucose utilization in the liver without inducing any increase in insulin secretion in response to glucose.

293. A glucose kinase activator compound defined as a compound which at a compound concentration of at or below 30 μ M at a glucose concentration of 2 mM in the Glucokinase Activation Assay (I) disclosed herein gives 1.5 - fold higher glucokinase activity than measured at a glucose concentration of 2 mM in the Glucokinase Activation Assay (I) without compound, which glucokinase activator compound shows a significantly higher activity in isolated hepatocytes compared to the activity of the glucokinase compound in Ins-1 cells.

294. A compound according to any one of the claims 1 to 291, which compound increases glucose utilization in the liver without inducing any increase in insulin secretion in response to glucose.

295. A compound according to any one of the claims 1 to 291, which compound shows a significantly higher activity in isolated hepatocytes compared to the activity of the compound in Ins-1 cells.

5 296. A compound according to any one of the claims 292 to 295, which compound shows a significantly higher activity in isolated hepatocytes measured as described in the Glucokinase Activity Assay (II) compared to the activity of the compound in Ins-1 cells measured as described in the Glucokinase Activity Assay (III).

10 297. A compound according to claim 296, which compound shows an activity in isolated hepatocytes measured as described in the Glucokinase Activity Assay (II) which activity is at least 1.1 fold higher, such as at least 1.2 fold higher, for instance at least 1.3 fold higher, such as at least 1.4 fold higher, for instance 1.5 fold higher, such as at least 1.6 fold higher, for instance at least 1.7 fold higher, such as at least 1.8 fold higher, for instance at least 1.9 fold higher, such as at least 2.0 fold higher, for instance at least a 3.0 fold higher, such as at least a 4.0 fold higher, for instance at least 5.0 fold higher, such as at least 10 fold higher
15 than the activity of the compound in Ins-1 cells measured as described in the Glucokinase Activity Assay (III).

298. A compound according to claim 296, which compound shows no activity in the Ins-1 cells measured as described in the Glucokinase Activity Assay (III).

20 299. A method of preventing hypoglycaemia comprising administration of a compound according to any one of the claims 292 to 298.

300. The use of a compound according to any one of the claims 292 to 298 for the preparation of a medicament for the prevention of hypoglycaemia.

25 301. A compound according to any one of claims 1 to 298, which is an agent useful for the treatment of an indication selected from the group consisting of hyperglycemia, IGT, insulin resistance syndrome, syndrome X, type 2 diabetes, type 1 diabetes, dyslipidemia, hypertension, and obesity.

302. A compound according to any one of claims 1 to 301 for use as a medicament.

303. A compound according to any one of claims 1 to 301 for treatment of hyperglycemia, for treatment of IGT, for treatment of Syndrome X, for treatment of type 2 diabetes, for treatment

of type 1 diabetes, for treatment of dyslipidemia, for treatment of hyperlipidemia, for treatment of hypertension, for treatment of obesity, for lowering of food intake, for appetite regulation, for regulating feeding behaviour, or for enhancing the secretion of enteroincretins, such as GLP-1.

- 5 304. A pharmaceutical composition comprising, as an active ingredient, at least one compound according to any one of claims 1 to 303 together with one or more pharmaceutically acceptable carriers or excipients.

305. A pharmaceutical composition according to claim 304 in unit dosage form, comprising from about 0.05 mg to about 1000 mg, preferably from about 0.1 mg to about 500 mg and
10 especially preferred from about 0.5 mg to about 200 mg of the compound according to any one of claims 1 to 303.

306. Use of a compound according to any one of the claims 1 to 303 for increasing the activity of glucokinase.

307. Use of a compound according to any one of claims 1 to 303 for the preparation of a
15 medicament for the treatment of metabolic disorders, for blood glucose lowering, for the treatment of hyperglycemia, for the treatment of IGT, for the treatment of Syndrome X, for the treatment of impaired fasting glucose (IFG), for the treatment of type 2 diabetes, for the treatment of type 1 diabetes, for delaying the progression of impaired glucose tolerance (IGT) to type 2 diabetes, for delaying the progression of non-insulin requiring type 2 diabetes
20 to insulin requiring type 2 diabetes, for the treatment of dyslipidemia, for the treatment of hyperlipidemia, for the treatment of hypertension, for lowering of food intake, for appetite regulation, for the treatment of obesity, for regulating feeding behaviour, or for enhancing the secretion of enteroincretins.

308. Use of a compound according to any one of claims 1 to 303 for the preparation of a
25 medicament for the adjuvant treatment of type 1 diabetes for preventing the onset of diabetic complications.

309. Use of a compound according to any one of claims 1 to 303 for the preparation of a medicament for increasing the number and/or the size of beta cells in a mammalian subject,

for treatment of beta cell degeneration, in particular apoptosis of beta cells, or for treatment of functional dyspepsia, in particular irritable bowel syndrome.

310. Use according to any one of the claims 307 to 309 in a regimen which comprises treatment with a further antidiabetic agent.

5 311. Use according to any one of the claims 307 to 310 in a regimen which comprises treatment with a further antihyperlipidemic agent.

312. Use according to any one of claims 307 to 311 in a regimen which comprises treatment with a further antiobesity agent.

10 313. Use according to any one of claims 307 to 312 in a regimen which comprises treatment with a further antihypertensive agent.

15 314. Use of a compound according to any one of the claims 1 to 303 or a pharmaceutical composition according to claim 304 or claim 305 for the treatment of metabolic disorders, for blood glucose lowering, for the treatment of hyperglycemia, for treatment of IGT, for treatment of Syndrome X, for the treatment of impaired fasting glucose (IFG), for treatment of type 2 diabetes, for treatment of type 1 diabetes, for delaying the progression of impaired glucose tolerance (IGT) to type 2 diabetes, for delaying the progression of non-insulin requiring type 2 diabetes to insulin requiring type 2 diabetes, for treatment of dyslipidemia, for treatment of hyperlipidemia, for treatment of hypertension, for the treatment or prophylaxis of obesity, for lowering of food intake, for appetite regulation, for regulating feeding behaviour, or for enhancing the secretion of enteroincretins.

20 315. Use of a compound according to any one of the claims 1 to 303 or a pharmaceutical composition according to claim 304 or claim 305 for the adjuvant treatment of type 1 diabetes for preventing the onset of diabetic complications.

25 316. Use of a compound according to any one of the claims 1 to 303 or a pharmaceutical composition according to claim 304 or claim 305 for increasing the number and/or the size of beta cells in a mammalian subject, for treatment of beta cell degeneration, in particular apoptosis of beta cells, or for treatment of functional dyspepsia, in particular irritable bowel syndrome.

INTERNATIONAL SEARCH REPORT

International Application No

PCT/DK 03/00449

A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 A61K31/433 A61P3/10 A61P3/06 A61P3/04 C07D277/48
 C07D277/52 C07D213/75 C07D417/12 C07D277/56 C07D417/06
 C07D417/14 C07D285/16 C07D487/04 C07D277/54

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 C07D A61K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

EPO-Internal, WPI Data, PAJ, CHEM ABS Data, INSPEC

C. DOCUMENTS CONSIDERED TO BE RELEVANT

| Category * | Citation of document, with indication, where appropriate, of the relevant passages | Relevant to claim No. |
|------------|--|-----------------------|
| X | <p>REGEL E ET AL: "C-ACYLIERUNG VON 5 GLIEDRIGEN N-HETEROCYCLLEN, I ACYLIERUNG AN C-2 VON IMIDAZOLEN UND BENZIMIDAZOLEN C-ACYLATION OF 5-MEMBERED N-HETEROCYCLES, I. - ACYLATION AT C-2 OF IMIDAZOLES AND BENZIMIDAZOLES"</p> <p>JUSTUS LIEBIGS ANNALEN DER CHEMIE, VERLAG CHEMIE GMBH. WEINHEIM, DE, 1977, pages 145-158, XP009003467 ISSN: 0075-4617 see compound 1347 page 154; table 6</p> <p>---</p> | 1 |
| X | <p>US 3 152 136 A (HARRIS GUY H ET AL) 6 October 1964 (1964-10-06) examples</p> <p>---</p> <p>---/---</p> | 1 |

☒ Further documents are listed in the continuation of box C.☒ Patent family members are listed in annex.

* Special categories of cited documents:

- "A" document defining the general state of the art which is not considered to be of particular relevance
- "E" earlier document but published on or after the international filing date
- "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- "O" document referring to an oral disclosure, use, exhibition or other means
- "P" document published prior to the international filing date but later than the priority date claimed

- "T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- "X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- "Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- "&" document member of the same patent family

Date of the actual completion of the international search

5 September 2003

Date of mailing of the international search report

23/09/2003

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
 NL - 2280 HV Rijswijk
 Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
 Fax: (+31-70) 340-3016

Authorized officer

Schmid, J-C

INTERNATIONAL SEARCH REPORT

Int. Patent Application No.

PCT/DK 03/00449

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

| Category * | Citation of document, with indication, where appropriate, of the relevant passages | Relevant to claim No. |
|------------|--|-----------------------|
| X | US 3 317 534 A (FUMIO YONEDA ET AL) 2 May 1967 (1967-05-02) examples --- | 1 |
| X | DE 21 17 807 A (SHELL) 28 October 1971 (1971-10-28) examples --- | 1 |
| X | WO 97 24328 A (BAYER AG ;ES SAYED MAZEN (DE); FROBEL KLAUS (DE); GRIX SUZANNA (GB) 10 July 1997 (1997-07-10) page 19, line 3 page 32, line 1 page 35, line 1 page 40, line 1 page 47, line 2 page 52, line 2 claims --- | 1,158, 195 |
| X | EP 1 211 246 A (IHARA CHEMICAL IND CO ;KUMIAI CHEMICAL INDUSTRY CO (JP)) 5 June 2002 (2002-06-05) claims; table 29 --- | 1,158, 195 |
| X | PURCHASE C F ET AL: "Tetrazole-substituted ureas as inhibitors of acyl-CoA:cholesterol O-acyltransferase (ACAT). A novel preparation of ureas from weakly nucleophilic amines" BIOORGANIC & MEDICINAL CHEMISTRY LETTERS, OXFORD, GB, vol. 6, no. 15, 6 August 1996 (1996-08-06), pages 1753-1758, XP004135596 ISSN: 0960-894X examples --- | 1,160, 195 |
| X | DE 27 12 630 A (BAYER AG) 28 September 1978 (1978-09-28) examples 17,24,27 --- | 1,160, 195 |
| X | ATWAL ET AL: "Cardioselective Antiischemic ATP-Sensitive Potassium Channel Openers. 4. Structure-Activity Studies on Benzopyranylcyanoguanidines: Replacement of the Benzopyran Portion" JOURNAL OF MEDICINAL CHEMISTRY, AMERICAN CHEMICAL SOCIETY, WASHINGTON, US, vol. 39, no. 1, January 1996 (1996-01), pages 304-313, XP002116120 ISSN: 0022-2623 examples --- | 1,160, 195 |

-/--

INTERNATIONAL SEARCH REPORT

Int. Patent Application No

PCT/DK 03/00449

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

| Category * | Citation of document, with indication, where appropriate, of the relevant passages | Relevant to claim No. |
|------------|---|-----------------------|
| X | WHITE A D ET AL: "HETEROCYCLIC UREAS: INHIBITORS OF ACYL-COA:CHOLESTEROL O-ACYLTRANSFERASE AS HYPOCHOLESTEROLEMIC AGENTS" JOURNAL OF MEDICINAL CHEMISTRY, AMERICAN CHEMICAL SOCIETY. WASHINGTON, US, vol. 39, no. 22, 25 October 1996 (1996-10-25), pages 4382-4395, XP002918005 ISSN: 0022-2623 examples; table 5 ---- | 1,160, 195 |
| X | WO 99 32111 A (BAYER AG) 1 July 1999 (1999-07-01) examples 2,110,136; table 1 ---- | 1,160, 195 |
| X | WO 99 32106 A (BAYER AG) 1 July 1999 (1999-07-01) examples 92,127,128,159,160,351 ---- | 1,160, 195 |
| X | WO 01 57008 A (BASF AG ;ERICSSON ANNA (US); SCOTT BARBARA (US); ARNOLD LEE D (US)) 9 August 2001 (2001-08-09) examples 118,146 ---- | 1,160, 195 |
| X | WO 99 24035 A (SQUIBB BRISTOL MYERS CO) 20 May 1999 (1999-05-20) examples 64,68-73,84-88,93-95,109,110 ---- | 1,160, 195 |
| X | US 5 556 969 A (CHAMBERS MARK S ET AL) 17 September 1996 (1996-09-17) claims 1,6 ---- | 1,160, 195 |
| X | WO 02 14311 A (AMGEN INC) 21 February 2002 (2002-02-21) examples ---- | 1,160, 195 |
| E | WO 03 070727 A (AMGEN INC) 28 August 2003 (2003-08-28) examples ---- | 1,160, 195 |
| L | WO 03 055482 A (NOVO NORDISK A/S) 10 July 2003 (2003-07-10) examples 131,133 ---- | 1-316 |
| P,X | WO 02 070494 A (ICOS CORP) 12 September 2002 (2002-09-12) examples ---- | 1,160, 195 |
| A | WO 02 46173 A (HOFFMANN LA ROCHE) 13 June 2002 (2002-06-13) the whole document ----- | 1-316 |

INTERNATIONAL SEARCH REPORT

International application No.
PCT/DK 03/00449

Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)

This International Search Report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☒ Claims Nos.:
because they relate to subject matter not required to be searched by this Authority, namely:
Although claims 314 to 316 are directed to a method of treatment of the human/animal body, the search has been carried out and based on the alleged effects of the compound/composition.
2. ☒ Claims Nos.:
because they relate to parts of the International Application that do not comply with the prescribed requirements to such an extent that no meaningful International Search can be carried out, specifically:
see FURTHER INFORMATION sheet PCT/ISA/210
3. ☐ Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this International Search Report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this International Search Report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this International Search Report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
- ☐ No protest accompanied the payment of additional search fees.

FURTHER INFORMATION CONTINUED FROM PCT/ISA/ 210

Continuation of Box I.2

The initial phase of the search revealed a very large number of documents relevant to the issue of novelty. So many documents were retrieved that it is impossible to determine which parts of the claim(s) may be said to define subject-matter for which protection might legitimately be sought (Article 6 PCT). For these reasons, a meaningful search over the whole breadth of the claim(s) is impossible. Consequently, the search has been restricted to the compounds of formula 1 wherein L1 is -CH₂-, -O-, -S-, -SO-, SO₂-, -NR₁₁-, C(O), -C(=N-OR₁₂)-, G1 is aryl or heteroaryl, L2 is -N(R₂₀)-, L3 is -C(O)- and G3 is heteroaryl, fused heterocycl(heteroaryl or fused cycloalkylheteroalkyl).

The applicant's attention is drawn to the fact that claims, or parts of claims, relating to inventions in respect of which no international search report has been established need not be the subject of an international preliminary examination (Rule 66.1(e) PCT). The applicant is advised that the EPO policy when acting as an International Preliminary Examining Authority is normally not to carry out a preliminary examination on matter which has not been searched. This is the case irrespective of whether or not the claims are amended following receipt of the search report or during any Chapter II procedure.

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/DK 03/00449

| Patent document cited in search report | Publication date | Patent family member(s) | Publication date |
|---|---------------------|----------------------------|--|
| US 3152136 | A | 06-10-1964 | NONE |
| US 3317534 | A | 02-05-1967 | NONE |
| DE 2117807 | A | 28-10-1971 | CA 946845 A1 07-05-1974 CH 552339 A 15-08-1974 DE 2117807 A1 28-10-1971 ES 390121 A1 01-05-1974 FR 2089546 A5 07-01-1972 GB 1318291 A 23-05-1973 JP 54007857 B 10-04-1979 NL 7104858 A 19-10-1971 PH 9683 A 10-02-1976 US 3736330 A 29-05-1973 ZA 7102295 A 26-01-1972 |
| WO 9724328 | A | 10-07-1997 | AU 1372897 A 28-07-1997 WO 9724328 A1 10-07-1997 |
| EP 1211246 | A | 05-06-2002 | AU 6876800 A 10-04-2001 CA 2384354 A1 15-03-2001 EP 1211246 A1 05-06-2002 CN 1372552 T 02-10-2002 WO 0117975 A1 15-03-2001 |
| DE 2712630 | A | 28-09-1978 | DE 2712630 A1 28-09-1978 BE 867173 A1 18-09-1978 BR 7801766 A 19-12-1978 DK 132578 A 24-09-1978 FR 2384763 A1 20-10-1978 JP 53119875 A 19-10-1978 NL 7803046 A 26-09-1978 US 4174398 A 13-11-1979 |
| WO 9932111 | A | 01-07-1999 | AU 739642 B2 18-10-2001 AU 1997199 A 12-07-1999 CA 2315720 A1 01-07-1999 DE 1041982 T1 07-06-2001 EP 1041982 A1 11-10-2000 ES 2154253 T1 01-04-2001 JP 2001526223 T 18-12-2001 WO 9932111 A1 01-07-1999 |
| WO 9932106 | A | 01-07-1999 | AU 2198999 A 12-07-1999 BG 104597 A 28-02-2001 BR 9814374 A 14-05-2002 CA 2315717 A1 01-07-1999 CN 1290164 T 04-04-2001 CZ 20002350 A3 15-08-2001 DE 1047418 T1 03-05-2001 EP 1047418 A1 02-11-2000 ES 2153340 T1 01-03-2001 GR 2001300007 T1 28-02-2001 HU 0101704 A2 28-12-2001 JP 2001526220 T 18-12-2001 NO 20003232 A 21-08-2000 PL 343083 A1 30-07-2001 SK 9632000 A3 12-03-2001 |

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/DK 03/00449

| Patent document cited in search report | Publication date | Patent family member(s) | Publication date |
|---|---------------------|---|--|
| WO 9932106 A | | TR 200002618 T2 WO 9932106 A1 | 20-04-2001 01-07-1999 |
| WO 0157008 A | 09-08-2001 | AU 3669801 A BG 107062 A BR 0108085 A CA 2398754 A1 CN 1422262 T EP 1254123 A1 HU 0300359 A2 JP 2003521543 T NO 20023713 A SK 12712002 A3 WO 0157008 A1 US 2003153568 A1 | 14-08-2001 30-04-2003 18-03-2003 09-08-2001 04-06-2003 06-11-2002 28-06-2003 15-07-2003 04-10-2002 04-02-2003 09-08-2001 14-08-2003 |
| WO 9924035 A | 20-05-1999 | AU 744281 B2 AU 1371999 A BR 9814956 A CA 2309319 A1 CN 1290165 T EP 1037632 A1 HU 0102101 A2 JP 2001522800 T NO 20002121 A NZ 503491 A PL 340727 A1 TR 200001312 T2 TW 510898 B US 2002123484 A1 WO 9924035 A1 ZA 9810219 A | 21-02-2002 31-05-1999 03-10-2000 20-05-1999 04-04-2001 27-09-2000 28-11-2001 20-11-2001 09-05-2000 28-08-2002 26-02-2001 21-09-2000 21-11-2002 05-09-2002 20-05-1999 22-06-2000 |
| US 5556969 A | 17-09-1996 | NONE | |
| WO 0214311 A | 21-02-2002 | AU 8490901 A CA 2418273 A1 EP 1309589 A2 WO 0214311 A2 US 2002193405 A1 US 2002173507 A1 | 25-02-2002 21-02-2002 14-05-2003 21-02-2002 19-12-2002 21-11-2002 |
| WO 03070727 A | 28-08-2003 | US 2002193405 A1 WO 03070727 A1 | 19-12-2002 28-08-2003 |
| WO 03055482 A | 10-07-2003 | EP 1336607 A1 WO 03047626 A1 WO 03055482 A1 US 2003138416 A1 | 20-08-2003 12-06-2003 10-07-2003 24-07-2003 |
| WO 02070494 A | 12-09-2002 | WO 02070494 A1 US 2003069284 A1 | 12-09-2002 10-04-2003 |
| WO 0246173 A | 13-06-2002 | AU 2190202 A CA 2429642 A1 WO 0246173 A1 US 2002111372 A1 US 2002103241 A1 | 18-06-2002 13-06-2002 13-06-2002 15-08-2002 01-08-2002 |

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/DK 03/00449

| Patent document cited in search report | Publication date | Patent family member(s) | Publication date |
|---|---------------------|----------------------------|---------------------|
| WO 0246173 | A | US 2002103199 A1 | 01-08-2002 |
| | | US 2002107396 A1 | 08-08-2002 |
| ----- | | | |